

1996

Disease Management

Carolyn DeMoranville
Cranberry Station, carolynd@umext.umass.edu

Hilary A. Sandler
Cranberry Station, hsandler@umass.edu

Tom Bicki


Follow this and additional works at: <https://scholarworks.umass.edu/cranberrybmp>

 Part of the [Life Sciences Commons](#)

DeMoranville, Carolyn; Sandler, Hilary A.; and Bicki, Tom, "Disease Management" (1996). *Cranberry Station Best Management Practices Guide - 2000 Edition*. 8.

Retrieved from <https://scholarworks.umass.edu/cranberrybmp/8>

This Public Service and Outreach is brought to you for free and open access by the Cranberry Station Outreach and Public Service Activities at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Cranberry Station Best Management Practices Guide - 2000 Edition by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.



BEST MANAGEMENT PRACTICES GUIDE FOR MASSACHUSETTS CRANBERRY PRODUCTION

Disease Management

Applying an integrated scheme for the control of plant diseases can be a viable approach for many cranberry growers. Sanitation, proper irrigation schedules, resistant varieties, fungicides, and various cultural techniques can be used as components of a disease management program. Success of any control measure hinges upon the correct identification of the causal agent (the pathogen). Some diseases have distinctive field symptoms, but most disease identification should be confirmed by a diagnostic clinic or plant pathology laboratory. By utilizing more than one control measure, a grower may be able to reduce the chance of significant crop loss from infection by plant pathogens.

Disease management implies more than the application of chemicals at the appropriate time in the season. Knowledge of the life cycle of the pathogen, symptoms, as well as the conditions which predispose the plant to infection, contribute to effective management of cranberry diseases. Implementing cultural practices, such as trash removal, sanding, or improving bog drainage, offers opportunities to broaden the baseline defense against crop loss due to plant disease. Follow the recommended practices to encourage the natural defense mechanisms in the plant, make conditions in the bog less favorable for disease development, and decrease the presence and activity of plant pathogens.

Recommended Practices

◆ Scout for disease problems.

Measure areas of affected vines or mark with flags. Make adjustments in your disease management program before appreciable economic loss occurs.

Keep annual records which document the presence or increase of specific diseases for each bog.

Incorporate keeping quality forecasts into your fruit rot management program.

◆ Obtain proper diagnosis of new disease problems.

If the disease problem is unfamiliar or atypical of symptoms observed in the past, or if you are not getting expected response to control measures, consult with a plant pathologist. Improper diagnosis can delay the implementation of appropriate control measures and result in considerable economic loss.

Refer to the *Compendium of Blueberry and Cranberry Diseases* for aids in identification and further information.

◆ Adopt cultural disease control practices.

Sand on a regular basis to bury pathogen-infested leaf litter.

Use late water floods to reduce the incidence of fruit rot for that season.

Plant new bogs with vines taken from healthy bogs.

When leaving a *Phytophthora*-infested site, wash down picking equipment and boots with a 5% bleach solution. Take all precautions to reduce the movement of soil, water, and plant material from diseased bogs to non-infested bogs in order to limit the spread of pathogens.

Prune lush vines to promote good air circulation, reduce localized humidity, and speed the drying-out process.

Properly dispose of trash piles following harvest to remove sources of fruit rot fungal spores.

◆ **Remove trash from the bog area during or after harvest.**

Cranberry leaves, stems, and fruits left behind after harvest are colonized by fruit rot fungi. Trash can serve as a source of primary inoculum for infection of uprights, blossoms, or fruit in subsequent seasons.

For dry-harvested bogs, use a post-harvest flood in the fall. Any remaining trash should be removed from the winter flood before it is released.

Remove trash from a water-harvested bog during harvest or as soon after as possible.

Do not leave trash in piles around the bog. Move trash to the furthest reasonable distance from the production site. Avoid stockpiling trash upwind from the bog wherever possible.

◆ **Proper use of water is an important component to successful disease management.**

Run sprinkler systems for at least 4-5 hours in the **early morning** when irrigation is necessary. Short (1-2 hr), frequent intervals of watering on established bogs promote shallow root growth which is more susceptible to heat and drought stress.

Use two or three irrigation sessions to provide approximately 1" of water per week (if no rainfall occurs). Newly planted bogs may need to be watered daily in short episodes, (about 2 hr) until vines are established, see Cranberry Bog Construction and Renovation BMP.

Schedule your irrigation such that you do not extend the period of time that the vines are continuously wet. Infection by fruit rot fungi is favored when the fruit and vines are continuously wet for long periods.

Reduce the movement of water from diseased bogs to non-infested bogs in order to limit the spread of pathogens.

◆ **Use practices which minimize plant stress or lush growth. Avoid promoting conditions which favor pathogen infection.**

Avoid excessive nitrogen applications and improve bog drainage. Vine density and poor drainage often prevent rapid drying in the vine canopy and favor infection by fruit rot fungi.

Reduce plant stress from drought and heat. Irrigate vines to provide evaporative cooling. Run sprinklers for at least 1 hour and continue irrigating to keep the vines damp until the sheltered temperature falls below 85° F. See Water Management BMP.

When conditions favor oxygen deficiency (e. g., snow cover, clouded ice, or sand on ice) withdraw water from beneath the ice to provide air circulation. Winter injury may be minimized by keeping the bog flooded when soil is frozen and/or cold, windy conditions are predicted. Vines under stress (including heat and drought) are more susceptible to upright dieback.

Properly apply insecticides, herbicides, and fertilizers throughout the season. Plants weakened by pest pressures, phytotoxicity, or poor nutrition management are more susceptible to disease.

◆ **Plant new or renovated bogs with varieties which exhibit disease resistance.**

The most commonly used varieties in Massachusetts are listed from the most resistant to least resistant to fruit rot organisms: Howes, Stevens, Franklin, Bergman, Early Black, McFarlin, Ben Lear, and Crowley.

Black Veil, Foxboro Howes, Matthews, Shaw's Success, and Wilcox have shown *good fruit rot resistance* in experimental field trials. Current availability of these varieties is limited.

No variety appears to be resistant to *Phytophthora cinnamomi*, although Stevens and Black Veil appear to be less susceptible to attack by the pathogen than other varieties.

To confirm a variety is true to type, bring uprights and *attached fruit* to the Cranberry Experiment Station.

◆ **Minimize mechanical injury to fruit during dry harvesting.**

Make sure harvesting equipment is properly adjusted. The incidence of storage rot may be increased when fruit are injured by improperly operating harvesting machines. Injured fruit are more susceptible to infection because the fungi can enter through the wounds.

Exercise caution when dumping berries into empty bins. Take all precautions to minimize fruit bruising.

◆ **Fungicides may need to be integrated with cultural controls to obtain adequate disease management.**

For most fungal disease problems in cranberries, control is obtained by *preventing* initial attack by the pathogen. A steady increase or noticeable change in disease problems over a few years may indicate a need to adjust your disease management program.

Do not combine fungicides with insecticides or combine two different fungicides (i.e. Ridomil and chlorothalonil) in the same spray tank. Do not use additional stickers with chlorothalonil products. Refer to the Cranberry Chart Book for more information.

Add a spray adjuvant to applications of maneb/mancozeb formulations to improve distribution and deposition of these compounds.

If you use fungicides late in the season, use less persistent fungicides at the lowest effective rate to reduce fungicide residue on fruit.

For each application system, determine and use the optimum amount of water, pressure, injection timing, etc., needed to obtain a good distribution of fungicide.

◆ **Based on conditions on your bogs, use the lowest effective rate and number of fungicide applications for fruit rot control.**

When the keeping quality forecast, as reported by the UMass Cranberry Experiment Station, is favorable and/or a bog has a history of low rot, consider using reduced rates of fungicides that control fruit rot fungi. Fewer applications of mid to high range rates may also be considered.

Reduced rates or number of applications may be used in years with a good to excellent keeping quality forecast.

If a late water flood was used, the number of fungicide applications for fruit rot control can be reduced without compromising fruit rot control.

Consider using less fungicide per season (lower rates or fewer number of applications) if berries are grown for the processed fruit market rather than the fresh fruit market. You are only protecting against field rot as compared to field and storage rot for fresh fruit.

In years of good to excellent keeping quality forecasts, no fungicides should be applied on late water bogs with berries grown for the processed market.

◆ **Keep accurate records.**

Accurate records of fungicide applications are essential for farm planning and performance evaluation. Fungicide application dates, formulations and rates used, keeping quality forecasts (preliminary and final), and fruit rot history should be recorded on an annual basis.

Maintain records of any clinical diagnosis performed.

For further information:

Caruso, F. L. and Ramsdell, D. C. 1995. **Compendium of blueberry and cranberry diseases.** American Phytopathological Society Press, St. Paul, MN. 84 pp.

Cranberry Bog Construction and Renovation and Water Management BMPs in this series.

Cranberry chart book - management guide for Massachusetts. University of Massachusetts Cranberry Experiment Station.

