

11-2011

2011 Quinstar Use Survey

Hilary A. Sandler

University of Massachusetts - Amherst, hsandler@umass.edu

Follow this and additional works at: http://scholarworks.umass.edu/cranberry_research_repts



Part of the [Agriculture Commons](#), and the [Plant Sciences Commons](#)

Sandler, Hilary A., "2011 Quinstar Use Survey" (2011). *Cranberry Station Research Reports and Surveys*. 6.
http://scholarworks.umass.edu/cranberry_research_repts/6

This Article is brought to you for free and open access by the Cranberry Station Research Reports and Surveys at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Cranberry Station Research Reports and Surveys by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.



Hilary Sandler

Distributed November 2011, UMass Cranberry Station

Mailed to 40 growers; 22 responded = 55% return rate.

% farm treated for dodder: 48% said 100%; 14% said 90-99%; 14% said 60-70%

Describe dodder situation:

- 20% controlled with one application of Casoron (30-50 lb)
- 32% with 2 appl of Callisto (6-8 oz)
- 33% with 1 app of QS (8-12 oz)
- 23% with 2 app of QS (8 oz)
- 37% said managed only when Kerb was available.
- 32% said it is worse no matter what is done
- 14% said it is going to be a serious financial issue
- 41% said it's the biggest pest problem

Rank important contributors to lack of control (numbers of responses)

Herbicides not effective	11 noted as #1
Lots of seeds	6 noted as #1
Timing was off	6 noted as #2

How/when did dodder get on your farm?

Flood waters, birds, equipment, outside harvesting, passing of harvest boxes, no idea, geese; most said a long time ago (17 responded).

QuinStar use patterns

66% applied twice; 33% applied once. All used 4L. Almost all chemigated.
 Number of days between applications: 30 to 36 days (11 responses)
 16 used 8 oz; 3 used 12 oz (first and only application).

First application: 5 applied May 13; 2 applied May 6 and May 22; 1 applied May 3,5,6,8,12,18,19,20,24,28,29

Second application: 2 on June 5,8,10, 13, 16; 1 on May 29, June 2,9,27

%acres treated for dodder: 29% treated 100%; 20% treated 90-99%

History of dodder? 43% said very widespread; 43% moderate; and 14% had small patches

What was your control with what you did? 33% said good to excellent; 48% said fair to moderate; 19% said poor

How did you time your spray?

53% found seedlings and sprayed right away.

29% found seedlings and waited 4 to 7 days.

18% talked to other growers.

29% talked to extension specialist.

12% talked to ag supplier or consultants.

Vine damage? 55% reported slight; 25% moderate; 30% reported cupping.; 28% said vines grew out of noticed damage.

Seed reduction? 40% said yes, 15% said no, 45% not sure

Use again in 2012? 50% yes, 15% no, and 35% not sure.

	<u>What will you do if QS is available?</u>	<u>What if NOT available?</u>	(# responses)
Use Casoron only	0	1	
Use Callisto only	2	4	
Use QS only	1	-	
Casoron & QS	2	-	
Casoron & Callisto	3	12	
QS & Callisto	6	-	
All 3	5	-	
Floods	1	1	
Flame cultivation	0	1	
Hand pull	0	3	
Rake	5	4	
Seeds in flood	8	4	
Remove weeds w dodder	5	6	
MRLs	-	2	

Correlations:

- No relationship between CONTROL (Excellent=1, Good=2, Moderate=3, and 4=Poor) and date of first application, number of app, days btw apps, history of dodder infestation, and applying right away ($P>0.15$).
- Fruit destination (1=fresh, 2=processed, 3=both) was negatively correlated with control ($P=0.016$). Fresh fruit tended to have worse control. Fruit destination was positively correlated with history of infestation ($P=0.038$; widespread, moderate, patchy). Fresh fruit tended to have heavier dodder infestations.
- History was positively correlated ($P=0.002$) with timing methods (1=right away, 2=number of days later, 3=spoke to someone, 4=used calendar). Growers with worse infestations tend to time their applications as soon as they saw seedlings emerge.