University of Massachusetts - Amherst
ScholarWorks@UMass Amherst

Cranberry Chart Book - Management Guide

Cranberry Station Outreach and Public Service Activities

1-1-2010

2010 Chart Book: Disease Management At A Glance

Frank L. Caruso
University of Massachusetts - Amherst, fcaruso@umext.umass.edu

Follow this and additional works at: http://scholarworks.umass.edu/cranchart

Caruso, Frank L., "2010 Chart Book: Disease Management At A Glance" (2010). *Cranberry Chart Book - Management Guide.* Paper 24.
http://scholarworks.umass.edu/cranchart/24

This Teaching is brought to you for free and open access by the Cranberry Station Outreach and Public Service Activities at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Cranberry Chart Book - Management Guide by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

DISEASE MANAGEMENT 2010 AT A GLANCE

Prepared by Frank L. Caruso

University of Massachusetts Amherst, Cranberry Station, PO Box 569, East Wareham, MA 02538

ALWAYS READ THE LABEL and REFER TO 2010 CHART BOOK

FUNGICIDE	RATE/A	DISEASE	NOTES
Aliette WDG	5 lb	Phytophthora root rot	30 day PHI, maximum 4 uses
Ridomil Gold SL Ridomil Gold GR	1-1.75 pt 20-35 lb		45 day PHI, maximum 3 uses Do not apply EC by air.
Phostrol Pro Phyt	24 pt 4 pt		3 day PHI, maximum 4 uses No PHI, maximum 3 uses
Bravo Weather Stik, Chloronil 720, Chlorothalonil 720 SC, Equus 720 SST Bravo Ultrex, Equus DF Echo 720 Echo 90 DF	4-6.5 pt 3.8-6 lb 4-7 pt 3.25-5.75 lb	upright dieback	50 day PHI, maximum 3 uses per season, 12 hr reentry, DO NOT add spreader-sticker or Dipel.
Champ Formula 2 Flowable Champ DP Dry Prill Champion WP, Champ WG	5.3 pt 5.3 lb 8 lb		No PHI, 24 hr reentry, 5 uses max
Bravo Weather Stik, Chloronil 720, Chlorothalonil SC , Equus 720 SST Echo 720 Echo 90DF Bravo Ultrex, Equus DF Chlorothalonil 720 Equus 500 ZN	4-6.5 pt 4-7 pt 3.25-5.75 lb 3.8-6.0 lb 4-6.67 pt 5.75-9.25 pt	fruit rot (start 10–20% bloom)	50 day PHI, maximum 3 uses per season. Do NOT add spreader- sticker or Dipel
Ferbam Granuflo	6 lb		Maximum 5 uses per season, 50 day PHI, 24 hr reentry.
Abound	6.2-15.4 oz		Should not use more than 2 times per season. Is very phytotoxic to McIntosh apples.
Indar 75 WSP Indar 2F	4 oz 6-12 oz		Should not use more than 2 times per season. 30 day PHI.
Top Cop with Sulfur	2 qt	fruit rot (starting mid-bloom)	Max. 3 uses per season, no PHI
Dithane M-45 or DF Rainshield Penncozeb 75DF or 80WP Manzate Pro-Stick	3-6 lb		30 day PHI, 24 hr reentry, spray adjuvants may be added
Maneb 75 DF, Maneb 80 WP	4.8-6 lb		
Manex, Dithane F-45 Rainshield Manzate Flowable, Penncozeb 4 FL	2.4-4.8 qt		30 day PHI, 24 hr reentry, spray adjuvants may be added
ManKocide Cuprofix MZ Disperss	10.5 lb 7.5-14 lb		30 day PHI, 24 hr reentry
3 lb copper flowable Champ Formula 2 Flowable Champ DP Dry Prill Kocide DF, Champ WG Nu-Cop 50DF, Champion WP, Nu-Cop 50 WP Kocide 2000 Nu-Cop HB Nu-Cop 3L Kocide 3000 Copper-Count-N	10.67 pt 5.33 pt 5.6 lb 8 lb 8 lb 6 lb 4 lb 5.33-10.67 pt 3.5 lb 8 qt	fruit rot (starting late bloom)	Maximum 3 uses per season (except 5 for Champ), no PHI, 24 hr reentry
Ferbam Granuflo	9 lb/100 gal	fairy ring	Apply 1 gal/sq ft in June-July, one application, 24 hr reentry.
Sul-Po-Mag or K-Mag 0-0-22	4000 lb/A or 1.5 oz/sq ft		Mid-August through October
Abound	15.4 oz		First app at bud break, followed 2-4 weeks later, if necessary
Indar 75 WSP Indar 2F	4 oz 12 oz		First app at bud break, followed 2-4 weeks later, if necessary