

3-2013

## 2013 Chart Book: Disease Management At-A-Glance

Frank L. Caruso

*University of Massachusetts Amherst Cranberry Station, fcaruso@umass.edu*

Follow this and additional works at: <http://scholarworks.umass.edu/cranchart>



Part of the [Agriculture Commons](#), and the [Plant Sciences Commons](#)

---

Caruso, Frank L., "2013 Chart Book: Disease Management At-A-Glance" (2013). *Cranberry Chart Book - Management Guide*. 160.  
<http://scholarworks.umass.edu/cranchart/160>

This Public Service and Outreach is brought to you for free and open access by the Cranberry Station Outreach and Public Service Activities at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Cranberry Chart Book - Management Guide by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact [scholarworks@library.umass.edu](mailto:scholarworks@library.umass.edu).

## DISEASE MANAGEMENT 2013 AT A GLANCE

Prepared by Frank L. Caruso

University of Massachusetts Amherst, Cranberry Station  
PO Box 569, East Wareham, MA 02538

### ALWAYS READ THE LABEL and REFER TO 2013 CHART BOOK

FUNGICIDE	RATE/A	DISEASE	NOTES
Aliette WDG (fosetyl-al)	5 lb	<b>Phytophthora root rot</b>	3 day PHI, maximum 4 uses
Legion 80WDG	5 lb		
Phostrol (phosphites)	5-6 pt		3 day PHI, maximum 4 uses
Pro Phyt (phosphites)	4 pt		No PHI, maximum 3 uses
Ridomil Gold SL (mefenoxam)	1-1.75 pt		45 day PHI, maximum 3 uses Do not apply SL by air
Ridomil Gold GR (mefenoxam)	20-35 lb		45 day PHI, maximum 3 uses
Ultra Flourish (mefenoxam)	2-3.5 pt		
<hr/>			
<b>CHLOROTHALONILS</b>			
Bravo Ultrex	3.8-6 lb	<b>upright dieback</b>	ALL formulations are Zone II restricted Limit 3 applications per season 12 hour reentry interval DO NOT add spreader-sticker Do not mix with Dipel 50 day PHI
Bravo Weather Stik	4-6.5 pt		
Chloronil 720	4-6.5 pt		
Chlorothalonil 720 SC	4-6.5 pt		
Echo 720	4-7 pt		
Echo 90 DF	3.25-5.75 lb		
Equus DF	3.8-6 lb		
Equus 720 SST	4-6.5 pt		
Equus 500 ZN	5.75-9.25 pt		
<hr/>			
<b>COPPERS</b>			
Champ DP Dry Prill	5.3 lb		No PHI, 48 hr reentry, 5 uses max, except 3 for Champ WG
Champ Formula 2 Flowable	5.3 pt		
Champ WG	4.2 lb		
<hr/>			
Abound (azoxystrobin)	15.5 oz/30-100 gal	<b>fairy ring</b>	First app at bud break, followed 2-4 weeks later, if necessary
Ferbam Granuflo	9 lb/100 gal		
Indar 75 WSP (fenbuconazole)	4 oz/30-100 gal		First app at bud break, followed 2-4 weeks later, if necessary
Indar 2F (fenbuconazole)	12 oz/30-100 gal		
Sul-Po-Mag or K-Mag 0-0-22	4000 lb/A or 1.5 oz/sq ft		Mid-August through October, Use alone or after lime

SEE OTHER SIDE FOR FRUIT ROT 

## DISEASE MANAGEMENT 2013 AT A GLANCE

Prepared by Frank L. Caruso

University of Massachusetts Amherst, Cranberry Station  
PO Box 569, East Wareham, MA 02538

### ALWAYS READ THE LABEL and REFER TO 2013 CHART BOOK

FUNGICIDE	RATE/A	DISEASE	NOTES
Abound (azoxystrobin)	6.0-15.5oz	<b>fruit rot (start 10–20% bloom)</b>	Use only 2 times per season Very phytotoxic to McIntosh apples Hold water 14 days
<b><u>CHLOROTHALONILS</u></b>			
Bravo Ultrex	3.8-6 lb		ALL formulations are Zone II restricted Limit 3 applications per season 12 hour reentry interval DO NOT add spreader-sticker Do not mix with Dipel 50 day PHI
Bravo Weather Stik	4-6.5 pt		
Chloronil 720	4-6.5 pt		
Chlorothalonil 720 SC	4-6.5 pt		
Echo 720	4-7 pt		
Echo 90 DF	3.25-5.75 lb		
Equus DF	3.8-6 lb		
Equus 720 SST	4-6.5 pt		
Equus 500 ZN	5.75-9.25 pt		
Ferbam Granuflo	6 lb		
Indar 75 WSP (fenbuconazole)	4 oz		Do not use more than 2 times per season, 30 day PHI
Indar 2F (fenbuconazole)	6-12 oz		
<hr/>			
<b><u>COPPERS</u></b>			
Cuprofix MZ Disperss	7.5-14 lb	<b>fruit rot (starting mid-bloom)</b>	30 day PHI, 24 hr reentry No PHI, 48 hr reentry Max 3 uses per season, no PHI
Cuprofix Ultra 40 Disperss	5 lb		
Top Cop with Sulfur	2 qt		
<b><u>MANCOZEBS</u></b>			
Dithane DF Rainshield	3-6 lb		30 day PHI, 24 hr reentry, spray adjuvants may be added
Dithane F-45 Rainshield	2.4-4.8 qt		
Dithane M-45	3-6 lb		
ManKocide	7 lb		
Manzate Flowable	2.4-4.8 qt		
Manzate Pro-Stick	3-6 lb		
Penncozeb 4 FL	2.4-4.8 qt		
Penncozeb 75DF	3-6 lb		
Penncozeb 80WP	3-6 lb		
<hr/>			
<b><u>COPPERS</u></b>			
Badge SC	7 pt	<b>fruit rot (starting late bloom)</b>	Maximum 3 uses per season (except 5 for Champ DP Dry Prill and Champ Formula 2 Flowable), no PHI 24 hr reentry except 48 for Kocide 2000 Kocide 3000, Champ DP Dry Prill, Champ Formula 2 Flowable, Champ WG, Nu-Cop HB, Nu-Cop 3L and 12 hr for Nordox WG
Champ DP Dry Prill	5.6 lb		
Champ Formula 2 Flowable	5.33 pt		
Champ WG	4.2 lb		
Copper-Count-N	8 qt		
Kocide 2000	6 lb		
Kocide 3000	3.5 lb		
Nordox	8 lb		
Nordox 75WG	5 lb		
Nu-Cop HB	4.2 lb		
Nu-Cop 3L	4.33-5.67 pt		
Nu-Cop 50DF	8 lb		
3 lb copper flowable	10.67 pt		

SEE OTHER SIDE FOR OTHER DISEASES →