

2001

The weak ope and dimension-eight operators

E Golowich

golowich@physics.umass.edu

Follow this and additional works at: http://scholarworks.umass.edu/physics_faculty_pubs



Part of the [Physical Sciences and Mathematics Commons](#)

Recommended Citation

Golowich, E, "The weak ope and dimension-eight operators" (2001). *HIGH ENERGY PHYSICS, VOLS I AND II*. 465.
http://scholarworks.umass.edu/physics_faculty_pubs/465

This Article is brought to you for free and open access by the Physics at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Physics Department Faculty Publication Series by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

THE WEAK OPE AND DIMENSION-EIGHT OPERATORS

E. GOLOWICH

Physics Department, University of Massachusetts, Amherst MA 01003, USA

E-mail: golowich@physics.umass.edu

We discuss recent work which identifies a potential flaw in standard treatments of weak decay amplitudes, including that of ϵ'/ϵ . The point is that (contrary to conventional wisdom) dimension-eight operators contribute to weak amplitudes at order $G_F\alpha_s$ and without $1/M_W^2$ suppression. The effect of dimension-eight operators is estimated to be at the 100% level in a sum rule determination of the operator $\mathcal{Q}_7^{(6)}$ for $\mu = 1.5$ GeV, suggesting that presently available values of μ are too low to justify the neglect of these effects.

1 Motivation

1.1 Calculating Kaon Weak Amplitudes

The modern approach to calculating a kaon weak nonleptonic amplitude \mathcal{M} involves use of the operator product expansion,

$$\mathcal{M} = \sum_d \sum_i \mathcal{C}_i^{(d)}(\mu) \langle \mathcal{Q}_i^{(d)} \rangle_\mu, \quad (1)$$

in which the nonleptonic weak hamiltonian \mathcal{H}_W is expressed as a linear combination of local operators $\mathcal{Q}_i^{(d)}$. There is a sum over the dimensions (starting here at $d = 6$) of the local operators and a sum over all operators of a common dimension. In practice, the following hybrid methodology is employed:

1. The Wilson coefficients $\mathcal{C}_i^{(d)}(\mu)$ are calculated in \overline{MS} renormalization.
2. The operator matrix elements $\langle \mathcal{Q}_i^{(d)} \rangle_\mu$ are calculated in cutoff renormalization at the energy scale μ . The term ‘cutoff’ means specifically that μ serves as a ‘separation scale’ which distinguishes between short-distance and long-distance physics. Three different approaches falling into this category are quark models, $1/N_c$ expansion methods, and lattice-QCD evaluations.^a

The reason for this hybrid approach is that it is not practical to carry out the (low energy) kaon matrix element evaluations with

\overline{MS} renormalization. Typical choices for the scale μ fall in the range $0.5 \leq \mu(\text{GeV}) \leq 3$, the lower part used in quark-model and $1/N_c$ evaluations and the upper part in lattice simulations.

The purpose of this talk is to describe some recent results:¹

1. In a pure cutoff scheme, dimension-eight operators occur in the weak hamiltonian at order $G_F\alpha_s/\mu^2$, μ being the separation scale. This can be explicitly demonstrated (see Sect. 2) in a calculation involving a LR weak hamiltonian.
2. In dimensional regularization (DR), the $d = 8$ operators do *not* appear explicitly in the hamiltonian at order $G_F\alpha_s$. However, the use of a cutoff scheme for the calculation of the matrix elements of dimension-six operators requires a careful matching onto DR for which dimension-eight operators *do* play an important role.

These findings mean that hybrid evaluations, in the sense described above, of kaon matrix elements at low μ will contain (unwanted) contributions from dimension-eight operators. At the very least, this will introduce an uncertainty of unknown magnitude into the evaluation.

2 Cutoff Renormalization

^aA list of references is given elsewhere.¹

2.1 ϵ'/ϵ in the Chiral Limit

The determination of ϵ'/ϵ can be shown to depend upon the matrix elements $\langle(\pi\pi)_0|\mathcal{Q}_6^{(6)}|K\rangle$ and $\langle(\pi\pi)_2|\mathcal{Q}_8^{(6)}|K\rangle$.² In the chiral limit of vanishing light-quark mass, the latter matrix element (as well as that of operator $\mathcal{Q}_7^{(6)}$) can be inferred from certain vacuum expectation values, $\langle 0|\mathcal{O}_{1,8}^{(6)}|0\rangle \equiv \langle\mathcal{O}_{1,8}^{(6)}\rangle$, where $\mathcal{O}_{1,8}^{(6)}$ are dimension-six four-quark operators.³ The use of soft-meson techniques to relate physical amplitudes to those in the world of zero light-quark mass is a well-known procedure of chiral dynamics.

2.2 Sum Rules for $\langle\mathcal{O}_{1,8}^{(6)}\rangle$

Numerical values for $\langle\mathcal{O}_{1,8}^{(6)}\rangle$ in cutoff renormalization can be obtained from the following sum rules,³

$$\begin{aligned} \frac{16\pi^2}{3}\langle\mathcal{O}_1^{(6)}\rangle_\mu^{(c.o.)} &= \int_0^\infty ds s^2 \ln \frac{s+\mu^2}{s} \Delta\rho \\ 2\pi\langle\alpha_s\mathcal{O}_8^{(6)}\rangle_\mu^{(c.o.)} &= \int_0^\infty ds s^2 \frac{\mu^2}{s+\mu^2} \Delta\rho, \end{aligned} \quad (2)$$

where $\Delta\rho(s)$ is the difference of vector and axialvector spectral functions, and $\Delta\Pi(Q^2)$ is the corresponding difference of isospin polarization functions ($\mathcal{I}m \Delta\Pi = \pi\Delta\rho$).

2.3 Physics of a LR Operator

One can probe the influence of $d = 8$ operators by considering the K-to- π matrix element $\mathcal{M}(p)$,

$$\mathcal{M}(p) = \langle\pi^-(p)|\mathcal{H}_{LR}|K^-(p)\rangle, \quad (3)$$

where \mathcal{H}_{LR} is a LR hamiltonian obtained by flipping the chirality of one of the quark pairs in the usual LL hamiltonian \mathcal{H}_W . The reason for defining such a LR operator is that, in leading chiral order, its K-to- π matrix element is nonzero and yields information on $\langle\mathcal{O}_1^{(6)}\rangle$ and $\langle\mathcal{O}_8^{(6)}\rangle$.

To demonstrate this, we proceed to the chiral limit to find

$$\begin{aligned} \mathcal{M} &\equiv \mathcal{M}(0) = \lim_{p=0} \mathcal{M}(p) \\ &= \frac{3G_F M_W^2}{32\sqrt{2}\pi^2 F_\pi^2} \int_0^\infty dQ^2 \frac{Q^4}{Q^2 + M_W^2} \Delta\Pi. \end{aligned} \quad (4)$$

This result is *exact* — it is not a consequence of any model. Information about $\langle\mathcal{O}_1^{(6)}\rangle$ and $\langle\mathcal{O}_8^{(6)}\rangle$ is obtained by performing an operator product expansion on $\Delta\Pi(Q^2)$. Working to first order in α_s we have

$$\begin{aligned} \mathcal{M} &= \frac{G_F}{2\sqrt{2}F_\pi^2} \left[\langle\mathcal{O}_1^{(6)}\rangle_\mu^{(c.o.)} \right. \\ &\quad \left. + \frac{3}{8\pi} \ln \frac{M_W^2}{\mu^2} \langle\alpha_s\mathcal{O}_8^{(6)}\rangle_\mu + \frac{3}{16\pi^2} \frac{\mathcal{E}_\mu^{(8)}}{\mu^2} + \dots \right] \end{aligned} \quad (5)$$

The three additive terms in Eq. (5) are proportional respectively to the quantities $\langle\mathcal{O}_1^{(6)}\rangle$, $\langle\mathcal{O}_8^{(6)}\rangle$ and $\mathcal{E}^{(8)}$. The last of these ($\mathcal{E}^{(8)}$) contains the effect of the $d = 8$ contributions.^b For dimensional reasons, $\mathcal{E}^{(8)}$ must be accompanied by an inverse squared energy. This turns out to be the factor μ^{-2} .

In Table 1 we display the numerical values (in units of 10^{-7} GeV^2) of the three terms of Eq. (5) for various choices of μ . Observe for the lowest values that the dimension-eight term dominates the contribution from $\langle\mathcal{O}_1^{(6)}\rangle$. Only when one proceeds to a sufficiently large value like $\mu = 4 \text{ GeV}$ is the $d = 8$ influence suppressed.

3 Dimensional Regularization

Suppose one wishes to express the entire analysis in terms of \overline{MS} quantities. To do so requires converting matrix elements in cut-off renormalization to those in \overline{MS} renormalization. Recall, in dimensional regularization

^bAlthough the $d = 8$ LL operators arising from $\mathcal{Q}_2^{(6)}$ have been determined¹, to our knowledge the individual $d = 8$ LR operators comprising $\mathcal{E}^{(8)}$ have not.

Table 1. Eq. (5) in units of 10^{-7} GeV².

μ (GeV)	Term 1	Term 2	Term 3
1.0	-0.12	-3.84	0.64
1.5	-0.28	-3.49	0.30
2.0	-0.44	-3.24	0.17
4.0	-0.89	-2.63	0.04

one calculates in d dimensions and for dimensional consistency introduces a scale $\mu_{\text{d.r.}}$.

The dimensionally regularized matrix element for $\langle \mathcal{O}_1^{(6)} \rangle$ is found from the d -dimensional integral,³

$$\langle \mathcal{O}_1^{(6)} \rangle_{\mu}^{(\text{d.r.})} = \langle \mathcal{O}_1^{(6)} \rangle_{\mu}^{(\text{c.o.})} + \frac{d-1}{(4\pi)^{d/2}} \frac{\mu_{\text{d.r.}}^{4-d}}{\Gamma(d/2)} \int_{\mu^2}^{\infty} dQ^2 Q^d \Delta\Pi(Q^2). \quad (6)$$

The term in Eq. (6) containing the integral is proof that the dimensionally regularized matrix element $\langle \mathcal{O}_1^{(6)} \rangle_{\mu}^{(\text{d.r.})}$ will contain *short-distance* contributions. As written, this term becomes divergent for four dimensions and also is scheme-dependent. In the $\overline{\text{MS}}$ approach, the divergent factor $2/\epsilon - \gamma + \ln(4\pi)$ is removed. The NDR scheme involves a certain procedure for treating chirality in d -dimensions. The final result is a relation (given here to $\mathcal{O}(\alpha_s)$) between the cutoff and $\overline{\text{MS}}$ -NDR matrix elements,

$$\begin{aligned} \langle \mathcal{O}_1^{(6)} \rangle_{\mu}^{(\overline{\text{MS}}\text{-NDR})} &= \langle \mathcal{O}_1^{(6)} \rangle_{\mu}^{(\text{c.o.})} \\ &+ \frac{3}{8\pi} \left[\ln \frac{\mu_{\text{d.r.}}^2}{\mu^2} - \frac{1}{6} \right] \langle \alpha_s \mathcal{O}_8^{(6)} \rangle_{\mu} \\ &+ \frac{3}{16\pi^2} \cdot \frac{\mathcal{E}_{\mu}^{(8)}}{\mu^2} + \dots \end{aligned} \quad (7)$$

The effect of the $d = 8$ contribution to the weak OPE now appears in the $d = 6$ $\overline{\text{MS}}$ -NDR operator matrix element. Note also that the parameter $\mu_{\text{d.r.}}$ is distinct from the separation scale μ .

4 Evaluation of $B_{7,8}^{(3/2)}$

To suppress the effect of dimension-eight operators on the determinations of Eq. (2), one should evaluate the two sum rules for $\langle \mathcal{O}_{1,8} \rangle_{\mu}^{(\text{c.o.})}$ at a large value of μ (*e.g.* $\mu \geq 4$ GeV) and then use renormalization group equations to run the matrix elements down to lower values of μ (*e.g.* $\mu = 2$ GeV).⁴ Alternative approaches might involve the finite energy sum rule framework⁵ or QCD-lattice simulations at sufficiently large μ .

5 Concluding Remarks

This talk has dealt with an important aspect of calculating kaon weak matrix elements, the role of dimension-eight operators. In this regard, Eq. (7) is of special interest. It reveals that the relation between $\overline{\text{MS}}$ -NDR and cutoff matrix elements will involve not only mixing between operators of a given dimension but also mixing between operators of differing dimensions. The net result of our work is that existing work on ϵ'/ϵ will be affected, especially for methods which take $\mu \leq 2$ GeV.

Acknowledgments

This work was supported in part by the National Science Foundation.

References

1. V. Cirigliano, J.F. Donoghue and E. Golowich, *Dimension-eight operators in the weak OPE*, hep-ph/0007196.
2. For a review see A. J. Buras, *Weak hamiltonian, CP violation and rare decays*, hep-ph/9806471.
3. J.F. Donoghue and E. Golowich, *Phys. Lett. B* **478**, 172 (2000).
4. V. Cirigliano, J.F. Donoghue and E. Golowich, work in progress.
5. E. Golowich and K. Maltman, work in progress.