

2003

A characterization of Dynkin elements

PE Gunnells

University of Massachusetts - Amherst, gunnells@math.umass.edu

E Sommers

University of Massachusetts - Amherst, esommers@math.umass.edu

Follow this and additional works at: http://scholarworks.umass.edu/math_faculty_pubs

Recommended Citation

Gunnells, PE and Sommers, E, "A characterization of Dynkin elements" (2003). *MATHEMATICAL RESEARCH LETTERS*. 789.
http://scholarworks.umass.edu/math_faculty_pubs/789

This Article is brought to you for free and open access by the Mathematics and Statistics at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Mathematics and Statistics Department Faculty Publication Series by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

A CHARACTERIZATION OF DYNKIN ELEMENTS

PAUL E. GUNNELLS AND ERIC SOMMERS

ABSTRACT. We give a characterization of the Dynkin elements of a simple Lie algebra. Namely, we prove that one-half of a Dynkin element is the unique point of minimal length in its N -region. In type A_n this translates into a statement about the regions determined by the canonical left Kazhdan-Lusztig cells, which leads to some conjectures in representation theory.

1. INTRODUCTION

Let G be a connected simple algebraic group over the complex numbers and \mathfrak{g} its Lie algebra. We begin by recalling the Dynkin-Kostant classification of nilpotent G -orbits in \mathfrak{g} (see [4]).

Let $H = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ and $E = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ be elements of $\mathfrak{sl}_2(\mathbf{C})$. Let \mathcal{A}_{hom} denote the G -conjugacy classes of Lie algebra homomorphisms from $\mathfrak{sl}_2(\mathbf{C})$ to \mathfrak{g} . Then the map $\Omega : \mathcal{A}_{hom} \rightarrow \{\text{nilpotent } G\text{-orbits in } \mathfrak{g}\}$ given by $\Omega(\phi) = \text{Ad}(G)\phi(E)$ is a bijection: surjectivity is just the Jacobson-Morozov theorem and injectivity follows from a theorem of Kostant. Let $\Upsilon : \mathcal{A}_{hom} \rightarrow \{\text{semi-simple } G\text{-orbits in } \mathfrak{g}\}$ be the map $\Upsilon(\phi) = \text{Ad}(G)\phi(H)$. A theorem of Mal'cev shows that Υ is injective. Hence nilpotent orbits in \mathfrak{g} are parametrized by the image of Υ . This finite set was completely determined by Dynkin [5].

If \mathcal{O} is a nilpotent orbit, then an element in the conjugacy class $\Upsilon(\Omega^{-1}(\mathcal{O}))$ is called a *Dynkin element* for \mathcal{O} (or for any element $e \in \mathcal{O}$). If we fix a Cartan subalgebra \mathfrak{h} of \mathfrak{g} and a set of simple roots $\Pi \subset \mathfrak{h}^*$, then there is a unique Dynkin element h for \mathcal{O} which lies in \mathfrak{h} and such that $\alpha(h) \geq 0$ for all $\alpha \in \Pi$. We often refer to this element as **the** Dynkin element for \mathcal{O} ; the numbers $\alpha(h)$ are assigned to the Dynkin diagram of G and yield the weighted Dynkin diagram of \mathcal{O} . To summarize, this diagram completely determines the Dynkin elements for \mathcal{O} and hence \mathcal{O} itself.

The aim of this paper is to give a new characterization of the Dynkin elements. We begin by partitioning the dominant Weyl chamber into regions indexed by nilpotent orbits (which we call N -regions). These regions arise by generalizing in the most straightforward fashion the notion of S -cells (also called geometric cells) from the finite Weyl group to the affine Weyl group; they are closely related to Lusztig's \tilde{S} -cells [12]. Our main theorem states that the point of minimal length in the closure of the N -region corresponding to an orbit \mathcal{O} is one-half the Dynkin element for \mathcal{O} . In type A_n , the closure of an N -region coincides with the region determined by a canonical left Kazhdan-Lusztig cell, so our theorem becomes a statement about Kazhdan-Lusztig cells.

The Dynkin elements for G are important for the representation theory of an algebraic group of type dual to G over a local field. This is a part of conjectures by Arthur as

Date: 20 January 2003.

explained in Vogan [20]. Arthur's conjectures posit that one-half of a Dynkin element for a group of type dual to the type of G gives rise to a spherical representation of G which is unitary. This has been proved in some cases: see [20] for references. In particular, when the local field is the complex numbers, the central role played by one-half of a Dynkin element in the unitarity question for complex Lie groups is one of the key ideas in the work of Barbasch and Vogan [1]. It leads them to introduce the notion of special unipotent representations for these groups. On the algebraic side, McGovern has shown in the classical groups that these elements yield completely prime primitive ideals in the universal enveloping algebra of the Lie algebra of type dual to \mathfrak{g} (see the last section for a precise statement) [14]. The main theorem of this paper suggests a generalization about where to look for other semi-simple conjugacy classes in \mathfrak{g} which may have this property. We suspect these elements will be important in representation theory and may in particular be useful for defining unipotent representations for complex Lie groups.

Here is a brief outline of the paper. In section 2, we introduce definitions and state the main theorem, Theorem 2.1. In section 3, we give a proof. In section 4, we make a definition of a property relating Dynkin elements to B -stable ideals in the nilradical of a Borel subalgebra. We use this definition to explore further properties of the Dynkin elements as minimal points of certain convex regions. In the final section, we speculate on how generalizing the idea of the main theorem to Kazhdan-Lusztig cells and \tilde{S} -cells for G of general type may lead to an important new class of infinitesimal characters.

We thank J. Humphreys, G. Lusztig, W. McGovern, I. Mirkovic, A. Ram, D. A. Vogan, Jr., and the referee for helpful discussions and/or suggestions. We gratefully acknowledge the support of NSF grants DMS-0245580 (P.G.) and NSF grants DMS-0201826 and DMS-9729992 (E.S.).

2. MAIN THEOREM

Let (Φ, X, Φ^\vee, Y) be the root datum of a connected simple algebraic group G defined over the complex numbers with respect to a maximal torus T . Let W be the Weyl group. Fix a Borel subgroup B containing T and let Φ^+ be the determined positive roots and Π the simple roots. Let $\mathfrak{g}, \mathfrak{b}, \mathfrak{t}$ be the Lie algebras of G, B, T , respectively. Let $V = Y \otimes \mathbf{R}$. We make V into a Euclidean space by fixing a W -invariant positive-definite inner product (\cdot, \cdot) , with the induced norm denoted $\|\cdot\|$. The inner product is unique up to a non-zero scalar, since V is an irreducible representation of W . The closure of the dominant Weyl chamber in V is

$$\mathfrak{C} := \{x \in V \mid \langle \alpha, x \rangle \geq 0 \text{ for all } \alpha \in \Phi^+\},$$

where $\langle \cdot, \cdot \rangle$ is the extension of the pairing of weights and coweights.

Inspired by the work of Shi [15], [16], we partition V into regions, called ST -regions (ST stands for sign type), as follows. Let $x \in V$. For each positive root $\alpha \in \Phi^+$, let X_α equal $+$, 0 , or $-$, according to whether $\langle \alpha, x \rangle \geq 1$, $0 \leq \langle \alpha, x \rangle < 1$, or $\langle \alpha, x \rangle < 0$, respectively. The set of values $\{X_\alpha \mid \alpha \in \Phi^+\}$ which arise in this fashion is called an admissible sign type. Now V can be partitioned according to sign type (we drop the word admissible): two elements $x, y \in V$ belong to the same ST -region if they determine the same sign type. It is clear that the interior of an ST -region coincides with a connected component of the hyperplane arrangement studied by Shi [16].

In this note we are interested in the dominant sign types (and corresponding ST -regions). These are the sign types with no minus signs; the corresponding regions lie in \mathfrak{C} . The dominant sign types have an important interpretation in terms of the Lie algebra \mathfrak{g} of G . Namely, let $\mathfrak{g}_\alpha \subset \mathfrak{g}$ be the root space for $\alpha \in \Phi$. Then for each dominant sign type $\mathcal{S} = \{X_\alpha\}$ the subspace

$$\mathfrak{n}_{\mathcal{S}} := \bigoplus_{X_\alpha=+} \mathfrak{g}_\alpha$$

is a B -stable ideal in the nilradical \mathfrak{n} of \mathfrak{b} ; and every such ideal arises in this way. The fact that every ideal arises in this way can be traced back to Shi's characterization of admissible sign types [16]; more recently, it was proved in a different manner by Cellini-Papi [3].

We now define a partition of \mathfrak{C} into regions called N -regions indexed by the nilpotent orbits in \mathfrak{g} ; these are closely related to Lusztig's \tilde{S} -cells in the affine Weyl group (see the last section). Each N -region will be a union of certain ST -regions. Let \mathcal{S} be a dominant sign type and let $\mathfrak{n}_{\mathcal{S}}$ be the corresponding B -stable ideal. Let $\mathcal{O}_{\mathcal{S}}$ denote the unique nilpotent orbit of \mathfrak{g} such that $\mathcal{O}_{\mathcal{S}} \cap \mathfrak{n}_{\mathcal{S}}$ is dense in $\mathfrak{n}_{\mathcal{S}}$. Given a nilpotent orbit \mathcal{O} we define the N -region indexed by \mathcal{O} to be

$$N_{\mathcal{O}} := \{x \in \mathfrak{C} \mid \mathcal{O}_{\mathcal{S}_x} = \mathcal{O}\}$$

where \mathcal{S}_x is the sign type determined by x .

We now recall the definition of the Dynkin elements of G (with respect to T and B). Given a nilpotent orbit \mathcal{O} , let $e \in \mathcal{O}$ and let $\{e, h, f\}$ be an \mathfrak{sl}_2 -triple in \mathfrak{g} with $h \in \mathfrak{h} = \text{Lie}(T)$. Identifying V with the real subspace of \mathfrak{h} spanned by the coroots Φ^\vee , we always have $h \in V$ (h , in fact, lies in the integral lattice generated by the coroots; this follows from the representation theory of \mathfrak{g} and of \mathfrak{sl}_2). By altering our choice of e , we can ensure that $h \in \mathfrak{C}$. Then as noted in the introduction, h depends only on \mathcal{O} and is called the Dynkin element for \mathcal{O} .

Our main result is

Theorem 2.1. *Let \mathcal{O} be a nilpotent orbit in \mathfrak{g} . Let $N_{\mathcal{O}}$ be the corresponding N -region in \mathfrak{C} . Let x be a point of minimal Euclidean length in the closure of $N_{\mathcal{O}}$. Then $2x$ is the Dynkin element for \mathcal{O} . In particular, $N_{\mathcal{O}}$ is non-empty and x is unique.*

3. PROOF

The theorem follows easily from the next two propositions. If $h \in \mathfrak{C}$ is the Dynkin element for the orbit \mathcal{O} , we call

$$\mathfrak{n}_h = \bigoplus_{i \geq 2} \mathfrak{g}_i$$

the **Dynkin ideal** of \mathcal{O} , where \mathfrak{g}_i is the i -eigenspace for the action of $\text{ad}(h)$ on \mathfrak{g} . This is a B -stable ideal in \mathfrak{n} .

Proposition 3.1. *Let $h \in \mathfrak{C}$ be the Dynkin element for the orbit \mathcal{O} . Then $\frac{1}{2}h$ lies in $N_{\mathcal{O}}$.*

Proof. Let \mathcal{S} be the sign type for $\frac{1}{2}h$. Then clearly $\mathfrak{n}_{\mathcal{S}} = \mathfrak{n}_h$ from the definitions. The result follows from the well-known fact that $\mathcal{O} \cap \mathfrak{n}_h$ is dense in \mathfrak{n}_h [2]. \square

Fix a dominant sign type \mathcal{S} and let $\mathfrak{n}_{\mathcal{S}}$ be the corresponding ideal.

Proposition 3.2. *Let $e \in \mathfrak{n}_S$ and let $h \in \mathfrak{C}$ be the Dynkin element of the G -orbit through e . Then $\|\frac{1}{2}h\| \leq \|x\|$ for any x in the closure of the ST -region corresponding to S . If equality holds, then $x = \frac{1}{2}h$.*

Proof. The proof will be a slight modification of an argument due to Barbasch and Vogan ([1] Lemma 5.7). Since all W -invariant positive-definite inner products (\cdot, \cdot) on V agree up to a non-zero scalar, we may as well assume that (h, α^\vee) is an integer for all $\alpha^\vee \in \Phi^\vee$; this is possible since h lies in the coroot lattice. Let $\lambda \in \mathfrak{h}^*$ be the unique element such that $\lambda(x) = (h, x)$ for all $x \in V$. Then λ is integral and it is also dominant since $h \in \mathfrak{C}$.

Let (F_λ, π_λ) be the irreducible finite-dimensional representation of \mathfrak{g} of highest weight λ . The lowest weight of F_λ is $w_0(\lambda)$ where w_0 is the longest element of W (with respect to the simple reflections defined by Π). Let $\mathfrak{s} = \{e', h, f\}$ be an $\mathfrak{sl}_2(\mathbb{C})$ -triple with $e' \in \mathcal{O}_e$. Restricting F_λ to \mathfrak{s} , we see that the irreducible subrepresentation of \mathfrak{s} of largest dimension has dimension $m + 1$ where $m = \lambda(h)$ (as $h \in \mathfrak{C}$). This us allows to determine the index of nilpotency of $\pi_\lambda(e')$ on F_λ : we have

$$\pi_\lambda(e')^m \neq 0, \text{ but } \pi_\lambda(e')^{m+1} = 0.$$

Because e and e' are G -conjugate, the same holds for $\pi_\lambda(e)$. The key point, to be exploited shortly, is that $m = \lambda(h) = \|h\|^2$.

On the other hand, let x be in the ST -region for S (or in fact, its closure). Viewing $x \in \mathfrak{h}$, we can filter F_λ by the eigenspaces of the semisimple operator $\pi_\lambda(x)$ with corresponding eigenvalues greater than a fixed eigenvalue. The length of this filtration is $\lambda(x) - w_0\lambda(x)$, as $x \in \mathfrak{C}$.

Now if $\alpha \in \Phi^+$ satisfies $\alpha(x) \geq 1$, then any non-zero e_α in \mathfrak{g}_α acts on F_λ by raising degrees by at least 1. Write $e = \sum e_\alpha$ where $e_\alpha \in \mathfrak{g}_\alpha$. The fact that $e \in \mathfrak{n}_S$ and x belongs to the closure of the ST -region for S means that $\alpha(x) \geq 1$ for all α appearing in the sum for e . Consequently e acts by raising degrees by at least 1. It follows that $\pi_\lambda(e)^{M+1} = 0$ where $M \leq \lambda(x) - w_0\lambda(x)$. The latter quantity is bounded by $2\|h\|\|x\|$ by Cauchy-Schwartz and the fact that $\|w_0(h)\| = \|h\|$.

Hence $m \leq M \leq 2\|h\|\|x\|$. It follows that $\|h\|^2 \leq 2\|h\|\|x\|$ or $\|\frac{1}{2}h\| \leq \|x\|$.

If equality holds, then $(h, x) = \|h\|\|x\|$ and so x must be a positive multiple of h . Hence, $x = \frac{1}{2}h$. \square

The proof of the theorem follows: every element in the closure of $N_{\mathcal{O}}$ has size at least the size of one-half the Dynkin element of \mathcal{O} and this minimum is achieved at one-half the Dynkin element. Uniqueness is a consequence of the last statement of the proposition.

We have the following corollary:

Corollary 3.3. *If \mathcal{O}' lies in the closure of the nilpotent orbit \mathcal{O} and $\mathcal{O}' \neq \mathcal{O}$, then the size of the Dynkin element of \mathcal{O}' is strictly smaller than the size of the Dynkin element of \mathcal{O} .*

Proof. Let $\mathfrak{n}_h = \bigoplus_{i \geq 2} \mathfrak{g}_i$ be the Dynkin ideal for the Dynkin element h of \mathcal{O} . The G -saturation of \mathfrak{n}_h is by definition the set of elements $\text{Ad}(g)e$ where $g \in G$ and $e \in \mathfrak{n}_h$; this is exactly the closure of \mathcal{O} . Hence \mathcal{O}' intersects \mathfrak{n}_h and the result follows from the previous proposition applied to the sign type determined by $\frac{1}{2}h$. \square

4. PROPERTY D

In this section, we define a property for B -stable ideals that, if it holds in general, leads to a further understanding of the Dynkin elements. Furthermore, it should also be of interest in the study of B -stable ideals and their associated orbits.

Let \mathfrak{n}_S be the ideal corresponding to the dominant sign-type S . Let \mathcal{O}_S be the associated nilpotent orbit; that is, the one for which $\mathcal{O}_S \cap \mathfrak{n}_S$ is dense in \mathfrak{n}_S .

Definition 4.1. We say \mathfrak{n}_S has **property D** if there exists $e \in \mathcal{O}_S \cap \mathfrak{n}_S$ and $h \in \mathfrak{b}$ such that e and h can be extended to an $\mathfrak{sl}_2(\mathbf{C})$ -triple $\{e, h, f\}$. We say \mathfrak{n}_S has **strong property D** if the above holds for any orbit \mathcal{O} which intersects \mathfrak{n}_S .

We suspect that every B -stable ideal in \mathfrak{n} possesses strong property D for all G . Note that it is not true in general that for every element $e \in \mathfrak{n}_S$ there exists $h \in \mathfrak{b}$ such that e and h can be extended to an $\mathfrak{sl}_2(\mathbf{C})$ -triple.

Lemma 4.2. *Property D holds for every Dynkin ideal.*

Proof. Let $h \in \mathfrak{h}$ be the Dynkin element for the orbit \mathcal{O} and let $\{e, h, f\}$ be an $\mathfrak{sl}_2(\mathbf{C})$ -triple. In particular, $e \in \mathcal{O}$. Then the equation $[h, e] = 2e$ implies $e \in \mathfrak{n}_h$, and thus e and h satisfy the conditions for property D. \square

Let I and I' be two B -stable ideals in \mathfrak{n} . Suppose that $I = I' \oplus \mathfrak{g}_\beta$ for some $\beta \in \Phi^+$. For $\alpha \in \Pi$, let P_α denote the minimal parabolic subgroup containing B whose Lie algebra contains $\mathfrak{g}_{-\alpha}$. Assume that I is invariant under the action of P_α . It is not hard to see that $\langle \beta, \alpha^\vee \rangle \leq 0$.

Lemma 4.3. *Assume that $\langle \beta, \alpha^\vee \rangle < 0$ above. Then if property D (or the strong version) holds for I , it also holds I' .*

Proof. Let $e \in I$ and $h \in \mathfrak{b}$ satisfy the conditions for strong property D for the orbit \mathcal{O}_e . Since h lies in some Cartan subalgebra of \mathfrak{b} , we may assume $h \in \mathfrak{h}$ since all Cartan subalgebras of \mathfrak{b} are conjugate under B and I is B -stable.

Let L_α be the Levi subgroup of P_α containing T . Consider the representation of L_α on I and let M be the irreducible constituent containing \mathfrak{g}_β : it has lowest weight β and highest weight $\beta - \langle \beta, \alpha^\vee \rangle \alpha$, and these are distinct by the hypothesis.

Write $e = \sum e_\gamma$, a sum of non-zero root vectors. If $\gamma = \beta$ does not appear in the sum, we are done (since then $e \in I'$). If $\gamma = \beta - \langle \beta, \alpha^\vee \rangle \alpha$ does not appear in the sum, we proceed by letting $n \in N_G(T)$ represent the simple reflection $s_\alpha \in W$. Then $\text{Ad}(n)e \in I'$ and $\text{Ad}(n)h \in \mathfrak{h}$ and we are done. Finally if both the highest weight and lowest weight for M appear in the sum for e , we must have $\alpha(h) = 0$ (as $[h, e] = 2e$) and thus h lies in the center of the Lie algebra of L_α . From the representation theory of $SL_2(\mathbf{C})$ with respect to the representation M , there exists $g \in L_\alpha$ such that $\text{Ad}(g)e \in I'$. In this case, $\text{Ad}(g)h = h$ and the proof is complete. \square

The proof, in particular, implies that I and I' have the same associated orbit. The idea for considering two ideals related in the above fashion is closely connected to an equivalence relation on ideals defined in [19].

Proposition 4.4. *When G is of type A_n , strong property D holds for every B -stable ideal in \mathfrak{n} .*

Proof. Let P be a parabolic subgroup containing B and \mathfrak{n}_P the nilradical of its Lie algebra. It is not difficult (G is of type A) to show that \mathfrak{n}_P has strong property D by exhibiting explicit elements $e \in \mathcal{O}$ in a standard form for each orbit \mathcal{O} in the closure of the Richardson orbit for P . Of course, this uses the classification of nilpotent orbits in type A and knowledge of the closure relations.

Then the first theorem in [19] and the previous lemma imply that strong property D holds for every B -stable ideal in \mathfrak{n} . If we wanted only property D , Lemma 4.2 would allow us to avoid the classification of nilpotent orbits. \square

We have also checked that strong property D holds for all B -stable ideals in \mathfrak{n} for many \mathfrak{g} , including G_2 , F_4 , E_6 , and E_7 . We suspect that strong property D always holds. One application of property D is the following:

Proposition 4.5. *Let \mathcal{S} be a dominant sign type. If $\mathfrak{n}_{\mathcal{S}}$ possesses property D , then there is a convex region containing the closure of the ST -region of \mathcal{S} whose unique minimal point is one-half a Dynkin element for $\mathcal{O}_{\mathcal{S}}$ (here and only here, we do not assume that the Dynkin element lies in the closed dominant Weyl chamber).*

Proof. Let $e \in \mathfrak{n}_{\mathcal{S}}$ and $h \in \mathfrak{b}$ satisfy the conditions of property D for the orbit \mathcal{O}_e ; let $f \in \mathfrak{g}$ be such that $\{e, h, f\}$ is an $\mathfrak{sl}_2(\mathbb{C})$ -triple. We may assume $h \in \mathfrak{h}$ as in Lemma 4.3 (note that h need not lie in \mathfrak{C}).

Write $e = \sum e_{\alpha}$ where $e_{\alpha} \in \mathfrak{g}_{\alpha}$ and denote by $\text{supp}(e)$ the set of α which appear in this sum. Clearly, $\alpha(\frac{1}{2}h) = 1$ for all $\alpha \in \text{supp}(e)$.

Define the following convex region in V :

$$R_e = \{v \in V \mid \alpha(v) \geq 1 \text{ for all } \alpha \in \text{supp}(e)\}.$$

Since $e \in \mathfrak{n}_{\mathcal{S}}$, R_e contains the closure of the ST -region for \mathcal{S} . Then the proof of Proposition 3.2 yields that every point in R_e has length at least the length of $\frac{1}{2}h$. Moreover, $\frac{1}{2}h$ actually belongs to R_e . Uniqueness also follows from Proposition 3.2 (or the convexity of R_e). \square

5. RELATION TO CELLS AND REPRESENTATION THEORY

Let ${}^L\mathfrak{g}$ be the Lie algebra which is of type dual to \mathfrak{g} . Let ${}^L\mathfrak{h} \subset {}^L\mathfrak{g}$ be a Cartan subalgebra. We identify ${}^L\mathfrak{h}^* \cong \mathfrak{h}$. Let ${}^L\mathfrak{b} \subset {}^L\mathfrak{g}$ be the Borel subalgebra determined by \mathfrak{b} .

As noted in the introduction, if $\lambda \in \mathfrak{h}$ is one-half a Dynkin element, then the weight $\lambda \in {}^L\mathfrak{h}^* \cong \mathfrak{h}$ is important in representation theory. For example, let $L(\mu)$ be the irreducible representation of ${}^L\mathfrak{g}$ of highest weight $\mu - \rho$, where ρ is one-half the sum of the positive roots in ${}^L\mathfrak{g}$ determined by ${}^L\mathfrak{b}$. Then if λ is one-half a Dynkin element, the annihilator $J(\lambda)$ of $L(\lambda)$ is a maximal ideal in $U({}^L\mathfrak{g})$, the universal enveloping algebra of ${}^L\mathfrak{g}$, and this ideal is known to be completely prime in the classical groups by the work of McGovern ([14] Corollary 6.21). The idea for studying these particular weights arises in the work of Barbasch and Vogan on the unitary dual for complex Lie groups; they are known to yield (in many cases, see [1]) unitary representations. These representations are called special unipotent representations. In what follows, we locate other weights by varying the type of cell under consideration; we hope that these weights might be useful for the unitarity question for complex Lie groups (and more generally for Arthur's conjectures

for groups over local fields). As a first step, one might hope that the purely algebraic work of McGovern extends to these weights.

We begin by re-phrasing our main result in type A_n . Let θ be the highest root of Φ^+ and let s_0 be the affine reflection in the hyperplane $H_{\theta,1}$ in V where θ takes the value 1. Let s_1, \dots, s_n denote the reflections in the hyperplanes $H_{\alpha,0}$ where α takes the value 0, as α runs through the simple roots in Π . Let $\tilde{S} = \{s_0, s_1, \dots, s_n\}$. Then the affine Weyl group of G is the group generated by the elements in \tilde{S} ; this is a Coxeter group with length function denoted $l(-)$. For $w \in W_a$, let $R(w) = \{s \in \tilde{S} \mid l(ws) < l(w)\}$.

Following [13], we call $\mathcal{C} \subset W_a$ a canonical left Kazhdan-Lusztig cell if it is the intersection of a two-sided Kazhdan-Lusztig cell and the set of elements $w \in W_a$ such that $R(w) \subset \{s_0\}$. The canonical left cells are indexed by nilpotent orbits for the group dual to G ; this is Lusztig's parametrization of two-sided cells [11]. For example, the cell consisting solely of the identity element in W_a corresponds to the regular nilpotent orbit in ${}^L\mathfrak{g}$.

Let A be the fundamental alcove for W_a defined by

$$A := \{x \in V \mid \alpha(x) > 0 \text{ for } \alpha \in \Pi, \theta(x) < 1\}$$

and let \mathcal{C} be a canonical left Kazhdan-Lusztig cell in W_a . Define $\bar{\mathcal{C}}$ to be the closure of

$$\{w(A) \mid w \in \mathcal{C}\}$$

where the action of W_a on alcoves is the one in [9]. In particular, $\bar{\mathcal{C}} \subset \mathfrak{C}$.

By the classification of left Kazhdan-Lusztig cells in type A_n (due to Shi [15] and Lusztig [10]) and by a comment of Lusztig in [10] (proved by Lawton in [8]), $\bar{\mathcal{C}}$ exactly coincides with the closure of $N_{\mathcal{O}}$ where \mathcal{O} is the orbit dual to the one parametrizing \mathcal{C} . We remark that the Lawton-Lusztig result is closely related to property D.

Our main theorem now becomes:

Theorem 5.1. *Let \mathcal{O} be a nilpotent orbit in $\mathfrak{sl}_n(\mathbb{C})$ and let \mathcal{C} be the canonical left cell attached to the orbit which is dual to \mathcal{O} . Then the element of minimal length in $\bar{\mathcal{C}}$ is exactly one-half the Dynkin element for \mathcal{O} .*

Already in B_2 and G_2 when \mathcal{C} is a canonical left cell, it is no longer true that $\bar{\mathcal{C}}$ is a union of the closures of ST -regions. Nevertheless, the element λ of minimal length in $\bar{\mathcal{C}}$ is interesting: the maximal ideal $J(\lambda)$ turns out to be completely prime. We can find these weights explicitly in rank 2 by looking at the pictures in [9].

Example 5.2. In type B_2 , the elements are (written as a weighted diagram – see the introduction – with the first simple root being long): $(0, 0)$, $(0, \frac{1}{2})$, $(1, \frac{1}{2})$, $(1, 1)$, coming from the cell attached by Lusztig to the regular, the subregular, the minimal, and the zero orbit, respectively. Only the third element is not one-half a Dynkin element; the corresponding maximal ideal $J(\lambda)$ is the Joseph ideal, the unique completely prime primitive ideal in B_2 whose associated variety is the minimal orbit [6].

Example 5.3. In type G_2 , the elements are (written as a weighted diagram, with the first simple root being long): $(0, 0)$, $(\frac{1}{2}, 0)$, $(\frac{1}{2}, \frac{1}{2})$, $(1, \frac{1}{3})$, $(1, 1)$, coming from the cell attached to the regular, the subregular, the eight-dimensional, the minimal, and the zero orbit respectively. The third and fourth elements are not one-half of a Dynkin element. The

fourth element yields the Joseph ideal [6] and the third element yields a maximal ideal which Joseph showed is completely prime and whose associated variety is the eight-dimensional nilpotent orbit [7].

In the rest of the paper, G returns to being of general type. The above examples suggest

Conjecture 5.4. *Let \mathcal{C} be a canonical left Kazhdan-Lusztig cell in W_a . Let $\lambda \in {}^L\mathfrak{h}^* \cong \mathfrak{h}$ be an element of minimal length in $\bar{\mathcal{C}}$ (which we conjecture is unique). Then $J(\lambda)$ is a completely prime, maximal ideal of $\mathbf{U}({}^L\mathfrak{g})$ whose associated variety is the orbit which indexes \mathcal{C} .*

When G is not of type A_n , Joseph has shown that there is a unique completely prime, primitive ideal whose associated variety is the minimal orbit. Hence the conjecture predicts that when G is not of type A_n , the point of minimal length in the canonical left cell region attached to the minimal orbit is the infinitesimal character of the Joseph ideal.

We also note that it seems likely that the element of minimal length in $\bar{\mathcal{C}}$ is one-half a Dynkin element if and only if the orbit ${}^L\mathcal{O}$ indexing \mathcal{C} is special; in which case, the minimal element should be one-half the Dynkin element for the smallest orbit in \mathfrak{g} which maps under Lusztig-Spaltenstein duality to ${}^L\mathcal{O}$.

We now investigate another way to get weights which may be of interest in representation theory. They may be especially useful for defining unipotent representations for complex Lie groups. It would be interesting to see if there is any connection between these weights and the weights produced in [14] in the classical groups.

In [12], Lusztig defined the notion of \tilde{S} -cells in W_a . They are parametrized by pairs (\mathcal{O}, C) where \mathcal{O} is a nilpotent orbit in \mathfrak{g} and C is a conjugacy class in $A(\mathcal{O})$ (the fundamental group of \mathcal{O}). Let \mathcal{D} be an \tilde{S} -cell and let $\bar{\mathcal{D}} \subset V$ be the closure of

$$\{w(A) \mid w \in \mathcal{D} \text{ and } w(A) \subset \mathfrak{E}\}.$$

We call $\bar{\mathcal{D}}$ a (canonical left) \tilde{S} -region. It is not too hard to see that the closure of an N -region is a union of \tilde{S} -regions. More explicitly, if \mathcal{O} is a nilpotent orbit in \mathfrak{g} ,

$$\bar{N}_{\mathcal{O}} = \bigcup \bar{\mathcal{D}}_{(\mathcal{O}, C)}$$

where the union is over all conjugacy classes C in $A(\mathcal{O})$. It follows from our main theorem that one-half the Dynkin element of \mathcal{O} is the minimal element of $\bar{\mathcal{D}}_{(\mathcal{O}, C)}$ for some conjugacy class $C \subset A(\mathcal{O})$ (in fact, $C = 1$ but we omit the proof). This leads us to conjecture

Conjecture 5.5. *Let $\lambda \in {}^L\mathfrak{h}^* \cong \mathfrak{h}$ be an element of minimal length in a canonical left \tilde{S} -region. Then $J(\lambda)$ is completely prime.*

In types B_2 and G_2 the only elements we obtain by looking at \tilde{S} -regions which are not one-half a Dynkin element are the elements listed above coming from the Kazhdan-Lusztig cells (we found this from the B_2 -picture in [12] and calculating the canonical left \tilde{S} -cells in G_2). This leads us to wonder if the elements from the first conjecture are a subset of the elements coming from the second conjecture. This would follow from a conjecture of the second author [18] which we now describe.

Let $\mathcal{N}_{o,c}$ denote the set of all pairs (\mathcal{O}, C) as above. Let ${}^L\mathcal{N}_o$ denote the nilpotent orbits for the dual group ${}^L G$ in ${}^L\mathfrak{g}$. Let $d : \mathcal{N}_{o,c} \rightarrow {}^L\mathcal{N}_o$ be the map from [17].

Conjecture 5.6. *Let C be the canonical left Kazhdan-Lusztig cell indexed by the orbit ${}^L\mathcal{O} \subset {}^L\mathfrak{g}$. Then*

$$\bar{C} = \bigcup \bar{D}_{(\mathcal{O}, C)}$$

where the union is over all pairs (\mathcal{O}, C) such that $d_{(\mathcal{O}, C)} = {}^L\mathcal{O}$.

REFERENCES

- [1] D. Barbasch and D. A. Vogan, Jr., *Unipotent representations of complex semisimple Lie groups*, Ann. of Math. **121** (1985), 41–110.
- [2] R. Carter, *Finite groups of Lie type. Conjugacy classes and complex characters*, Wiley Classics Library, John Wiley & Sons, Ltd, Chichester, 1993.
- [3] P. Cellini and P. Papi, *ad-nilpotent ideals of a Borel subalgebra*, J. of Algebra **225** (2000), no. 1, 130–141.
- [4] D. H. Collingwood and W. M. McGovern, *Nilpotent Orbits in Semisimple Lie Algebras*, Van Nostrand Reinhold Mathematics Series, Van Nostrand Reinhold Co., New York, 1993.
- [5] E. B. Dynkin, *Semisimple subalgebras of semisimple Lie algebras*, AMS Translations **6** (1965), 111–244.
- [6] A. Joseph, *The minimal orbit in a simple Lie algebra and its associated maximal ideal*, Ann. Sci. Ecole Norm. Sup. (4) **9** (1976), no. 1, 1–29.
- [7] ———, *Kostant’s problem and Goldie rank*, Noncommutative harmonic analysis and Lie groups (Marseille, 1980), vol. 880, Lecture Notes in Math, no. 1, Springer, Berlin-New York, 1981, pp. 249–266.
- [8] G. Lawton, *Two-sided cells in the affine Weyl group of type A_{n-1}* , J. Algebra **120** (1989), no. 1, 74–89.
- [9] G. Lusztig, *Cells in affine Weyl groups*, Algebraic groups and related topics (Kyoto/Nagoya, 1983), Adv. Stud. Pure Math., 6, North-Holland, Amsterdam, 1985, pp. 255–287.
- [10] ———, *The two-sided cells of the affine Weyl group of type A_n* , Infinite-dimensional groups with applications, Math. Sci. Res. Inst. Publ., 4, Springer, New York, 1985, pp. 275–283.
- [11] ———, *Cells in affine Weyl groups. IV*, J. Fac. Sci. Univ. Tokyo Sect. 1A Math. **36** (1989), 297–328.
- [12] ———, *Affine Weyl groups and conjugacy classes in Weyl groups*, Transformation Groups **1** (1996), no. 1-2, 83–97.
- [13] G. Lusztig and N. H. Xi, *Canonical left cells in affine Weyl groups*, Adv. in Math. **72** (1988), no. 2, 284–288.
- [14] W. M. McGovern, *Completely Prime Maximal Ideals and Quantization*, Memoirs of the AMS **108** (1994), no. 519.
- [15] J.-Y. Shi, *The Kazhdan-Lusztig cells in certain affine Weyl groups*, Lecture Notes in Mathematics, no. 1179, Springer-Verlag, 1986.
- [16] ———, *Sign types corresponding to an affine Weyl group*, J. London Math. Soc. **35** (1987), no. 1, 56–74.
- [17] E. Sommers, *Lusztig’s canonical quotient and generalized duality*, J. of Algebra (2001), no. 243, 790–812.
- [18] ———, *Conjectures for cells in an affine Weyl groups*, in preparation, 2002.
- [19] ———, *Equivalence classes of ideals in the nilradical of a Borel subalgebra*, preprint, 2002.
- [20] D. A. Vogan, Jr., *Three-dimensional subgroups and unitary representations*, Challenges for the 21st century (Singapore, 2000), World Sci. Publishing, River Edge, NJ, 2001, pp. 213–250.

UNIVERSITY OF MASSACHUSETTS—AMHERST, AMHERST, MA 01003
E-mail address: gunnells@math.umass.edu

UNIVERSITY OF MASSACHUSETTS—AMHERST, AMHERST, MA 01003; INSTITUTE FOR ADVANCED STUDY,
PRINCETON, NJ 08540
E-mail address: esommers@math.umass.edu