

2010

# Type-1.5 superconductivity in multiband systems: the effects of interband couplings

Johan Carlstrom

Egor Babaev

*University of Massachusetts - Amherst*, babaev1@physics.umass.edu

Martin Speight

Follow this and additional works at: [http://scholarworks.umass.edu/physics\\_faculty\\_pubs](http://scholarworks.umass.edu/physics_faculty_pubs)



Part of the [Physics Commons](#)

---

## Recommended Citation

Carlstrom, Johan; Babaev, Egor; and Speight, Martin, "Type-1.5 superconductivity in multiband systems: the effects of interband couplings" (2010). *Physics Department Faculty Publication Series*. 1090.

[http://scholarworks.umass.edu/physics\\_faculty\\_pubs/1090](http://scholarworks.umass.edu/physics_faculty_pubs/1090)

This Article is brought to you for free and open access by the Physics at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Physics Department Faculty Publication Series by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact [scholarworks@library.umass.edu](mailto:scholarworks@library.umass.edu).

# Type-1.5 superconductivity in multiband systems: the effects of interband couplings

Johan Carlström<sup>1</sup>, Egor Babaev<sup>1,2</sup> and Martin Speight<sup>3</sup>

<sup>1</sup>*Department of Theoretical Physics, The Royal Institute of Technology, Stockholm, SE-10691 Sweden*

<sup>2</sup>*Department of Physics, University of Massachusetts Amherst, MA 01003 USA*

<sup>3</sup>*School of Mathematics, University of Leeds, Leeds LS2 9JT, UK*

In contrast to single-component superconductors, which are described at the level of Ginzburg-Landau theory by a single parameter  $\kappa$  and are divided in type-I  $\kappa < 1/\sqrt{2}$  and type-II  $\kappa > 1/\sqrt{2}$  classes, two-component systems in general possess three fundamental length scales and have been shown to possess a separate “type-1.5” superconducting state<sup>1,2</sup>. In that state, as a consequence of the extra fundamental length scale, vortices attract one another at long range but repel at shorter ranges, and therefore should form clusters in low magnetic fields. In such clusters one can define a negative interface energy inside a cluster and at the same time one can define a positive interface energy associated with the cluster’s boundary. In this work we present a detailed study of the appearance of type-1.5 superconductivity and the interpretation of the fundamental length scales in the case of two active bands with substantial interband couplings such as intrinsic Josephson coupling, mixed gradient coupling and density-density interactions. We show that in the presence of substantial intercomponent interactions of the above types the system supports type-1.5 superconductivity with fundamental length scales being associated with the mass of the gauge field and two masses of normal modes represented by mixed combinations of the density fields.

## I. INTRODUCTION

According to Ginzburg-Landau theory, a conventional superconductor near  $T_c$  is described by a single complex order parameter field. The physics of these systems is governed by two fundamental length scales, the magnetic field penetration depth  $\lambda$  and the coherence length  $\xi$ , and the ratio  $\kappa$  of these determines the response to an external field, sorting them into two categories as follows; type-I when  $\kappa < 1/\sqrt{2}$  and type-II when  $\kappa > 1/\sqrt{2}$ <sup>3</sup>.

Type-I superconductors expel weak magnetic fields, while strong fields give rise to formation of macroscopic normal domains with magnetic flux<sup>4</sup>. The response of type-II superconductors is completely different; below some critical value  $H_{c1}$ , the field is expelled. Above this value a superconductor forms a lattice or a liquid of vortices that have a supercurrent circulating around a normal core and carry magnetic flux through the system. Finally, at a second critical value,  $H_{c2}$  superconductivity is destroyed.

These different responses are usually viewed as consequences of the vortex interaction in these systems, the energy cost of a boundary between superconducting and normal states and the thermodynamic stability of vortex excitations. In a type-II superconductor the energy cost of a boundary between the normal and the superconducting state is negative, while the interaction between vortices is repulsive<sup>3</sup>. This leads to a formation of stable vortex lattices and liquids. In type-I superconductors the situation is the opposite; vortex interaction is attractive (thus making them unstable against collapse into one large vortex), while the boundary energy between normal and superconducting states is positive. From a thermodynamic point of view the principal difference between type-I and type-II states is the following: (i) In type-II superconductors the external magnetic field strength required to make formation of vortex excitations energetically preferred,  $H_{c1}$ , is smaller than the thermodynamical magnetic field  $H_{ct}$  (the field whose energy density is equal to the condensation energy of a superconductor, i.e. the field at which the uniform superconducting state becomes thermodynamically unstable); (ii) In type-I superconductors the field strength required to create a vortex excitation is larger than the thermodynamical critical magnetic field i.e. vortices cannot form. One can distinguish also a special “zero measure” boundary case where  $\kappa$  has a critical value exactly at the type-I/type-II boundary, which in the most common GL model parameterization corresponds to  $\kappa = 1/\sqrt{2}$ . In that case vortices do not interact<sup>5</sup> in the Ginzburg-Landau theory.

The above circumstances result in a situation where, in a strong external magnetic field, type-I superconductors usually have a tendency to minimize boundary energy between the normal and superconducting states, leading to a formation of large inclusions of normal phase which frequently have laminar structure<sup>4</sup>.

Recently there has been increased interest in superconductors with several superconducting components. The main situations where multiple superconducting components arise are (i) multiband superconductors<sup>6-11</sup>, (ii) mixtures of independently conserved condensates such as the projected superconductivity in metallic hydrogen and hydrogen rich alloys<sup>12-14</sup> and (iii) superconductors with other than s-wave pairing symmetries. In this work we focus on the cases (i) and (ii). The principal difference between the cases (i) and (ii) is the absence of the intercomponent Josephson coupling in case (ii).

In two-band superconductors (i) the superconducting components originate from electronic Cooper pairing in different bands<sup>6</sup>. Therefore these condensates could not *a priori* be expected to be independently conserved. This, at the level of effective models should manifest itself in a rather generic presence of intercomponent Josephson coupling.

In two-band superconductors (ii) the superconducting components originate from electronic Cooper pairing in different bands<sup>6</sup>. Therefore these condensates could not *a priori* be expected to be independently conserved. This, at the level of effective models should manifest itself in a rather generic presence of intercomponent Josephson coupling.

In the case (ii) two superconducting components were predicted to originate from electronic and protonic Cooper pairing in metallic hydrogen or hydrogen-rich alloys. In the projected liquid metallic deuterium or deuterium-rich alloys, electronic superconductivity was predicted to coexist at ultra high pressures with deuteronic condensation<sup>12-14</sup>. Because electrons cannot be converted to protons or deuterons the condensates are independently conserved, and therefore in the effective model intercomponent Josephson coupling is forbidden on symmetry grounds. These states are currently a subject of a renewed experimental pursuit. They are expected to arise at high but experimentally accessible pressures ( $\approx 400GPa$ ). Current static compression experiments achieve pressures of  $\approx 350GPa$  with pressures of an order of  $1TPa$  being anticipated in diamond anvil cells experiments due to the recent availability of ultra hard diamonds. Similar two-charged component models were discussed in the context of the physics of neutron stars where they represent coexistent protonic and  $\Sigma^-$ -hyperon Cooper pairs in the neutron star interior<sup>15</sup>.

This wide variety of systems raises the need to understand and classify the possible magnetic responses of multicomponent superconductors. It was discussed recently that in multicomponent systems the magnetic response is much more complex than in ordinary systems, and that the type-I/type-II dichotomy is not sufficient for classification. Rather, in a wide range of parameters, as a consequence of the existence of three fundamental length scales, there is a separate superconducting regime where vortices have long-range attractive short-range repulsive interaction and form vortex clusters immersed in domains of two-component Meissner state<sup>1,2</sup>. Recent experimental works<sup>16,17</sup> have put forward the suggestion that this state is realized in the two-band material  $MgB_2$ , which sparked growing interest in this topic. In particular questions were raised over whether this “type-1.5” superconducting regime (as it was termed by Moshchalkov et al<sup>16</sup>, for recent works see<sup>18</sup>) is possible even in principle in the case of various non vanishing couplings (e.g. intrinsic Josephson coupling, mixed gradient couplings etc) between superconducting components in different bands.

In this work we report a detailed study of the appearance of type-1.5 superconductivity especially focusing on the case of multiband superconductivity, demonstrating the persistence of this type of superconductivity in the presence of various kinds of intercomponent couplings (such as interband Josephson coupling, mixed gradient coupling, and density-density coupling).

## II. TYPE-1.5 SUPERCONDUCTIVITY

The possibility of a new type of superconductivity, distinct from the type-I and type-II in multicomponent systems<sup>1,2</sup> is based on the following considerations. In principle the boundary problem in the Ginzburg-Landau type of equations in the presence of phase winding is

not, from a rigorous point of view, reducible to a one-dimensional problem in general. Furthermore as discussed in<sup>1,2</sup> in general in two-component models there are *three fundamental length scales* which renders the model impossible to parametrize in terms of a single dimensionless parameter  $\kappa$ . In the case where the condensates are not coupled by interband Josephson coupling but only by the vector potential these length scales are the two independent coherence lengths (set by the inverse masses of the corresponding scalar density fields) and magnetic field penetration length (set by the inverse mass acquired by the gauge field). In contrast, in the case where the condensates are coupled by interband Josephson terms, one cannot distinguish (in the GL sense) independent coherence lengths attributed to different condensates. Nonetheless, in this case the density variations can also possess two fundamental length scales<sup>2</sup>, in contrast to single-component theories. We elaborate on this fact below. In<sup>1,2</sup> vortex solutions in two-component theories were found which have non-monotonic vortex interaction, with a long range attractive part determined by a dominant at these distances density-density interaction and a short range repulsive part produced by current-current and electromagnetic interactions. An important circumstance which was demonstrated was that these vortices are thermodynamically stable in spite of the existence of the attractive tail in the interaction.

A non-monotonic intervortex interaction potential should result in the formation of vortex clusters in low magnetic field immersed into the vortexless areas, a state referred to in<sup>1</sup> as the “semi-Meissner state”.

If the vortices form clusters one cannot use the usual one-dimensional argument concerning the superconductor-to-normal domain boundary to classify the magnetic response of the system. First of all, the energy per vortex in such a case depends on whether a vortex is placed in a cluster or not: i.e. formation of a single isolated vortex might be energetically unfavorable, while formation of vortex clusters is favorable, because in cluster where vortices are placed in a minimum of the interaction potential the energy per flux quantum is smaller than that for an isolated vortex. Thus besides the energy of a vortex in a cluster, there appears an additional energy characteristic associated with the boundary of a cluster. In other words, in this situation, to determine whether a system forms vortices in response to a magnetic field, it is not sufficient to study the one-dimensional boundary problem nor the single-vortex problem, in contrast to single component systems. Moreover, in a cluster one may define a negative energy boundary while the energy cost associated with the cluster’s boundary might be positive. A magnetic phase distinct from the vortex and Meissner states is therefore possible in these circumstances.

The existence of thermodynamically stable type-1.5 superconducting regimes ultimately depends on the existence of a nonmonotonic intervortex interaction potential. It is an important question how generic this effect

is. In this work we mainly focus on multiband realizations of multicomponent superconductivity and report a detailed study of the effects of interband Josephson coupling, mixed gradient coupling, and density-density coupling terms on vortex interactions in two band superconductors. We show that (i) when these couplings are present, the system still can possess three fundamental length scales, in contrast to the two length scales in the usual single-component GL theory; (ii) non-monotonic interaction is possible in a wide parameter range in these models.

### III. THE MODEL

We study the type-1.5 regime using the following two-component Ginzburg-Landau (TCGL) free energy functional.

$$F = \frac{1}{2}(D\psi_1)(D\psi_1)^* + \frac{1}{2}(D\psi_2)(D\psi_2)^* \quad (1)$$

$$-\nu \text{Re}\left\{(D\psi_1)(D\psi_2)^*\right\} + \frac{1}{2}(\nabla \times \mathbf{A})^2 \quad (2)$$

$$+\alpha_1|\psi_1|^2 + \frac{1}{2}\beta_1|\psi_1|^4 + \alpha_2|\psi_2|^2 + \frac{1}{2}\beta_2|\psi_2|^4 \quad (3)$$

$$-\eta_1|\psi_1||\psi_2|\cos(\theta_1 - \theta_2) + \eta_2|\psi_1|^2|\psi_2|^2. \quad (4)$$

Here  $D = \nabla + ie\mathbf{A}$ , and  $\psi_a = |\psi_a|e^{i\theta_a}$ ,  $a = 1, 2$ , represent two superfluid components which, in a two-band superconductor correspond to two superfluid densities in different bands. The equivalence mapping between our units and the standard textbook units is given in Appendix A. Such models were microscopically derived in<sup>8-10</sup>. The first two terms represent standard Ginzburg-Landau gradient terms, the second term represents mixed gradient interactions which were shown to originate in two-band superconductors from impurity scattering<sup>8,9</sup>. The next term is the magnetic field energy density and the remaining terms represent an effective potential. Here we note that  $\alpha_1$  and  $\alpha_2$  can invert sign at different temperatures. The regime where  $\alpha_1$  is positive while  $\alpha_2$  is negative corresponds to the situation where one of the bands has no superconductivity of its own but nonetheless bears some superfluid density due to interband Josephson tunneling, which is represented here by the term  $\eta_1|\psi_1||\psi_2|\cos(\theta_1 - \theta_2)$ . The type-1.5 behaviour in this regime was studied in<sup>2</sup>. In this work we will mainly focus on the situation where both bands are active, i.e.  $\alpha_{1,2} < 0$ . For generality we also add a higher order density-density coupling term  $\eta_2|\psi_1|^2|\psi_2|^2$ . We also consider the case of independently conserved condensates where the third and ninth terms in (4) are forbidden on symmetry grounds, that is,  $\nu = \eta_1 = 0$  (see also remark<sup>20</sup>).

A microscopic derivation of the TCGL model (4) requires the fields  $|\psi_a|$  to be small. However it does not in principle require  $\alpha_a$  to change sign at the same temperature. Moreover, as in the case of single-component GL

theory, we expect the model (4) to give in many cases a qualitatively acceptable picture in lower temperature regimes as well. In fact, our analysis can in some cases give a qualitative picture for the case where one of the fields does not possess a GL-type effective potential because the regime where one of the bands is in a London limit (i.e. it does not possess a GL effective potential but has a small vortex core modeled by a sharp cutoff) can be recovered from our analysis as a limiting case. As will be clear from the analysis below that regime also supports type-1.5 superconductivity.

The only vortex solutions of the model (4) which have finite energy per unit length are the integer  $N$ -flux quantum vortices which have the following phase windings along a contour  $l$  around the vortex core:  $\oint_l \nabla\theta_1 = 2\pi N$ ,  $\oint_l \nabla\theta_2 = 2\pi N$ . Vortices with differing phase windings carry a fractional multiple of the magnetic flux quantum and have energy divergent with the system size. These solutions were investigated in detail in<sup>21</sup>.

In what follows we investigate only integer flux vortex solutions which are the energetically cheapest objects to produce by means of an external field in a bulk superconductor.

### IV. VORTEX ASYMPTOTICS

The key to understanding the interaction of well separated vortices is to analyze the large  $r$  asymptotics of the vortex solution. We will analyze this problem in the context of a general TCGL model whose free energy takes the form

$$F = \frac{1}{2}(D_i\psi_1)^*D_i\psi_1 + \frac{1}{2}(D_i\psi_2)^*D_i\psi_2 + \frac{1}{2}(\partial_1 A_2 - \partial_2 A_1)^2 + F_p \quad (5)$$

where  $F_p$  contains all the non-gradient terms (in particular, but not restricted to, Josephson and density-density interaction terms). This free energy is consistent with (4) in the case  $\nu = 0$ . We will show in section IV D how to handle mixed gradient terms. The precise form of  $F_p$  is not crucial for our analysis in this section. By gauge invariance, it can depend on the condensates only via  $|\psi_1|$ ,  $|\psi_2|$  and (if the condensates are not independently conserved) on  $\theta_1 - \theta_2$ . We will assume that  $F_p$  takes its minimum value (which we normalize to be 0) when  $|\psi_1| = u_1 > 0$ ,  $|\psi_2| = u_2 > 0$  and  $\theta_1 - \theta_2 = 0$ . So, either there is no phase coupling ( $F_p$  is independent of  $\theta_1 - \theta_2$ ) and the choice of  $\theta_1 - \theta_2 = 0$  is arbitrary, or the phase coupling is such as to encourage phase locking. (Note that the case of phase anti-locked fields can trivially be recovered from our analysis by mapping  $\psi_2 \mapsto -\psi_2$ ).

The field equations are obtained from  $F$  by demanding that the total free energy  $E = \int F dx_1 dx_2$  is stationary with respect to all variations of  $\psi_1$ ,  $\psi_2$  and  $A_i$ . A routine

calculation yields

$$D_i D_i \psi_a = 2 \frac{\partial F_p}{\partial \psi_a^*} \quad (6)$$

$$\partial_i (\partial_i A_j - \partial_j A_i) = e \sum_{a=1}^2 \text{Im} (\psi_a^* D_j \psi_a). \quad (7)$$

This triple of coupled nonlinear partial differential equations supports solutions of the form

$$\begin{aligned} \psi_a &= f_a(r) e^{i\theta} \\ (A_1, A_2) &= \frac{a(r)}{r} (-\sin \theta, \cos \theta) \end{aligned} \quad (8)$$

where  $f_1, f_2, a$  are real profile functions. Note that in some cases mixed gradient terms favour non-axially symmetric solutions. In this section we consider only axially symmetric vortices. Fields within the above ansatz satisfy the field equations if and only if the profile functions  $f_1(r), f_2(r), a(r)$  satisfy the coupled ordinary differential equation system

$$f_a'' + \frac{1}{r} f_a' - \frac{1}{r^2} (1 + ea)^2 f_a = \left. \frac{\partial F_p}{\partial |\psi_a|} \right|_{(u_1, u_2, 0)} \quad (9)$$

$$a'' - \frac{1}{r} a' - e(1 + ea)(f_1^2 + f_2^2) = 0. \quad (10)$$

The solution we require, the vortex, has boundary behaviour  $f_a(r) \rightarrow u_a$ ,  $a(r) \rightarrow -1/e$  as  $r \rightarrow \infty$ . So, for large  $r$ , the quantities

$$\epsilon_a(r) = f_a(r) - u_a, \quad \alpha(r) = a(r) + \frac{1}{e} \quad (11)$$

are small and so should, to leading order, satisfy the *linearization* of (9),(10) about  $(u_1, u_2, -1/e)$ . That is, at large  $r$ ,

$$\epsilon_a'' + \frac{1}{r} \epsilon_a' = \sum_{b=1}^2 \mathcal{H}_{ab} \epsilon_b \quad (12)$$

$$\alpha'' - \frac{1}{r} \alpha' - e^2 (u_1^2 + u_2^2) \alpha = 0 \quad (13)$$

where  $\mathcal{H}$  is the Hessian matrix of  $F_p(|\psi_1|, |\psi_2|, 0)$  about its minimum

$$\mathcal{H}_{ab} = \left. \frac{\partial^2 F_p}{\partial |\psi_a| \partial |\psi_b|} \right|_{(u_1, u_2, 0)}. \quad (14)$$

So  $\alpha$  decouples from  $\epsilon_1, \epsilon_2$  asymptotically, and we see immediately that

$$\alpha(r) = q_0 r K_1(\mu_A r), \quad \mu_A = e \sqrt{u_1^2 + u_2^2} \quad (15)$$

where  $K_n$  denotes the  $n$ th modified Bessel's function of the second kind<sup>22</sup>, and  $q_0$  is an unknown real constant. Hence, at large  $r$ ,

$$\mathbf{A} \sim \left( -\frac{1}{er} + q_0 K_1(\mu_A r) \right) (-\sin \theta, \cos \theta). \quad (16)$$

Since, for all  $n$ ,

$$K_n(s) \sim \sqrt{\frac{\pi}{2s}} e^{-s} \quad \text{as } s \rightarrow \infty, \quad (17)$$

it follows that the magnetic field decays exponentially as a function of  $r$ , with length scale (penetration depth)

$$\lambda \equiv \frac{1}{\mu_A} = \frac{1}{e \sqrt{u_1^2 + u_2^2}}. \quad (18)$$

By contrast, (12) represents, in general, a coupled pair of ordinary differential equations for  $\epsilon_1, \epsilon_2$ . Since  $(u_1, u_2, 0)$  is a *minimum* of  $F_p(|\psi_1|, |\psi_2|, \theta_1 - \theta_2)$ , the Hessian matrix  $\mathcal{H}$  is a *positive definite* symmetric  $2 \times 2$  real matrix. Hence its eigenvalues,  $\mu_1^2, \mu_2^2$  say, are real and positive, and its eigenvectors,  $v_1, v_2$  say, form an orthonormal basis for  $\mathbb{R}^2$ . Expanding  $\epsilon = (\epsilon_1, \epsilon_2)^T$  in the basis  $v_1, v_2$

$$\epsilon(r) = \chi_1(r) v_1 + \chi_2(r) v_2, \quad (19)$$

we see that  $\chi_1, \chi_2$  satisfy the *uncoupled* pair of ordinary differential equations

$$\chi_a'' + \frac{1}{r} \chi_a' = \mu_a^2 \chi_a, \quad (20)$$

whence

$$\chi_a(r) = q_a K_0(\mu_a r) \quad (21)$$

for some (unknown) constants  $q_1, q_2$ . Since  $v_1, v_2$  are orthonormal, there is an angle  $\Theta$ , which we call the *mixing angle*, such that the eigenvectors of  $\mathcal{H}$  are

$$v_1 = \begin{pmatrix} \cos \Theta \\ \sin \Theta \end{pmatrix}, \quad v_2 = \begin{pmatrix} -\sin \Theta \\ \cos \Theta \end{pmatrix}. \quad (22)$$

Hence, at large  $r$  the density fields behave as

$$\psi_1 \sim [u_1 + q_1 \cos \Theta K_0(\mu_1 r) - q_2 \sin \Theta K_0(\mu_2 r)] e^{i\theta} \quad (23)$$

$$\psi_2 \sim [u_2 + q_1 \sin \Theta K_0(\mu_1 r) + q_2 \cos \Theta K_0(\mu_2 r)] e^{i\theta} \quad (24)$$

where, once again,  $K_0$  is a Bessel function.

From this analysis it follows that

1. in general there are three fundamental length scales in the problem (in contrast to the two length scales of one-component Ginzburg-Landau theory) which manifest themselves in the vortex asymptotics, namely  $1/\mu_A$ ,  $1/\mu_1$  and  $1/\mu_2$ ;
2. these are constructed from the vacuum expectation values  $u_a$  of  $|\psi_a|$  (in the case of  $1/\mu_A$ ) and from the eigenvalues of  $\mathcal{H}$ , the Hessian matrix of  $F_p$  about the vacuum (i.e. the ground state);
3.  $1/\mu_A$  can be interpreted as the London penetration length of the magnetic field;

4. however, unless the mixing angle  $\Theta$  is a multiple of  $\pi/2$ ,  $1/\mu_1$  and  $1/\mu_2$  cannot be interpreted as the coherence lengths of  $\psi_1, \psi_2$  in the usual Ginzburg-Landau sense. This is because the normal modes of the field theory close to the vacuum are not  $|\psi_a| - u_a$ , but rather

$$\begin{aligned}\chi_1 &= (|\psi_1| - u_1) \cos \Theta - (|\psi_2| - u_2) \sin \Theta \\ \chi_2 &= (|\psi_1| - u_1) \sin \Theta - (|\psi_2| - u_2) \cos \Theta\end{aligned}$$

obtained by rotating through the mixing angle  $\Theta$ , which is also determined by  $\mathcal{H}$ ; Therefore in general (e.g. in the presence of intercomponent Josephson coupling) for a one-flux quantum axially symmetric vortex, the recovery of both fields  $\psi_a$  at very long range will be according to the same exponential law, set by the smaller of the masses  $\mu_1, \mu_2$ ;

5. this analysis tells us only about the vortex structure at large  $r$ . It gives no direct information on the vortex core, which is important to understand quantitatively the nature of the vortex interactions at intermediate and short distances which will be studied numerically in the section V.

Since the gauge field mediates a repulsive force between vortices, while the condensate fields mediate an attractive force, it is clear that we can read off from the above analysis the condition under which the intervortex force is attractive at long range: we require that  $1/\mu_A$  is *not* the longest of the three length scales, or, more explicitly, that (at least) one of the eigenvalues of  $\mathcal{H}$  should be less than  $\mu_A^2 = e^2(u_1^2 + u_2^2)$ . We can predict an explicit formula for the long range two-vortex interaction potential, using the point vortex formalism<sup>19</sup>. This rests on the observation that, far from its core, the fields of the vortex are identical to those of a hypothetical point particle in a linear theory with two Klein-Gordon fields ( $\chi_1$  and  $\chi_2$  above) of mass  $\mu_1, \mu_2$  and a vector field ( $A$ ) of mass  $\mu_A$ . The point particle carries scalar monopole charges  $2\pi q_1$  and  $2\pi q_2$  and a magnetic dipole moment  $2\pi q_0$ . Two such hypothetical particles held distance  $r$  apart would experience an interaction potential

$$V(r) = 2\pi [q_0^2 K_0(\mu_A r) - q_1^2 K_0(\mu_1 r) - q_2^2 K_0(\mu_2 r)]. \quad (25)$$

This formula reproduces the prediction explained above: the long range interaction will be attractive if (at least) one of  $\mu_1, \mu_2$  is less than  $\mu_A$ .

### A. The $U(1) \times U(1)$ symmetric model

We first illustrate the above analysis in the case of  $U(1) \times U(1)$  condensates coupled only by a gauge field<sup>1</sup> where

$$F_p = \alpha_1 |\psi_1|^2 + \frac{\beta_1}{2} |\psi_1|^4 + \alpha_2 |\psi_2|^2 + \frac{\beta_2}{2} |\psi_2|^4 + \text{constant}, \quad (26)$$

with both  $\alpha_1 < 0$  and  $\alpha_2 < 0$ . Here  $F_p$  is independent of  $\theta_1 - \theta_2$  and its minimum occurs at  $|\psi_a| = u_a$  where

$$u_a = \sqrt{\frac{-\alpha_a}{\beta_a}}. \quad (27)$$

The Hessian matrix of  $F_p$  at  $(u_1, u_2, 0)$  is

$$\mathcal{H} = \begin{pmatrix} -4\alpha_1 & 0 \\ 0 & -4\alpha_2 \end{pmatrix} \quad (28)$$

whose eigenvalues are  $\mu_a^2 = -4\alpha_a$ , with corresponding eigenvectors  $v_1 = (1, 0)^T$ ,  $v_2 = (0, 1)^T$ . Hence, the mixing angle is  $\Theta = 0$ , the penetration depth is

$$\lambda \equiv \frac{1}{\mu_A} = e^{-1} \left( \frac{-\alpha_1}{\beta_1} + \frac{-\alpha_2}{\beta_2} \right)^{-\frac{1}{2}} \quad (29)$$

and the density decay lengths are the usual coherence lengths

$$\xi_a \equiv \frac{1}{\mu_a} = \frac{1}{2\sqrt{-\alpha_a}}. \quad (30)$$

The criterion for long-range vortex attraction therefore amounts to the requirement that one of the coherence lengths is larger than the magnetic field penetration length,

$$\min\{2\sqrt{-\alpha_1}, 2\sqrt{-\alpha_2}\} < e\sqrt{\frac{-\alpha_1}{\beta_1} + \frac{-\alpha_2}{\beta_2}}. \quad (31)$$

Note this criterion indicates only a long range attraction. For a realization of the type-1.5 regime one should additionally require short range repulsion and thermodynamic stability of a vortex cluster (i.e. a vortex cluster should become energetically favorable to form in external fields smaller than the thermodynamical critical magnetic field). In general these aspects of the problem require numerical analysis.

### B. Josephson coupling

We next consider how this picture is influenced by the addition of an interband Josephson term which breaks the  $U(1) \times U(1)$  symmetry to  $U(1)$ ,

$$F_p = \hat{F}_p - \eta_1 |\psi_1| |\psi_2| \cos(\theta_1 - \theta_2), \quad (32)$$

where  $\hat{F}_p$  is the free energy defined in (26) and  $\eta_1 > 0$ , so that  $F_p$  is minimized when  $\theta_1 - \theta_2 = 0$ . Adding this term changes the vacuum expectation values  $u_a$  of the fields. To find  $u_1, u_2$  we must solve

$$\frac{\partial F_p}{\partial |\psi_1|} = \frac{\partial F_p}{\partial |\psi_2|} = 0, \quad (33)$$

that is,

$$+2\alpha_1 u_1 + 2\beta_1 u_1^2 = \eta_1 u_2 \quad (34)$$

$$+2\alpha_2 u_2 + 2\beta_2 u_2^2 = \eta_1 u_1. \quad (35)$$

Unfortunately, it is not possible to solve these equations explicitly, except in special cases. For particular values of the parameters  $\alpha_a, \beta_a, \eta_1$  they can easily be solved numerically, as can the eigenvalue problem for  $\mathcal{H}$ . Note that like in the case of uncoupled bands, there are in general three fundamental length scales also in the presence of the Josephson term which can then be computed. We present numerical analysis of this problem in the full Ginzburg-Landau model in section V. To make analytical advance in this section we treat the  $\eta_1$  dependence of the length scales perturbatively. That is, we will construct Taylor expansions for  $u_a(\eta_1)$ ,  $\mu_A(\eta_1)$ ,  $\mu_a(\eta_1)$  and  $\Theta(\eta_1)$ . To keep the presentation simple, we will work to order  $\eta_1^1$ , so the results will give the leading correction to the formulae of the previous section as the Josephson coupling  $\eta_1$  is “turned on”. Higher order corrections are easily computed but do not give much extra insight.

Let us denote quantities defined in the uncoupled model (at  $\eta_1 = 0$ ) with a hat, so  $\hat{u}_a = \sqrt{-\alpha_a/\beta_a}$  are the uncoupled vacuum expectation values of  $|\psi_a|$ , for example. Let  $u(\eta_1) = (u_1(\eta_1), u_2(\eta_1))^T$  and

$$G(|\psi_1|, |\psi_2|) = \left( \frac{\partial F_p / \partial |\psi_1|}{\partial F_p / \partial |\psi_2|} \right). \quad (36)$$

Then, by definition  $G(u(\eta_1)) = 0$  for all  $\eta_1$ , and  $\hat{G}(\hat{u}) = 0$ . Differentiating with respect to  $\eta_1$  (denoted by a prime), we see that

$$\begin{aligned} 0 &= \frac{\partial G}{\partial \eta_1}(\hat{u}) + \hat{\mathcal{H}}u'(0) \\ \Rightarrow u'(0) &= -\hat{\mathcal{H}}^{-1} \frac{\partial G}{\partial \eta_1}(\hat{u}) \\ &= - \begin{pmatrix} \hat{\mu}_1^{-2} & 0 \\ 0 & \hat{\mu}_2^{-2} \end{pmatrix} \begin{pmatrix} -\hat{u}_2 \\ -\hat{u}_1 \end{pmatrix}. \end{aligned} \quad (37)$$

Hence the ground state densities receive a correction linear in  $\eta_1$

$$\begin{aligned} u_1(\eta_1) &= \hat{u}_1 + \frac{\hat{u}_2}{\hat{\mu}_1^2} \eta_1 + O(\eta_1^2) \\ u_2(\eta_1) &= \hat{u}_2 + \frac{\hat{u}_1}{\hat{\mu}_2^2} \eta_1 + O(\eta_1^2). \end{aligned} \quad (38)$$

From this expression for ground state densities one can readily calculate the gauge field mass  $\mu_A(\eta_1)$ , whose inverse gives the London penetration length. One sees that

$$\mu_A(\eta_1)^2 = \hat{\mu}_A^2 + 2\hat{u}_1\hat{u}_2 \left( \frac{1}{\hat{\mu}_1^2} + \frac{1}{\hat{\mu}_2^2} \right) \eta_1 + O(\eta_1^2). \quad (39)$$

So the effect of the Josephson coupling is always to increase the vacuum expectation values of  $|\psi_a|$ , and hence to decrease the penetration depth  $1/\mu_A$ .

The other two length scales are the eigenvalues of  $\mathcal{H}$

where

$$\begin{aligned} \mathcal{H}_{ab} &= \frac{\partial^2}{\partial |\psi_a| \partial |\psi_b|} (\hat{F}_p - \eta_1 |\psi_1| |\psi_2|) \Big|_{u(\eta_1)} \\ &= \hat{\mathcal{H}}_{ab} + \eta_1 \frac{\partial^3 \hat{F}_p}{\partial |\psi_a| \partial |\psi_b| \partial |\psi_c|} \Big|_{\hat{u}} u'_c(0) \\ &\quad - \eta_1 (1 - \delta_{ab}) + O(\eta_1^2). \end{aligned} \quad (40)$$

Now

$$\frac{\partial^3 \hat{F}_p}{\partial |\psi_a| \partial |\psi_b| \partial |\psi_c|} = \begin{cases} 12\beta_a |\psi_a| & \text{if } a = b = c \\ 0 & \text{otherwise} \end{cases} \quad (41)$$

so

$$\begin{aligned} \mathcal{H} &= \hat{\mathcal{H}} + \eta_1 \begin{pmatrix} 12\beta_1 \hat{u}_1 u'_1(0) & -1 \\ -1 & 12\beta_2 \hat{u}_2 u'_2(0) \end{pmatrix} + O(\eta_1^2) \\ &= \begin{pmatrix} \hat{\mu}_1^2 + 3\eta_1 \hat{u}_2 / \hat{u}_1 & -\eta_1 \\ -\eta_1 & \hat{\mu}_2^2 + 3\eta_1 \hat{u}_1 / \hat{u}_2 \end{pmatrix} + O(\eta_1^2). \end{aligned} \quad (42)$$

So the eigenvalues  $\lambda = \mu_a^2$  satisfy the characteristic equation

$$\left( \hat{\mu}_1^2 + 3\eta_1 \frac{\hat{u}_2}{\hat{u}_1} - \lambda \right) \left( \hat{\mu}_2^2 + 3\eta_1 \frac{\hat{u}_1}{\hat{u}_2} - \lambda \right) + O(\eta_1^2) = 0. \quad (43)$$

Hence

$$\begin{aligned} \mu_1^2 &= \hat{\mu}_1^2 + 3\eta_1 \frac{\hat{u}_2}{\hat{u}_1} + O(\eta_1^2) \\ \mu_2^2 &= \hat{\mu}_2^2 + 3\eta_1 \frac{\hat{u}_1}{\hat{u}_2} + O(\eta_1^2). \end{aligned} \quad (44)$$

As with the penetration depth, the effect of Josephson coupling (at leading order), is to *decrease* the characteristic length scales  $1/\mu_a$  of both normal modes. Another effect of Josephson coupling is mixing the fields. Let us now compute the corresponding mixing angle  $\Theta(\eta_1)$ . Recall that this is, by definition, the angle such that

$$v(\eta_1) = \begin{pmatrix} \cos \Theta(\eta_1) \\ \sin \Theta(\eta_1) \end{pmatrix} \quad (45)$$

is the eigenvector of  $\mathcal{H}$  with eigenvalue  $\mu_1^2$ . We know that  $v(0) = \hat{v} = (1, 0)^T$ , that  $v(\eta_1) \cdot v(\eta_1) = 1$  and that

$$M(\eta_1)v(\eta_1) = 0, \quad \text{where } M(\eta_1) = \mathcal{H}(\eta_1) - \mu_1(\eta_1)^2 \mathbb{I}_2 \quad (46)$$

for all  $\eta_1$ . As computed above,

$$M(\eta_1) = \begin{pmatrix} 0 & 0 \\ 0 & \hat{\mu}_2^2 - \hat{\mu}_1^2 \end{pmatrix} + \eta_1 \begin{pmatrix} 0 & -1 \\ -1 & 3\left(\frac{\hat{u}_1}{\hat{u}_2} - \frac{\hat{u}_2}{\hat{u}_1}\right) \end{pmatrix} + O(\eta_1^2) \quad (47)$$

Differentiating (46) and  $v(\eta_1) \cdot v(\eta_1) = 1$  with respect to  $\eta_1$  yields

$$M'(0)\hat{v} + M(0)v'(0) = 0, \quad v(0) \cdot v'(0) = 0 \quad (48)$$

which can be solved for  $v'(0)$ . One finds that

$$v(\eta_1) = \begin{pmatrix} 1 \\ 0 \end{pmatrix} + \eta_1 \begin{pmatrix} 0 \\ (\hat{\mu}_2^2 - \hat{\mu}_1^2)^{-1} \end{pmatrix} + O(\eta_1^2). \quad (49)$$

Hence, the mixing angle is

$$\Theta(\eta_1) = \frac{\eta_1}{\hat{\mu}_2^2 - \hat{\mu}_1^2} + O(\eta_1^2). \quad (50)$$

Thus the Josephson term produces mode mixing. Clearly, this perturbative expansion is well-defined only if  $\hat{\mu}_1 \neq \hat{\mu}_2$ . In the case where  $\hat{\mu}_1 = \hat{\mu}_2$ ,  $\hat{\mathcal{H}} = \hat{\mu}_1^2 \mathbb{I}_2$  and the assertion that  $v(0) = (1, 0)^T$  is arbitrary: any orthonormal pair of vectors can be taken as the eigenvectors associated with  $\hat{\mu}_1^2, \hat{\mu}_2^2$ . Hence, the notion of ‘‘mixing angle’’ is ill-defined in this case, and is not amenable to perturbative calculation.

The normal modes are associated with the following combinations of the  $|\psi_a|$  fields

$$\begin{aligned} \chi_1 &= (|\psi_1| - u_1) \cos \left[ \frac{\eta_1}{\hat{\mu}_2^2 - \hat{\mu}_1^2} \right] - (|\psi_2| - u_2) \sin \left[ \frac{\eta_1}{\hat{\mu}_2^2 - \hat{\mu}_1^2} \right] \\ \chi_2 &= (|\psi_1| - u_1) \sin \left[ \frac{\eta_1}{\hat{\mu}_2^2 - \hat{\mu}_1^2} \right] - (|\psi_2| - u_2) \cos \left[ \frac{\eta_1}{\hat{\mu}_2^2 - \hat{\mu}_1^2} \right]. \end{aligned}$$

In principle one can associate ‘‘generalized coherence lengths’’ of these fields with the inverse masses of the model (44) which are functions of the coherence lengths ( $\hat{\mu}_a^{-1}$ ) and vacuum field densities ( $\hat{u}_a$ ) defined in the Josephson-uncoupled theory, and the strength of the Josephson coupling  $\eta_1$ . Note that, returning to the original fields  $|\psi_a|$ , the very long-range behavior of both of these density fields is governed by whichever of the fields  $\chi_{1,2}$  has the slower recovery rate. This implies that at very long range both fields  $|\psi_a|$  should have the *same* exponential recovery law set by the smaller of  $\mu_a$ . The physical meaning of mode mixing is that the variation of the original density fields  $|\psi_a|$  acquires two length scales and one should rotate the fields through the mixing angle  $\Theta$  to determine the normal modes  $\chi_{1,2}$  whose recovery rates are governed by different single exponential laws.

Another point to note here is that, from a quantitative point of view, turning on a very small Josephson coupling does not radically alter the integer flux vortices. For small  $\eta_1$ , there is a small correction to each of the length scales (all three length scales become smaller), and there is a small amount of normal mode mixing (measured by  $\Theta(\eta_1)$ ). Therefore for this class of (integer flux) vortices the addition of Josephson coupling does not represent any kind of singular perturbation.

### 1. Comparison with the case of passive second band

It is interesting to compare these results with the case where one of the bands,  $\psi_2$  say, is passive, and has superconductivity only by virtue of the Josephson coupling term<sup>2</sup>. In this case, the free energy  $F_p$  has  $\alpha_2 > 0$ . In the uncoupled model (at  $\eta_1 = 0$ ),  $F_p$  is minimized when  $|\psi_1| = \hat{u}_1 = \sqrt{-\alpha_1/\beta_1}$  and  $|\psi_2| = \hat{u}_2 = 0$ . The gauge field has mass  $\hat{\mu}_A = e\hat{u}_1$ , there is no mode mixing, and the condensates have masses  $\hat{\mu}_1 = 2\sqrt{-\alpha_1}$  and  $\hat{\mu}_2 = \sqrt{2\alpha_2}$ . The vortex solution has  $\psi_2 = 0$  everywhere

and is identical to the Abrikosov vortex of the single component GL theory with  $\psi_2$  set to zero. As the Josephson coupling is turned on,  $\psi_2$  acquires a vacuum density of order  $\eta_1^1$ , mode-mixing develops, and the three length scales acquire corrections. Repeating the arguments of the  $\alpha_2 < 0$  case, one finds that

$$\begin{aligned} u_1(\eta_1) &= \hat{u}_1 + O(\eta_1^2) \\ u_2(\eta_1) &= \frac{\hat{u}_1}{\hat{\mu}_2^2} \eta_1 + O(\eta_1^2) \\ \mu_A(\eta_1)^2 &= \hat{\mu}_A^2 + O(\eta_1^2) \\ \mu_1(\eta_1)^2 &= \hat{\mu}_1^2 + O(\eta_1^2) \\ \mu_2(\eta_1)^2 &= \hat{\mu}_2^2 + O(\eta_1^2) \\ \Theta(\eta_1) &= \frac{\eta_1}{\hat{\mu}_2^2 - \hat{\mu}_1^2} + O(\eta_1^2). \end{aligned} \quad (51)$$

A significant difference from the case of two active bands ( $\alpha_2 < 0$ ) is that all three of the length scales receive corrections only at order  $\eta_1^2$ . Nevertheless, there is mode-mixing at order  $\eta_1^1$ .

## C. Density-density coupling

A similar perturbative analysis of the case when there is bi-quadratic density-density coupling

$$F_p = \hat{F}_p + \frac{\eta_2}{2} |\psi_1|^2 |\psi_2|^2 \quad (52)$$

can be carried out. Since the calculations are similar, we merely record the results (once again, hatted parameters refer to the uncoupled  $\eta_2 = 0$  model):

$$\begin{aligned} u_1(\eta_2) &= \hat{u}_1 + \frac{\hat{u}_1 \hat{u}_2^2}{\hat{\mu}_1^2} \eta_2 + O(\eta_2^2) \\ u_2(\eta_2) &= \hat{u}_2 + \frac{\hat{u}_2 \hat{u}_1^2}{\hat{\mu}_2^2} \eta_2 + O(\eta_2^2) \\ \mu_A(\eta_2)^2 &= \hat{\mu}_A^2 + 2\hat{u}_1^2 \hat{u}_2^2 \left( \frac{1}{\hat{\mu}_1^2} + \frac{1}{\hat{\mu}_2^2} \right) \eta_2 + O(\eta_2^2) \\ \mu_1(\eta_2)^2 &= \hat{\mu}_1^2 + \left( 1 + 3 \frac{\hat{u}_2}{\hat{\mu}_1^2} \right) \hat{u}_2^2 \eta_2 + O(\eta_2^2) \\ \mu_2(\eta_2)^2 &= \hat{\mu}_2^2 + \left( 1 + 3 \frac{\hat{u}_1}{\hat{\mu}_2^2} \right) \hat{u}_1^2 \eta_2 + O(\eta_2^2) \\ \Theta(\eta_2) &= \frac{2\hat{u}_1 \hat{u}_2}{\hat{\mu}_1^2 - \hat{\mu}_2^2} \eta_2 + O(\eta_2^2). \end{aligned} \quad (53)$$

The effect of the extra term is to reduce (if  $\eta_2 > 0$ ) or increase (if  $\eta_2 < 0$ ) all three length scales and to introduce a small amount of mode mixing. We present numerical analysis of this kind of coupling in section V.



### D. Mixed gradient terms

In this section we consider the case where the free energy has gradient-gradient coupling terms,

$$F = \frac{1}{2}(D_i\psi_1)^*D_i\psi_1 + \frac{1}{2}(D_i\psi_2)^*D_i\psi_2 - \frac{\nu}{2}[(D_i\psi_1)^*D_i\psi_2 + (D_i\psi_2)^*D_i\psi_1] + F_p \quad (54)$$

where  $F_p(|\psi_1|, |\psi_2|, \theta_1 - \theta_2)$  is, as before, a non-negative function minimized at  $(u_1, u_2, 0)$ . In contrast to the previous two cases, this we can treat exactly, without resorting to power series expansion in the coupling parameter  $\nu$ . We can assume  $\nu > 0$  without loss of generality (the case  $\nu < 0$  is obtained by mapping  $\psi_2 \mapsto -\psi_2$ ), and we must have  $\nu < 1$ , or else  $F$  is not positive definite. This case does not fit into the general analysis presented above. Nonetheless a similar method, with some modification, can be applied.

The field equations are

$$D_i D_i(\psi_1 - \nu\psi_2) = 2\frac{\partial F_p}{\partial \psi_1^*} \quad (55)$$

$$D_i D_i(\psi_2 - \nu\psi_1) = 2\frac{\partial F_p}{\partial \psi_2^*} \quad (56)$$

$$\partial_i(\partial_i A_j - \partial_j A_i) = e \operatorname{Im}(\psi_1^* D_j(\psi_1 - \nu\psi_2) + \psi_2^* D_j(\psi_2 - \nu\psi_1)) \quad (57)$$

which support vortex solutions of the form (8) provided the profile functions obey the coupled system

$$\begin{aligned} \left( \frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{1}{r^2}(1+ea)^2 \right) P \begin{pmatrix} f_1 \\ f_2 \end{pmatrix} \\ = \begin{pmatrix} \partial F_p / \partial |\psi_1| \\ \partial F_p / \partial |\psi_2| \end{pmatrix} \Big|_{(f_1, f_2, 0)} \\ a'' - \frac{1}{r} a' - e(1+ea)(f_1^2 - 2\nu f_1 f_2 + f_2^2) = 0, \end{aligned} \quad (58)$$

where

$$P = \begin{pmatrix} 1 & -\nu \\ -\nu & 1 \end{pmatrix}. \quad (59)$$

The vortex boundary conditions are  $f_a(r) \rightarrow u_a$  and  $a(r) \rightarrow -1/e$  as  $r \rightarrow \infty$ . Note that these are independent of  $\nu$ . We again define  $\epsilon(r) = (f_1(r) - u_1, f_2(r) - u_2)^T$  and  $\alpha(r) = a(r) + 1/e$ , and linearize the system about  $\epsilon_1 = \epsilon_2 = \alpha = 0$ :

$$\begin{aligned} \left( \frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} \right) P\epsilon = \mathcal{H}\epsilon \\ \alpha'' - \frac{1}{r}\alpha' - e^2(u_1^2 - 2\nu u_1 u_2 + u_2^2)\alpha = 0. \end{aligned} \quad (60)$$

We immediately see that  $\mathbf{A}$  behaves asymptotically as in (16), but with

$$\mu_A = e\sqrt{u_1^2 - 2\nu u_1 u_2 + u_2^2}. \quad (61)$$

The effect of the gradient-gradient coupling is thus to *increase* the penetration depth  $1/\mu_A$ . Note that the effect would be opposite if there is a competing stronger Josephson term which enforces phase *anti-locking* in the vacuum, that is phase difference  $\theta_1 - \theta_2 = \pi$ .

To decouple the pair of equations for  $\epsilon_1, \epsilon_2$  we must expand  $\epsilon$  in a basis of eigenvectors, not of  $\mathcal{H}$ , but rather of

$$\tilde{\mathcal{H}}(\nu) = P(\nu)^{-1}\mathcal{H}. \quad (62)$$

Note that this matrix is *not* in general symmetric. Nonetheless, it can be shown that its eigenvalues are real and positive for all  $0 \leq \nu < 1$  (see the appendix B). Let  $\mu_1^2, \mu_2^2$  be the eigenvalues of  $\tilde{\mathcal{H}}$  and  $v_1, v_2$  be the corresponding eigenvectors. Then the condensate fields at large  $r$  take the form

$$\begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} \sim \left\{ \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} + q_1 K_0(\mu_1 r) v_1 + q_2 K_0(\mu_2 r) v_2 \right\} e^{i\theta} \quad (63)$$

Once again, the condition for long-range attraction is

$$\min\{\mu_1^2, \mu_2^2\} < \mu_A^2 \quad (64)$$

but now  $\mu_1^2, \mu_2^2$  are the eigenvalues of  $P(\nu)^{-1}\mathcal{H}$ , not  $\mathcal{H}$ , and  $\mu_A$  depends on  $u_1, u_2$  and  $\nu$ , as in (61). Note that

$$\mu_1^2 \mu_2^2 = \det \tilde{\mathcal{H}} = \det \mathcal{H} / \det P(\eta_1) = \frac{\det \mathcal{H}}{1 - \nu^2} \quad (65)$$

so the effect of the coupling must be to increase (at least) one of  $\mu_{1,2}$  and hence to decrease (at least) one of the normal mode recovery length scales. Since  $\tilde{\mathcal{H}}$  is not symmetric, there is no reason why  $v_1, v_2$  should be orthogonal, so it is not possible to define a single mixing angle in this case.

To illustrate, consider the simplest case, where  $F_p$  is defined as in (26). Then  $u_a = \sqrt{-\alpha_a/\beta_a}$  and the uncoupled  $\nu = 0$  model has  $\hat{\mu}_a^2 = -4\alpha_a$  and  $\hat{\mu}_A^2 = e^2(u_1^2 + u_2^2)$ . For  $\nu > 0$ , the vacuum expectation values do not change, but the penetration depth increases, since

$$\mu_A^2 = \hat{\mu}_A^2 - 2\nu u_1 u_2. \quad (66)$$

Furthermore

$$\tilde{\mathcal{H}}(\nu) = P(\nu)^{-1} \begin{pmatrix} \hat{\mu}_1^2 & 0 \\ 0 & \hat{\mu}_2^2 \end{pmatrix} = \frac{1}{1 - \nu^2} \begin{pmatrix} \hat{\mu}_1^2 & \nu \hat{\mu}_2^2 \\ \nu \hat{\mu}_1^2 & \hat{\mu}_2^2 \end{pmatrix} \quad (67)$$

whose eigenvalues are

$$\mu_{1,2}^2(\nu) = \frac{1}{2(1 - \nu^2)} \left( \hat{\mu}_1^2 + \hat{\mu}_2^2 \pm \sqrt{(\hat{\mu}_1^2 - \hat{\mu}_2^2)^2 + 4\nu^2 \hat{\mu}_1^2 \hat{\mu}_2^2} \right). \quad (68)$$

Now  $\mu_{1,2}^{-1}(\nu)$  are the new fundamental length scales which control the variation of the density fields (63). Without loss of generality, we may assume that  $\hat{\mu}_1 \geq \hat{\mu}_2$  (if  $\hat{\mu}_1 < \hat{\mu}_2$  then we simply swap the labels of the condensates). In

this case, for  $\nu > 0$  it is clear from the above expression that

$$\mu_1^2 > \frac{1}{2(1-\nu^2)} \left( \hat{\mu}_1^2 + \hat{\mu}_2^2 + \sqrt{(\hat{\mu}_1^2 - \hat{\mu}_2^2)^2} \right) = \frac{\hat{\mu}_1^2}{1-\nu^2} \quad (69)$$

so when  $0 < \nu < 1$ ,  $\mu_1(\nu) > \hat{\mu}_1$ . Recall that

$$\mu_1^2 \mu_2^2 = \det \tilde{\mathcal{H}} = \frac{\hat{\mu}_1^2 \hat{\mu}_2^2}{1-\nu^2}, \quad (70)$$

so

$$\mu_2^2 = \frac{\hat{\mu}_1^2}{\mu_1^2} \frac{\hat{\mu}_2^2}{1-\nu^2} < \hat{\mu}_2^2 \quad (71)$$

by (69), and hence  $\mu_2(\nu) < \hat{\mu}_2$  when  $0 < \nu < 1$ . In this case, the effect of gradient-gradient coupling is to decrease the smaller of the normal mode decay lengths,  $\mu_1^{-1}$ , and increase the larger,  $\mu_2^{-1}$ . Thus gradient coupling tends to increase the disparity in these length scales.

As in sections IV B and IV C it is instructive to see what happens to the parameters of the uncoupled model ( $\nu = 0$ ) as  $\nu > 0$  is turned on. This can be extracted from the above formulae by expanding in  $\nu$ , keeping only terms up to order  $\nu^1$ . One sees that

$$\begin{aligned} u_a(\nu) &= \hat{u}_a \quad \text{exactly} \\ \mu_A(\nu)^2 &= \hat{\mu}_A^2 - 2\nu \hat{u}_1 \hat{u}_2 \quad \text{exactly} \\ \mu_a(\nu)^2 &= \hat{\mu}_a^2 + O(\nu^2) \end{aligned} \quad (72)$$

so, to leading order in  $\nu$ , the only length scale that changes is the penetration depth  $1/\mu_A$ , which increases. The eigenvectors of  $\tilde{\mathcal{H}}$  are

$$v_1 = \begin{pmatrix} 1 \\ \frac{\hat{\mu}_1^2 \nu}{\hat{\mu}_1^2 - \hat{\mu}_2^2} \end{pmatrix} + O(\nu^2), \quad v_2 = \begin{pmatrix} \frac{-\hat{\mu}_2^2 \nu}{\hat{\mu}_1^2 - \hat{\mu}_2^2} \\ 1 \end{pmatrix} + O(\nu^2) \quad (73)$$

which, one should note, are *not* orthogonal: the angle between them is  $\frac{\pi}{2} - \nu + O(\nu^2)$ . It follows that the vortex has asymptotic densities (at large  $r$ )

$$\begin{aligned} |\psi_1| &\sim \hat{u}_1 + q_1 K_0(\hat{\mu}_1 r) - \frac{q_2 \hat{\mu}_2^2 \nu}{\hat{\mu}_1^2 - \hat{\mu}_2^2} K_0(\hat{\mu}_2 r) + O(\nu^2) \\ |\psi_2| &\sim \hat{u}_2 + q_2 K_0(\hat{\mu}_2 r) + \frac{q_1 \hat{\mu}_1^2 \nu}{\hat{\mu}_1^2 - \hat{\mu}_2^2} K_0(\hat{\mu}_1 r) + O(\nu^2) \end{aligned} \quad (74)$$

where  $q_1, q_2$  are unknown constants. So, while the ‘‘coherence lengths’’ remain unchanged to leading order, the normal modes with which they are associated do receive a correction at order  $\nu^1$ .

Finally, we remark that an alternative approach to handling gradient-gradient terms is to remove them from  $F$  from the outset by a linear redefinition of the fields: essentially one expands  $(\psi_1, \psi_2)^T$  in a basis of eigenvectors of  $P(\nu)^{23}$ . This is mathematically elegant, but tends to obscure the physical meaning of the non-gradient terms  $F_p$ .




FIG. 1. Non-monotonic vortex interaction in a system with a passive band (i.e. superconductivity is induced in the second band by an intrinsic proximity effect). In limit of zero Josephson coupling, the active band would have  $\kappa = 8\kappa_c$ , and thus would be deep into the type-II region. This figure shows that a perturbation in the form of a weak Josephson coupling to passive band in this case produces a minimum in the inter-vortex potential at a very large distance from the vortex center.




FIG. 2. Intervortex interaction potential for a set of systems with two active bands. The systems share the parameters given in the table ‘common parameters’. The green curve (4) corresponds to the case where the bands are coupled by the vector potential only. In this case, the ratio of the coherence lengths is  $\xi_2/\xi_1 = 4$ . The curve (2) shows the effect of the addition of Josephson term, the curve (5) shows the effect of addition of mixed gradient term. The curves (1) and (3) show the effect of the presence of both mixed gradient and Josephson terms with similar and opposite signs.





FIG. 3. Gradient, magnetic and potential energy contributions to the vortex interaction energy for the parameter set corresponding to the curve (3) of the Fig. 2 (black curve).



Curve	$\alpha_1$	$\beta_1$	$\alpha_2$	$\beta_2$	$\eta_1$	$\eta_2$	$e$	$\langle  \psi_1 ^2 \rangle$	$\langle  \psi_2 ^2 \rangle$
1	-1	1	-0.0069	0.0278	0	0	0.7	1	0.25
2	-1	1	-0.0069	0.0278	0.00556	0	0.7	1.0018	0.407
3	-1	1	-0.0069	0.0278	0.0111	0	0.7	1.004	0.526
4	-1	1	-0.0069	0.0139	0	0	0.7	1	0.5
5	-1	1	-0.0069	0.0278	0	-0.0044	0.7	1.00182	0.410

FIG. 4. Non monotonic vortex interaction in systems with two active bands and significant disparity in length scales associated with the density variation. In the absence of inter band coupling (curve 1), the ratio of the coherence length is  $\xi_2/\xi_1 = 12$ .




FIG. 5. Comparison of how Josephson coupling (1) and high order density coupling (2-4) affects vortex interaction energy. In (3),  $\eta_2$  is chosen so that the densities are approximately the same as in (1). In (4),  $\eta_2$  is chosen to give the same condensation energy as (1). Both these parameter values give larger vortex binding energy. Similar binding energy as (1) is acquired for a significantly smaller  $\eta_2$  (2). The large condensation energy associated with Josephson coupling is responsible for the shorter interaction range in (1).

## V. NUMERICAL SOLUTION OF THE NONLINEAR PROBLEM.

The linear analysis presented in the previous section can only provide information about the asymptotic tail of the intervortex interaction. To determine the actual full intervortex potential, especially in the case of strong interband coupling, it is necessary to treat the full nonlinear Ginzburg-Landau theory, something which is not possible analytically. In this section we present interaction energies for vortex pairs (including cases of relatively strong interband coupling), computed numerically using a local relaxation method. The numerical method which we use is the following: A lattice approximant of the energy is minimized with respect all the degrees of freedom in the full Ginzburg-Landau functional subject to the constraint that vortex positions remain fixed. This gives the intervortex interaction energy as a function of inter vortex separation. We used high resolutions grids, with the number of data points ranging from  $1600 \times 1600$  to  $2400 \times 1700$  and relaxed each configuration for 50–100 hours on an eight core cluster node.

In this section the length scale is given in units of  $2/\hat{\mu}_1$ , where, as in section IV,  $\hat{\mu}_1$  denotes the mass of the field  $|\psi_1|$  in the absence of interband coupling ( $\eta_1 = \eta_2 = \nu = 0$ ). Alternatively, the unit of length is  $\sqrt{2}\hat{\xi}_1$ , where  $\hat{\xi}_1 = \sqrt{2}/\hat{\mu}_1$  is the coherence length of the first band in




FIG. 6. Cross sections of two interacting vortices. The densities are given as  $1 - |\psi_i|^2 / \langle |\psi|^2 \rangle$ , i.e. the condensates are normalized to 1 and turned upside down. The magnetic field is also normalized. The left column corresponds to curve 3) and the right column correspond to curve 4) of the same figure. The inter-vortex separation is 3.0 in the upper row, giving repulsion, 5.4 in the mid row corresponding to the minimum energy, and 7.8 in the bottom row giving attraction. The curves 3 and 4 in fig 4 are the ones showing the largest binding energy, a property resulting from them having the largest vacuum expectation values in the second band (0.526, 0.5). A careful comparison shows that the recovery of the second band is slower in the right column where there is zero inter band coupling. This is a very generic result, Josephson coupling shrinks the disparity in length scales. The result of this effect is also seen when comparing the vortex interaction energies of the two system, the curve 3) reveals a shorter vortex interaction range than curve 4).

the uncoupled case. Recall that  $\hat{\xi}_1$  cannot be identified with physical coherence length when interband coupling is present. We also measure condensate density  $|\psi_a|$  in units of  $\hat{u}_1$ , the vacuum expectation value of  $|\psi_1|$  in the uncoupled case. As shown in appendix A, this amounts to using scale freedom to set  $\alpha_1 = -1$  and  $\beta_1 = 1$ . In the single component limit, the parameter  $e$  can then be interpreted as an inverse GL parameter. More precisely,  $\kappa = \sqrt{2}/e$ , so the critical value of  $e$  is  $e_c = 2$ . The inter vortex interaction energy is given in units of total vortex energy, i.e.  $2E_v$  where  $E_v$  is the energy of a single

isolated vortex. All energies are measured relative to the uniform Meissner state ( $\psi_a = u_a$ ,  $A = 0$ , in the notation of section IV).

#### A. Weak Josephson coupling to a passive band

Let us initially consider a strongly type-II superconductor with a single component, and see how vortex interaction in this system is modified by a weak Josephson coupling to a passive band<sup>2</sup> (i.e. a band which has no

Curve	$\alpha_1$	$\beta_1$	$\alpha_2$	$\beta_2$	$\eta_1$	$\eta_2$	$e$	$\langle  \psi_1 ^2 \rangle$	$\langle  \psi_2 ^2 \rangle$
1	-1	1	-0.0069	0.0139	0.0055	0	0.5	1.00237	0.73
2	-1	1	-0.0069	0.0139	0.0055	0	0.35	1.00237	0.73
3	-1	1	-0.0069	0.0139	0.0055	0	0.2	1.00237	0.73




FIG. 7. Transition from type-1.5 to type-II region in a system with two active bands. The binding energies are 0.0109 in the first curve and 0.0023 in the second curve. The third curve give monotonic repulsion. The small binding energy in the second case signals that the critical value for  $e$  is close to 0.35.

superconductivity of its own which in the context of GL theory manifests itself as a positive coefficient  $\alpha_2$ ).

The Fig. 1 shows the vortex interaction energy in such a system. In the limit of decoupled bands, the parameter  $e$  is here  $e_c/8$ . Therefore, for zero Josephson coupling this system would have  $\kappa \approx 5.7$ , putting it far into the type-II region. Weak Josephson coupling changes the length scales as discussed in the previous section and adds a qualitatively new feature: the inter vortex potential acquires a minimum, occurring at a separation of  $24\sqrt{2}\xi_1$ .

### B. Effects of Josephson and mixed gradient terms in case of two active bands

The figure 2 illustrates the effect of the mixed gradient term, as well as of Josephson coupling in a system with two active bands. The green curve (4) corresponds to two independent bands, interacting only trough the magnetic field. The curve (2) corresponds to a system with added Josephson coupling which increases the binding energy, but decreases the distance where the energy minimum is located and slightly reduces the range of interaction.

The inclusion of a mixed gradient term (shown as curve (5)) here has a similar effect on phase difference as the Josephson term. When the phases are locked  $\theta_1 - \theta_2 = 0$ , effectively this term gives a negative contribution to the energy associated with co-directed currents. Thus, for

this choice of the sign of  $\nu$ , the mixed gradient term also prefers phase locking  $\theta_1 - \theta_2 = 0$ .

Due to symmetry, changing  $\eta_1 \rightarrow -\eta_1$  and  $\nu \rightarrow -\nu$  does not qualitatively change the behavior of the system, as this only results in phase locking with  $\pi$  difference instead. While Josephson coupling increases the energy of vortices, and mixed gradients decreases it, their effect on interaction energy is the opposite. The decomposition of vortex interaction energy into a set of contributions from different terms given in Fig 3 illustrates why mixed gradients increases repulsion.

In contrast the curve (1) (blue curve), corresponds to the case where  $\nu$  and  $\eta_1$  have different signs, and so there is competition between the gradient mixing and the Josephson term with regard to the preferred phase difference. The mixed gradient term is minimal for a phase locking where  $\theta_1 - \theta_2 = \pi$ , while the Josephson term is minimal for  $\theta_1 - \theta_2 = 0$ . The result in these simulations was that the phase locking was determined by the dominating Josephson coupling, and that the gradient mixing resulted in increased cost for co-directed currents. This was the most energetically expensive vortex, but also exhibited the smallest inter vortex interaction energy.

### C. Solutions with large disparity in the characteristic length scales

Figure 4 shows a set of simulations done with two active bands and a larger disparity in characteristic length scales. We start with case when the condensates interact only through the magnetic field (blue curve (1)), and the density in the second band is 1/4 of the density in the first band and the coherence length ratio being  $\xi_2/\xi_1 = 12$  (so in the notation of section IV,  $\hat{u}_1 = 4\hat{u}_2$  and  $\hat{\mu}_1 = 12\hat{\mu}_2$ ). This allows non-monotonic interaction to occur at smaller  $e$  than above - here we simulate at  $e = 0.7$ . This gives the smallest binding energy in this set of simulations. Adding a Josephson coupling of  $\eta_1 = 0.00556$  [shown on curve (2)] gives a substantially higher density in the second band, and thus a stronger binding energy. Adding a stronger Josephson coupling [shown on the curve (3)] gives even larger density in the second band. It also shows some of the qualitative differences associated with this coupling. Namely, the binding occurs at a smaller separation, and the range of the interaction decreases. It is clearly visible that curve (3) crosses the other curves. The Josephson coupling causes the second condensates to recover faster (as follows also from the linear theory presented in the previous section), and thereby decreasing the range of attractive interaction.

Decreasing  $\hat{\beta}_2$  raises the density of condensate in the second band. In curve (4), we have reduced it by a factor 2, thereby increasing the vacuum expectation value of the density in the second band by a factor 2. This does however not change the length scale, as  $\xi$  is independent of  $\beta$ , but it does increase the energetic benefits of core

overlap in the second component. This case shows the largest binding energy in this set of simulations.

Finally, we consider the effect of a higher order density-density coupling  $\eta_2$  between the condensates which is shown on the curve (5). The parameter choice here gives approximately the same densities as (2) but smaller condensation energy. This should generally make the system (5) recover slower than system(2), resulting in longer range of the interaction. A more systematic comparison of Josephson coupling and higher order density coupling supporting this conclusion is given in Fig. 5.

The figure 6 displays cross sections of the condensate densities and magnetic flux in the systems 3-4 in Fig. 4, clearly illustrating the mechanism by which type-1.5 superconductivity appears.

Let us consider a different example of the appearance of the type-1.5 regime in the case where there is a substantial disparity in the dominant length scales associated with the variations of densities and magnetic field penetration length depth.

Again, our starting point is a reference case of two bands coupled only by vector potential where we choose  $\alpha_2$  so that the disparity in coherence length in absence of inter band coupling is  $\xi_2/\xi_1 = 12$ .

Now, we take  $\beta_2$  to be 0.0139 i.e. the same as in curve (4) of Fig. 4 and choose the Josephson coupling to be  $\eta_1 = 0.00556$ . Then, we successively decrease  $e$  to see where the system crosses over from type-1.5 to type-II. The outcome can be seen in Fig. 7. The first curve gives a binding energy of 0.011, the second gives 0.0023 and the third curve shows system crossing over into type-II regime by showing monotonic repulsion. Given the small binding energy of the second curve, the cross over from type-1.5 to type-II is close to  $e = 0.35$ .

The crosssection plots of two of these systems given in Fig 8 illustrate how the system crosses over from type-1.5 to type-II as  $e$  is decreased resulting in dominance of the repulsive interaction originating in the electromagnetic and current-current interaction.

## VI. CONCLUSIONS

In this paper we presented a detailed analytical and numerical study of the appearance of type-1.5 superconductivity in the case of two bands with various kinds of substantial interband couplings. In all the cases which we considered we demonstrated that the system possessed three fundamental length scales: one length scale  $1/\mu_A$  associated with London magnetic field penetration length while the other two fundamental scales  $1/\mu_{1,2}$  are associated with characteristic length scales controlling variations of density fields. In the limit of two condensates coupled only electromagnetically the length scales  $1/\mu_{1,2}$  are associated with standard GL coherence lengths<sup>1</sup>. However we show that introducing a nonzero Josephson and quartic density-density couplings makes both density fields decay according to the same expo-




FIG. 8. Cross sections plots showing condensate density and magnetic flux in type-1.5 superconductors with small  $e$ . a) Curve (1) if Fig. 7 at a separation of 7.2, corresponding to the energy minimum. b) The same system at a separation of 9.6. c) Curve (2) of the same figure at a separation of 9.6, corresponding to the energy minimum. The three distinct length scales associated with two band superconductors are indeed visible. Despite the inter-band Josephson coupling, there is a significant disparity in the recovery of the two condensates in all the plots. The third length scale, penetration depth, visibly differs between the two systems, and is responsible for the differences in inter vortex interaction.

nential law at very large distances from the core while, *at the same time the system still possesses two fundamental length scales which are associated now with variation of linear combinations of density fields rotated by a “mixing angle”*. The third fundamental length scale in that regime is the London penetration length and thus two-band systems with Josephson- and density-density interband couplings allow a well defined type-1.5 behavior. Next we studied the effect of mixed gradient terms and showed how the type-1.5 regime is described in that case. We showed that in the case of a substantial mixed gradient coupling the definition of three fundamental length scales requires additional care because it produces mode mixing which cannot be described by a single mixing angle. In the second part of the paper we presented a comparative numerical study of type-1.5 vortices in the different regimes with various intercomponent couplings. The results were demonstrated in the framework of a two-component Ginzburg-Landau model with local electrodynamics. However we expect that the described type-1.5 behavior is similarly present in lower-temperature regimes and in two-component models with non-local electrodynamics.

### ACKNOWLEDGEMENTS

We thank Alex Gurevich for discussions. EB was supported by Knut and Alice Wallenberg Foundation through the Royal Swedish Academy of Sciences, Swedish Research Council and by the US National Science Foundation CAREER Award No. DMR-0955902. JC was supported by the Swedish Research Council. MS was supported by the UK Engineering and Physical Sciences Research Council.

### Appendix A: Units

In this section we give an explicit mapping from our representation of the GL model to a more common textbook representation. Consider the Ginzburg-Landau model in the following quite usual units

$$\begin{aligned}
F = & \frac{\hbar^2}{2m_1} \left| \left( \nabla - i \frac{e^*}{\hbar c} A \right) \psi_1 \right|^2 \\
& + \frac{\hbar^2}{2m_2} \left| \left( \nabla - i \frac{e^*}{\hbar c} A \right) \psi_2 \right|^2 + \\
& - \nu \hbar^2 \text{Re} \left\{ \left( \nabla - i \frac{e^*}{\hbar c} A \right) \psi_1 \cdot \left( \nabla + i \frac{e^*}{\hbar c} A \right) \psi_2^* \right\} \\
& + \frac{1}{8\pi} (\nabla \times A)^2 \\
& + \alpha_1 |\psi_1|^2 + \frac{1}{2} \beta_1 |\psi_1|^4 + \alpha_2 |\psi_2|^2 + \frac{1}{2} \beta_2 |\psi_2|^4 \\
& - \eta_1 |\psi_1| |\psi_2| \cos(\theta_1 - \theta_2) + \eta_2 |\psi_1|^2 |\psi_2|^2 \quad (\text{A1})
\end{aligned}$$

Let us define the rescaled quantities

$$\begin{aligned}
\tilde{F} &= \frac{4\pi}{\hbar^2 c^2} F \\
\tilde{A} &= -\frac{A}{\hbar c} \\
\tilde{\psi}_a &= \sqrt{\frac{4\pi}{m_a c^2}} \psi_a \\
\tilde{\nu} &= \sqrt{m_1 m_2} \nu \\
\tilde{\alpha}_a &= \frac{m_a}{\hbar^2} \alpha_a \\
\tilde{\beta}_a &= \frac{m_a^2 c^2}{4\pi \hbar^2} \beta_a \\
\tilde{\eta}_1 &= \frac{\sqrt{m_1 m_2}}{\hbar^2} \eta_1 \\
\tilde{\eta}_2 &= \frac{m_1 m_2 c^2}{4\pi \hbar^2} \eta_2. \quad (\text{A2})
\end{aligned}$$

Then

$$\begin{aligned}
\tilde{F} = & \frac{1}{2} \left| (\nabla + ie^* \tilde{A}) \tilde{\psi}_1 \right|^2 + \frac{1}{2} \left| (\nabla + ie^* \tilde{A}) \tilde{\psi}_2 \right|^2 \\
& - \tilde{\nu} \text{Re} \left\{ (\nabla + ie^* \tilde{A}) \tilde{\psi}_1 \cdot (\nabla - ie^* \tilde{A}) \tilde{\psi}_2^* \right\} \\
& + \frac{1}{2} |\nabla \times \tilde{A}|^2 \\
& - \tilde{\alpha}_1 |\tilde{\psi}_1|^2 + \frac{\tilde{\beta}_1}{2} |\tilde{\psi}_1|^4 + \tilde{\alpha}_2 |\tilde{\psi}_2|^2 + \frac{\tilde{\beta}_2}{2} |\tilde{\psi}_2|^4 \\
& + \tilde{\eta}_1 |\tilde{\psi}_1| |\tilde{\psi}_2| \cos(\theta_1 - \theta_2) + \tilde{\eta}_2 |\tilde{\psi}_1|^2 |\tilde{\psi}_2|^2, \quad (\text{A3})
\end{aligned}$$

which, on dropping the tildes, coincides with the representation (4) used in this paper.

Throughout the paper, it is assumed that band 1 is active, that is,  $\alpha_1 < 0$ . It is convenient to rescale the expression (4) for  $F$  further so that  $\alpha_1$  is normalized to  $-1$  and  $\beta_1$  is normalized to 1. This can be achieved as follows. Recall (see section IV A) that in the absence of interband couplings (i.e. when  $\eta_1 = \eta_2 = \nu = 0$ ) condensate 1 has decay length-scale  $1/\hat{\mu}_1 = (-4\alpha_1)^{-1/2}$ . This scale is more usually specified by the coherence length

$$\hat{\xi}_1 = \frac{\sqrt{2}}{\hat{\mu}_1} = \frac{1}{\sqrt{-2\alpha_1}}. \quad (\text{A4})$$

We emphasize once more that, in the presence of interband couplings,  $\hat{\xi}_1$  is not the coherence length of condensate 1. This is the purpose of the hat, to remind us that this is a genuine coherence length only in the uncoupled case. Recall also that the vacuum density of condensate 1 in the uncoupled model is

$$\hat{u}_1 = \sqrt{\frac{-\alpha_1}{\beta_1}}. \quad (\text{A5})$$

Our second rescaling amounts to using  $\sqrt{2}\hat{\xi}_1$  as the unit of length and  $\hat{u}_1$  as the unit of condensate density (along with compensating rescalings of  $F$ ,  $e^*$  and  $A$ ). Explicitly,

let

$$\begin{aligned}
\bar{r} &= \frac{r}{\sqrt{2}\hat{\xi}_1} = \sqrt{-\alpha_1}r \\
\bar{F} &= \frac{2\hat{\xi}_1^2}{\hat{u}_1^4}F = \frac{\beta_1^2}{-\alpha_1^3}F \\
\bar{\psi}_a &= \frac{\psi_a}{\hat{u}_1} = \sqrt{\frac{\beta_1}{-\alpha_1}}\psi_a \\
\bar{A} &= \frac{A}{\hat{u}_1} \\
\bar{e} &= \frac{1}{\sqrt{2}}\hat{u}_1\hat{\xi}_1e^* = \frac{e^*}{\sqrt{\beta_1}} \\
\bar{\alpha}_2 &= 2\hat{\xi}_1^2\alpha_2 = \frac{\alpha_2}{-\alpha_1} \\
\bar{\beta}_2 &= 2\hat{\xi}_1^2\hat{u}_1^2\beta_2 = \frac{\beta_2}{\beta_1} \\
\bar{\eta}_1 &= 2\hat{\xi}_1^2\eta_1 = \frac{\eta_1}{-\alpha_1} \\
\bar{\eta}_2 &= 2\hat{\xi}_1^2\hat{u}_1^2\eta_2 = \frac{\eta_2}{\beta_1} \\
\bar{\nu} &= \nu.
\end{aligned} \tag{A6}$$

Substituting these into (4) yields

$$\begin{aligned}
\bar{F} &= \frac{1}{2} |(\bar{\nabla} + i\bar{e}\bar{A})\bar{\psi}_1|^2 + \frac{1}{2} |(\bar{\nabla} + i\bar{e}\bar{A})\bar{\psi}_2|^2 \\
&\quad - \bar{\nu} \text{Re} \{ (\bar{\nabla} + i\bar{e}\bar{A})\bar{\psi}_1 \cdot (\bar{\nabla} - i\bar{e}\bar{A})\bar{\psi}_2^* \} \\
&\quad + \frac{1}{2} |\bar{\nabla} \times \bar{A}|^2 \\
&\quad - |\bar{\psi}_1|^2 + \frac{1}{2} |\bar{\psi}_1|^2 - \bar{\alpha}_2 |\bar{\psi}_2|^2 + \frac{\bar{\beta}_2}{2} |\bar{\psi}_2|^2 \\
&\quad - \bar{\eta}_1 |\bar{\psi}_1| |\bar{\psi}_2| \cos(\theta_1 - \theta_2) + \bar{\eta}_2 |\bar{\psi}_1|^2 |\bar{\psi}_2|^2. \tag{A7}
\end{aligned}$$

This (having dropped the bars) is the GL energy used in section V for the purposes of numerical simulation.

Finally, we note that the single component GL model obtained from (A7) by setting  $\psi_2 \equiv 0$  has penetration depth  $\lambda = 1/\mu_a = 1/e$  and coherence length  $\xi = 1/\sqrt{2}$ , and hence GL parameter

$$\kappa = \lambda/\xi = \frac{\sqrt{2}}{e}. \tag{A8}$$

So, in the parameterization used in section V, one may regard  $e$  as an inverse GL parameter for the associated single band model. The value of  $e$  corresponding to Bogomolny limit of one-component theory is  $e_c = 2$  in this interpretation.

## Appendix B: The spectrum of $\tilde{\mathcal{H}}$

Let  $\mathcal{H}$  be a real, symmetric  $2 \times 2$  matrix both of whose eigenvalues are positive, and let  $\tilde{\mathcal{H}}(\nu) = P(\nu)^{-1}\mathcal{H}$  where  $P(\nu)$  is defined in equation (59). We wish to show that the eigenvalues  $\lambda_1(\nu), \lambda_2(\nu)$  of  $\tilde{\mathcal{H}}(\nu)$  are also real and positive for all  $0 \leq \nu < 1$ . First note that  $\tilde{\mathcal{H}}(0) = \mathcal{H}$ , so the conclusion holds at  $\nu = 0$ . Further,  $\lambda_a(\nu)$  depend continuously on  $\nu$ . Now  $\det \tilde{\mathcal{H}}(\nu) = \det \mathcal{H}/(1 - \nu^2) \neq 0$ , so neither eigenvalue ever vanishes. So  $\lambda_a(\nu)$  remain real and positive, unless, for some  $\nu = \nu_* \in (0, 1)$ , they coalesce and bifurcate into a complex conjugate pair. But then, at  $\nu = \nu_*$  we have  $\lambda_1(\nu_*) = \lambda_2(\nu_*) = \lambda_* \in (0, \infty)$  and hence  $\tilde{\mathcal{H}}(\nu_*) = \lambda_* \mathbb{I}_2$ . But then

$$\tilde{\mathcal{H}}(\nu) = P(\nu)^{-1}P(\nu_*)\tilde{\mathcal{H}}(\nu_*) = \lambda_*P(\nu)^{-1}P(\nu_*) \tag{B1}$$

which is symmetric, and hence has real eigenvalues, for all  $\nu \in (0, 1)$ . Hence a bifurcation to a complex conjugate pair of eigenvalues is not possible, and we conclude that  $\tilde{\mathcal{H}}(\nu)$  has real, positive eigenvalues for all  $\nu \in [0, 1)$ .

<sup>1</sup> E. Babaev & J.M. Speight Phys.Rev. B **72** 180502 (2005)

<sup>2</sup> E. Babaev, J. Carlstrom, J. M. Speight Phys. Rev. Lett. **105**, 067003 (2010)

<sup>3</sup> A. Abrikosov Sov. Phys. JETP **5**, 1174 (1957)

<sup>4</sup> L. Landau, Nature (London, United Kingdom) **141**, 688 (1938). R. P. Huebener, *Magnetic Flux Structures of Superconductors*, 2nd ed. (Springer-Verlag, New-York, 2001). R. Prozorov, A. F. Fidler, J. Hoberg, P. C. Canfield, Nature Physics **4**, 327 - 332 (2008)

<sup>5</sup> The noninteracting regime, which is frequently called ‘‘Bogomolny limit’’ is a property of Ginzburg-Landau model where, at  $\kappa = 1/\sqrt{2}$ , the core-core attractive interaction between vortices exactly cancels the current-current repulsive interaction. However, in a realistic system extremely close to the  $\kappa \rightarrow 1/\sqrt{2}$  limit one should go beyond the Ginzburg-Landau description to determine leftover intervortex interactions, because these are then determined not by the fundamental length scales of the GL theory but by small microscopic effects leading to non-universal

weak vortex interaction potentials. We do not consider the physics which arises in the Bogomolny limits in this work.

<sup>6</sup> H. Suhl, B. T. Matthias, and L. R. Walker Phys. Rev. Lett. **3**, 552 (1959)

<sup>7</sup> A. Liu, I.I. Mazin, J. Kortus, Phys. Rev. Lett. **87** 087005 (2001); I.I. Mazin, et al., Phys. Rev. Lett. **89** (2002) 107002.

<sup>8</sup> A. Gurevich, Phys. Rev. B **67** 184515 (2003) ;

<sup>9</sup> A. Gurevich, Physica C **056**160 (2007)

<sup>10</sup> M. E. Zhitomirsky and V.-H. Dao, Phys. Rev. B **69**, 054508 (2004).

<sup>11</sup> see e.g. K. Ishida, Y. Nakai, H., Hosono, J. Phys. Soc. Jpn. **78** 062001 (2009)

<sup>12</sup> N.W. Ashcroft, J. Phys. Condens. Matter **12**, A129 (2000); Phys. Rev. Lett. **92**, 187002 (2004) K. Mouloupoulos and N. W. Ashcroft Phys. Rev. Lett. **66** 2915 (1991),

<sup>13</sup> E. Babaev, A. Sudbø and N.W. Ashcroft Nature **431** 666 (2004), E. Smørgrav, J. Smiseth, E. Babaev, A. Sudbø. Phys. Rev. Lett. **94**, 096401 (2005)



- <sup>14</sup> E. V. Herland, E. Babaev, A. Sudbo arXiv:1006.3311 Phys. Rev. B in print
- <sup>15</sup> P. B. Jones, Mon. Not. Royal Astr. Soc. **371**, 1327 (2006); E. Babaev, Phys. Rev. Lett. **103**, 231101 (2009).
- <sup>16</sup> V.V. Moshchalkov M Menghini, T. Nishio, Q. H. Chen, A. V. Silhanek, V. H. Dao, L. F. Chibotaru, N. D. Zhigadlo, and J. Karpinski Phys. Rev. Lett. **102**, 117001 (2009)
- <sup>17</sup> T. Nishio, Q. Chen, W. Gillijns, K. De Keyser, K. Vervaeke, and V. V. Moshchalkov Phys. Rev. B **81**, 020506(R) (2010)
- <sup>18</sup> R. Geurts, M. V. Milosevic, F. M. Peeters Phys. Rev. B **81**, 214514 (2010) V. H. Dao, L. F. Chibotaru, T. Nishio, V. V. Moshchalkov arXiv:1007.1849; Shi-Zeng Lin, Xiao Hu arXiv:1007.1940
- <sup>19</sup> J.M. Speight, Phys. Rev. D **55** 3830 (1997) .
- <sup>20</sup> In the case of independently conserved condensates there could be different mixed gradient terms which are quadratic in gradients and quartic in  $\psi_a$ ; some of the effects of such terms on vortex physics were considered in E. Babaev Phys. Rev. B **79**, 104506 (2009), and in<sup>14</sup>
- <sup>21</sup> E. Babaev Phys.Rev.Lett. **89** (2002) 067001 E. Babaev, J. Jaykka, and M. Speight, Phys. Rev. Lett. **103**, 237002 (2009).
- <sup>22</sup> M. Abramowitz and I.A. Stegun, *Handbook of Mathematical Functions* (Dover, New York NY, USA, 1972) p. 374.
- <sup>23</sup> E. Babaev, J. Carlstrom, J.M. Speight not published.