



Comment on “Hausdorff Dimension of Critical Fluctuations in Abelian Gauge Theories”

Item Type	article
Authors	Prokof'ev, N;Svistunov, B
Download date	2025-04-08 03:48:30
Link to Item	https://hdl.handle.net/20.500.14394/40507

Comment on “Hausdorff Dimension of Critical Fluctuations in Abelian Gauge Theories”

In their Letter [1], Hove, Mo, and Sudbø derive a simple connection between the anomalous scaling dimension, η , of the U(1) universality class order parameter, $\phi(\mathbf{x})$, and the Hausdorff dimension, D_H , of critical loops:

$$\eta + D_H = 2. \quad (1)$$

In the loop representation, the correlator $G(\mathbf{r}) = \langle \phi(\mathbf{r})\phi^*(0) \rangle \propto r^{-(d-2+\eta)}$ describes the distribution of the end-points in open loops. For definiteness, one may think of the high-temperature-expansion loops for the lattice $|\phi|^4$ -model.

The analysis of Ref. [1] might seem absolutely compelling, being just a translation of the hyperscaling hypothesis into the loop language: *At the critical point there should be about one loop of diameter r per volume element r^d* [2]. Nevertheless, given the result $\eta = 0.0380(4)$ of Ref. [3], the relation (1) is in strong contradiction with the value $D_H = 1.7655(20)$ which we obtained for the 3D $|\phi|^4$ -model with suppressed leading corrections to scaling [3] (and also—with a bit less accuracy—for the standard bond-current model [4], and its special version with excluded loop overlaps and self-crossings). The simulations were done with the Worm algorithm [5].

The hidden flaw in the treatment of Ref. [1] is as follows. When introducing the self-similar expression

$$P(\mathbf{r}; N) \propto N^{-\rho} F(r/N^\Delta), \quad \Delta = 1/D_H \quad (2)$$

for the probability to find the ends of an open loop of length N being distance \mathbf{r} away from each other, which is then used to establish the connection between the open and closed loops, the authors take for granted that $F(0)$ is finite. While looking innocent, this is an *arbitrary* assumption, since the self-similar form (2) is valid only for $r \gg a$, where a is a microscopic cutoff (e.g., the lattice period). Strictly speaking, a closed loop of length N corresponds to $F(a/N^\Delta)$ rather than to $F(0)$, and one has to work with the generic asymptotic form

$$F(x) \propto x^\theta \quad \text{at} \quad x \ll 1, \quad (3)$$

with some exponent θ . With Eq. (3), the hyperscaling argument yields $\rho = (d - \theta)/D_H$, and from $G(r) \propto \int dN P(\mathbf{r}; N)$ one then obtains

$$\eta + D_H = 2 - \theta. \quad (4)$$

Using high-precision data for η and D_H mentioned above, we find $\theta = 0.1965(20)$.

It is instructive to explicitly verify Eq. (3) by simulating $P(r; N)$. In Fig. 1 we present results of such a simulation for the $|\phi|^4$ -model. We plot the value of $P(r, N)N^{d\Delta}$ as a function of r for three different values of N . In view of the self-similarity of $P(r, N)$, the qualitative difference between the cases of $\theta \neq 0$ and $\theta = 0$ is readily seen. In the former case, curves for different values of N should merge for $r/N^\Delta \ll 1$ —and they do in Fig. 1. In the latter case, as $r \rightarrow 0$ one should see a fan of curves with essentially different slopes and a common origin at $r = 0$.




FIG. 1: Distribution of open loops over radii for three different values of N . The Worm algorithm simulation [5] was done for the loop representation (high-temperature expansion) of the 3D lattice $|\phi|^4$ -model with $L = 192^3$ sites at the special critical point with suppressed leading corrections to scaling [3].

One important implication of Eq. (4) in the absence of additional relation between D_H , η and θ , is that the anomalous scaling dimension *can not* be deduced from simulations of closed loops which determine D_H only.

We are grateful to A. Kuklov, Z. Tešanović, and G. Williams for numerous discussions. This work was supported by the National Science Foundation under Grant DMR-0071767.

Nikolay Prokof'ev and Boris Svistunov,
Department of Physics, University of Massachusetts,
Amherst, MA 01003.

PACS numbers: 05.70.Jk, 64.60.Ak, 05.70.Fh

- [1] J. Hove, S. Mo, and A. Sudbø, Phys. Rev. Lett. **85**, 2368 (2000).
- [2] G.A. Williams, Phys. Rev. Lett. **82**, 1201 (1999).
- [3] M. Campostrini *et al.*, Phys. Rev. B, **63**, 214503 (2001).

- [4] M. Wallin *et al.*, Phys. Rev. B **49**, 12115 (1994).
- [5] N. Prokof'ev and B. Svistunov, Phys. Rev. Lett. **87**, 160601 (2001).