

2021

Dental Fluoride Varnish Application During Medical Visits Among Children Who Are Privately Insured

Kimberley H. Geissler PhD
University of Massachusetts Amherst

Andrew W. Dick PhD
RAND, Boston, Massachusetts

Sarah L. Goff PhD
University of Massachusetts Amherst

Christopher Whaley PhD
RAND, Santa Monica, California

Ashley M. Kranz PhD
RAND, Arlington, Virginia

Follow this and additional works at: https://scholarworks.umass.edu/hpp_faculty_pubs



Part of the [Health Policy Commons](#), [Internal Medicine Commons](#), and the [Public Health Education and Promotion Commons](#)

Recommended Citation

Geissler, Kimberley H. PhD; Dick, Andrew W. PhD; Goff, Sarah L. PhD; Whaley, Christopher PhD; and Kranz, Ashley M. PhD, "Dental Fluoride Varnish Application During Medical Visits Among Children Who Are Privately Insured" (2021). *JAMA Network Open*. 13.
<https://doi.org/10.1001/jamanetworkopen.2021.22953>

This Article is brought to you for free and open access by the Health Promotion and Policy at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Health Promotion and Policy Faculty Publication Series by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.



Dental Fluoride Varnish Application During Medical Visits Among Children Who Are Privately Insured

Kimberley H. Geissler, PhD; Andrew W. Dick, PhD; Sarah L. Goff, MD, PhD; Christopher Whaley, PhD; Ashley M. Kranz, PhD

Introduction

Fluoride varnish is effective at reducing tooth decay, which affects nearly a quarter of US children ages 2 to 5 years and more than half of children ages 6 to 8 years.^{1,2} To increase young children's receipt of preventive oral health services, the US Preventive Services Task Force recommends medical providers apply fluoride varnish to young children's teeth during well-child visits through 5 years of age.² Offering fluoride varnish in medical settings may increase young children's receipt of this service because 89% of children younger than 6 years of age had a preventive medical visit in 2019.³ However, fewer than 8% of young Medicaid-enrollees receive fluoride varnish in medical settings,⁴ and no studies have examined fluoride varnish applications during medical visits for children who are privately insured. Studying children who are privately insured is important because coverage of this service without cost-sharing has been mandated since 2015,⁵ and fewer than 1 in 3

+ Supplemental content

Author affiliations and article information are listed at the end of this article.

Table. Characteristics of Well-Child Visits With and Without Fluoride Varnish Applications

Characteristic	Overall (N = 328 661)	Observation, No. (%)		Unadjusted odds of fluoride varnish application, OR (95% CI) ^a
		Visit with fluoride varnish application (N = 15 756)	Visit without fluoride varnish application (N = 312 905)	
Visit included fluoride varnish application	15 756 (4.8)	15 756 (100)	312 905 (0)	NA
Age, y				
1	132 563 (40.3)	7438 (47.2)	125 125 (40.0)	13.6 (8.1-22.9) ^b
2	70 365 (21.4)	5749 (36.5)	64 616 (20.7)	20.3 (13.8-30.0) ^b
3	46 345 (14.1)	2069 (13.1)	44 276 (14.2)	10.7 (7.3-15.5) ^b
4	39 693 (12.1)	327 (2.1)	39 366 (12.6)	1.9 (1.6-2.3) ^b
5	39 695 (12.1)	173 (1.1)	39 522 (12.6)	[Reference]
Sex				
Male	169 001 (51.4)	8129 (51.6)	160 872 (51.4)	[Reference]
Female	159 660 (48.6)	7627 (48.5)	152 033 (48.6)	1.0 (0.95-1.0)
Insurance type				
Preferred provider organization	171 273 (52.1)	8795 (55.8)	162 478 (51.9)	1.2 (0.9-1.7)
Health maintenance organization	84 049 (25.6)	3363 (21.3)	80 686 (25.8)	0.9 (0.7-1.4)
Point of service	68 702 (20.9)	3401 (21.6)	65 301 (20.9)	1.2 (0.9-1.6)
Exclusive provider organization	4637 (1.4)	197 (1.3)	4440 (1.4)	[Reference]
Year of visit				
2016	111 156 (33.8)	4060 (25.8)	107 096 (34.2)	[Reference]
2017	110 892 (33.7)	5563 (35.3)	105 329 (33.7)	1.4 (1.1-1.8) ^b
2018	106 613 (32.4)	6133 (38.9)	100 480 (32.1)	1.6 (1.3-2.1) ^b
State of residence				
Connecticut	134 662 (41.0)	3829 (24.3)	130 833 (41.8)	[Reference]
Maine	80 174 (24.4)	6410 (40.7)	73 764 (23.6)	3.0 (1.5-6.1) ^b
New Hampshire	64 584 (19.7)	1401 (8.9)	63 183 (20.2)	0.8 (0.3-1.8)
Rhode Island	49 241 (15.0)	4116 (26.1)	45 125 (14.4)	3.1 (1.3-7.2) ^b

Abbreviation: OR, odds ratio.

^a OR and 95% CIs from logistic regression models where the unit of observation was a well-child medical visit; models were estimated separately for each of the independent variables of interest and included county-level cluster robust standard errors.

^b Indicates difference in OR is statistically significantly different from the reference group.

Open Access. This is an open access article distributed under the terms of the CC-BY License.

children under 5 years who are privately insured have an annual dental visit.⁶ We used data from 4 states to examine fluoride varnish application rates during well-child medical visits and identify characteristics associated with fluoride varnish receipt.


Methods

This cross-sectional study was approved by RAND's institutional review board, and a waiver of informed consent was granted because of the use of deidentified patient data. This study followed the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) reporting guideline.

We used 2016 to 2018 data for children who are privately insured from all payer-claims databases from Maine, Connecticut, New Hampshire, and Rhode Island. We limited the analytic sample to children ages 1 to 5 years and identified well-child visits using CPT codes 99381-3 and 99391-3. Well-child visit periodicity schedules vary by age; thus, our unit of analysis was the well-child visit. We identified fluoride varnish applications (ie, CPT code 99188 and CDT code D1206) during well-child visits with the same service date.

We calculated descriptive statistics and estimated the unadjusted and adjusted odds of a visit including fluoride varnish using logistic regression; we then calculated regression-adjusted

Figure. Regression-Adjusted Rates of Fluoride Varnish Applications During Well-Child Medical Visits, (N = 328 661)



The figure presents unadjusted and regression-adjusted probabilities. Unadjusted probabilities are derived from logistic regression models predicting fluoride varnish applications during well-child medical visits. Models are estimated separately for each of the independent variables of interest (ie, age, state, and year). Regression-adjusted results are from a model controlling for child sex, age, insurance type, visit year, and state. In all models, standard errors are clustered at the county level, and the delta method was used to calculate standard errors for predicted probabilities. We calculated probabilities of fluoride varnish application during a well-child medical visit for each

variable category of interest using model estimates to compute the mean of predicted probabilities for the entire sample after setting the covariate of interest (eg, Connecticut) while keeping all other covariates at their observed values.

^a Indicates difference in predicted probability is statistically significantly different from the reference group (child age = 5 years, state = Connecticut, year = 2016) at the 5% level.

probabilities of fluoride application (eMethods in the Supplement). Tests were 2-tailed, statistical significance was set at $P < .05$, and we used county-level cluster robust standard errors. Data analyses were performed using SAS Version 9.4 and Stata-MP version 16.1 (StataCorp). Analysis was performed during November 2020 to March 2021.

Results

The sample included 328 661 well-child visits for children aged 1 to 5 years (169 001 [51.4%] male; 132 563 [40.3%] were 1-year-olds). Of the visits, 134 662 (41.0%) occurred in Connecticut and 15 756 (4.8%) included fluoride varnish applications (Table). Fluoride varnish was more common among visits for younger children, as illustrated by unadjusted and regression-adjusted probabilities (Figure). A 2-year-old was 7.7 percentage points (pp) (95% CI, 5.9 pp-9.4 pp) more likely to receive fluoride varnish than a 5-year-old. From 2016 to 2018, the regression-adjusted probability of fluoride varnish application increased from 3.6% (95% CI, 2.8%-4.4%) to 5.8% (95% CI, 4.5%-7.1%). Fluoride varnish applications were most common in Rhode Island, with a regression-adjusted probability of 8.7% (95% CI, 5.1%-12.4%). New Hampshire had a lower rate, with a regression-adjusted probability of 2.2% (95% CI, 1.2%-3.3%) of a visit including fluoride varnish.

Discussion

Despite mandatory insurance coverage for fluoride varnish applications in medical settings, we found fewer than 5% of well-child visits for privately insured young children included this service, suggesting efforts are needed to increase pediatric medical providers' delivery of fluoride varnish. Young children have few dental visits⁶ and delivering fluoride varnish in medical settings can increase access to preventive oral health services. Although this study was limited to 4 states, these data included a variety of private insurers and clinicians. Differences in fluoride varnish applications across states may be driven by variation in access to dentists and Medicaid payment policy. This study was the first to assess delivery of an evidence-based service for children who are privately insured, which is recommended by the US Preventive Service Task Force and the American Academy of Pediatrics. Although increases over time were encouraging, very low rates of fluoride varnish in medical settings suggest substantial expansion of this service in medical settings is critical for improving children's oral health and overall well-being.

ARTICLE INFORMATION

Accepted for Publication: June 24, 2021.

Published: August 30, 2021. doi:10.1001/jamanetworkopen.2021.22953

Open Access: This is an open access article distributed under the terms of the [CC-BY License](#). © 2021 Geissler KH et al. *JAMA Network Open*.

Corresponding Author: Ashley M. Kranz, PhD, 1200 S Hayes St, Arlington, VA 22202-5050 (akranz@rand.org).

Author Affiliations: University of Massachusetts Amherst School of Public Health and Health Sciences, Amherst (Geissler, Goff); RAND, Boston, Massachusetts (Dick); RAND, Santa Monica, California (Whaley); RAND, Arlington, Virginia (Kranz).

Author Contributions: Dr Kranz had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: Geissler, Dick, Kranz.

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: Geissler, Kranz.

Critical revision of the manuscript for important intellectual content: Dick, Goff, Whaley, Kranz.

Statistical analysis: Geissler, Dick, Kranz.

Obtained funding: Dick, Kranz.

Administrative, technical, or material support: Geissler, Whaley.

Supervision: Whaley, Kranz.

Conflict of Interest Disclosures: Dr Geissler reported receiving grants from the National Institute of Dental and Craniofacial (NIDCR) Research during the conduct of the study. Dr Whaley reported receiving grants from the National Institute on Aging KO1 AGO61274 and grants from the Robert Wood Johnson Foundation during the conduct of the study. Dr Kranz reported receiving grants from NIDCR during the conduct of the study. No other disclosures were reported.

Funding/Support: This research was supported by grant R01 DE028530-01A1 from the National Institute of Dental and Craniofacial Research.

Role of the Funder/Sponsor: The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Disclaimer: The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Dental and Craniofacial Research or the National Institutes of Health.

Additional Contributions: The authors extend thanks to Grace Gahlon (RAND Corporation), Brenna O'Neill (RAND Corporation), and Ayachi Sharma (University of Massachusetts Amherst) for assistance reviewing the literature, statistical programming, and research assistance, respectively. These contributors were compensated for their time.

REFERENCES

1. Dye BA, Thornton-Evans G, Li X, Iafolla TJ. Dental caries and sealant prevalence in children and adolescents in the United States, 2011-2012. *NCHS Data Brief*. 2015;(191):1-8.
2. Moyer VA; US Preventive Services Task Force. Prevention of dental caries in children from birth through age 5 years: US Preventive Services Task Force recommendation statement. *Pediatrics*. 2014;133(6):1102-1111. doi:10.1542/peds.2014-0483
3. Child and Adolescent Health Measurement Initiative. 2019 National Survey of Children's Health data query. U.S. Department of Health and Human Services, Health Resources and Services Administration (HRSA), Maternal and Child Health Bureau (MCHB). Published 2021. Accessed June 14, 2021. <https://www.childhealthdata.org>
4. Geiger CK, Kranz AM, Dick AW, Duffy E, Sorbero M, Stein BD. Delivery of preventive oral health services by rurality: a cross-sectional analysis. *J Rural Health*. 2019;35(1):3-11. doi:10.1111/jrh.12340
5. HealthCare.Gov. Preventive care benefits for children US Center for Medicare & Medicaid Services. Published 2021. Accessed January 21, 2021. <https://www.healthcare.gov/preventive-care-children/>
6. Medicaid and CHIP Payment and Access Commission. Medicaid Access in Brief: Children's Dental Services. Published 2016. Accessed April 15, 2021. <https://www.macpac.gov/wp-content/uploads/2016/06/Medicaid-Access-in-Brief-Childrens-Dental-Services.pdf>

SUPPLEMENT.

eMethods.