2005

Directional cell expansion - turning toward actin

A Bannigan

TI Baskin

Follow this and additional works at: https://scholarworks.umass.edu/biology_faculty_pubs

Part of the [Biology Commons](#)

Recommended Citation

Bannigan, A and Baskin, TI, "Directional cell expansion - turning toward actin" (2005). *Current Opinion in Plant Biology*. 19.

<https://doi.org/10.1016/j.pbi.2005.09.002>

This Article is brought to you for free and open access by the Biology at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Biology Department Faculty Publication Series by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

Directional cell expansion — turning toward actin

Alex Bannigan and Tobias I Baskin

Significant recent progress toward understanding directional expansion in diffusely growing plant cells concerns actin. Tools for imaging actin, including both live-cell reporters and fixation protocols, have been improved. Proteins that interact with actin have been identified and their functions probed biochemically and genetically. Specifically, members of the actin-related protein2/3 (ARP2/3) complex and the Wiskott–Aldrich syndrome Verprolin-homologous (WAVE) complex have been identified. These proteins have salient functions in shaping trichomes and leaf pavement cells. Additionally, two targets of a *rho*-of-plants (ROP) G-protein have been discovered that exert opposing regulatory action on actin and microtubules, a pathway that appears to be responsible for establishing the undulating shapes of pavement cells. Finally, several mutants of the *fragile fiber* class have revealed a link between actin organization, cell wall synthesis, and phosphoinositol signaling.

Addresses

Biology Department, University of Massachusetts, Amherst,
Massachusetts 01003, USA

Corresponding author: Baskin, Tobias I (Baskin@bio.umass.edu)

Introduction

In plants, cell expansion is controlled by turgor pressure, cell wall properties, and the cytoskeleton. Traditionally, microtubules have been implicated in the direction of expansion, probably by orienting newly synthesized cellulose microfibrils [1,2]. By contrast, actin has been implicated in the rate (or overall extent) of expansion [3–6], probably through delivery of Golgi vesicles that contain membrane and wall materials to the site of growth. This traditional distinction is well illustrated by tip-growing cells, such as pollen tubes. Actin plays a primary role in the expansion of these cells whereas the role of microtubules is secondary [7–10]; consistently, the tips of these cells expands approximately isotropically (i.e. at the same rate in all directions) [11].

However, actin has recently emerged as an important contributor to the directionality of expansion in diffusely

growing cells, partly through collaboration with microtubules [6,12,13••]. There have been a wealth of papers revealing a role for actin and related proteins in directional cell expansion, papers that are expanding our limited understanding of actin's role in plant growth. This review focuses on the cooperation of actin and microtubules in shaping diffusely growing cells.

Advances in imaging actin

The improvement of imaging techniques in the past few years is a breakthrough in the study of actin. Actin is notoriously problematic to preserve with chemical fixation [12,14]. In animal cells, reference images have been produced by tagging actin, but this approach has yet to be successful for plant actin, whether tagged biochemically (e.g. with rhodamine) or genetically (e.g. with green fluorescent protein [GFP]). Recently, success in plants has been achieved by tagging actin-binding proteins.

Perhaps the first tagged actin-binding protein to be used in plants was mouse talin (GFP–mTalin) [15]. Recently, images of maize leaf cells expressing GFP–mTalin were shown to resemble material that was fixed and stained with phalloidin [16]. However, expression of GFP–mTalin can alter morphology and disrupt actin severely [17,18•,19]. Generally less disturbance to actin and plant growth has been achieved when actin-binding domains (ABDs) from *Arabidopsis* fimbrin are used for tagging. Full-length fimbrin–GFP gives bright fluorescence but reveals few filaments; however, a construct with a part of the protein that includes the binding domains, ABD1/2–GFP, gives dimmer fluorescence but greater detail [19]. Another construct that includes part of the fimbrin protein, GFP–fimbrinABD2, gives striking images of dense actin networks in *Arabidopsis* root, hypocotyl, and leaf epidermal cells, and permits studies of actin dynamics [18•]. Sheahan *et al.* [18•] noted a correspondence between the actin organization seen with this protein construct and that in images of carefully fixed cells. The complexity of the actin cytoskeleton is such that no single actin-binding protein can be expected to stain every bundle and filament, but the congruence of a live-cell reporter and fixation in a given investigation offers assurance that an approximation to the state of the cytoskeleton in the living cell has been reached.

As an alternative approach, Lovy-Wheeler and colleagues [14] improved fixation techniques, specifically for pollen tubes. They took advantage of the fact that cryofixation and freeze substitution provide unmatched structural preservation of single cells and used images from cryo-fixed pollen tubes as a reference to develop a protocol for

chemical fixation. The advantages of chemical fixation are that it requires no special apparatus, provides higher throughput, and can be applied immediately following live-cell imaging. The resulting protocol for chemical fixation led to images of actin that were all but indistinguishable from the cryofixed images, which, importantly, are somewhat different from images of living pollen tubes expressing fluorescent actin-binding proteins (K Wilsen *et al.*, unpublished).

Coordination of microtubules and actin in cells with complex shapes

In general, actin and microtubule arrays in the cell cortex are considered to have a close association [20], and it has been reported frequently that perturbing one can affect the stability of the other [21–24]. Recently, chemical disruption of actin was shown to disrupt severely the phragmoplast in synchronized tobacco tissue-culture cells and to prevent microtubule reorganization at the mitosis–interphase transition [25]. Also, a kinesin that was identified in cotton binds both microtubules and actin filaments and contains an actin-binding calponin-homology domain [26[•]]. Kinesins that have calponin domains form a small clade that is unique to plants; *Arabidopsis* has seven members, offering scope for interactions between actin and microtubules.

The interlinked status of actin and microtubules has been incisively revealed in pavement cells and trichomes. Pavement cells are characterized by lobed walls that fit together like a jigsaw puzzle across the surface of the leaf; the lobes can be gentle sinusoidal undulations, as in maize, or deeply curved and irregular, as in *Arabidopsis*. Trichomes grow out from the surface of the leaf to form spike-like cells that can be straight or branched. Trichomes in *Arabidopsis* have recently been demonstrated to grow diffusely rather than by tip growth [27]. The morphogenesis of pavement cells and trichomes has been comprehensively reviewed by Smith and Oppenheimer [13^{••}] and that of pavement cells and lobed mesophyll by Panteris and Galatis [28^{••}].

The prevailing theory to explain how the interlocking pattern of pavement cell lobes is formed has been that bands of microtubules cause localized wall thickenings, restricting cell wall expansion in the ‘neck’ region and forcing the intervening thinner wall regions to stretch under turgor pressure [6,28^{••}]. However, the inability of *Brick1*, a maize mutant in an actin-regulating protein, to form lobes despite the apparently correct formation of microtubule bands highlighted the importance of actin in lobe formation [16]. It is now clear that both actin and microtubules are crucial for the elaboration of lobes.

Likewise, building a trichome appears to require coordinated action from microtubules and actin. Mutants in any of a set of genes that encode actin-regulating proteins

have distorted trichomes ([13^{••}] and references below); however, cortical microtubules are aberrant in many of these mutants [27,29]. When these mutations are crossed into *zwichel*, a mutant that has a defect in a microtubule-dependent motor protein, the double mutants have a phenotype that indicates a synthetic interaction [27]. Additionally, endoplasmic microtubules, although less frequently scored than cortical microtubules, are abnormally organized and resistant to chemical depolymerization in at least one of the distorted trichome mutant backgrounds [30[•]]. Saedler *et al.* [30[•]] suggest that during trichome formation the cortical actin becomes more dynamic at a site of expansion, allowing vesicle delivery and wall bulging, while microtubules restrict the extent of bulging by hemming in the spread of the increased actin turnover. Intriguingly, localized control of actin dynamics is thought to be the basis of the formation of protrusions in moving animal cells [31,32[•]].

Modeling and remodeling actin: actin-binding proteins emerge from the genomic shadows

Actin behavior is governed by a suite of actin-associated protein complexes. Recent characterization of components of the actin-regulating ARP2/3 and WAVE complexes, as well as of the ROP-GTPase signaling pathway, has uncovered a complex web of control that potentially allows signal transduction to, and co-coordinated reorganization of, the actin cytoskeleton. These components are reviewed, in the context of cell polarity, by Xu and Scheres, this issue.

ARP2/3 complex

The ARP2/3 complex is pivotal for regulating actin polymerization in animals [32[•]]. When activated, it enhances the nucleation and polymerization of filamentous actin. It also binds to existing actin filaments, initiating branches and leading to the formation of dendritic actin arrays. Until recently, the ARP2/3 complex was unknown in plants.

Homologs of all seven subunits of the ARP2/3 complex have now been identified in plants on the basis of homology searches and loss-of-function mutants [30[•],33–35]. Mutations in the ARP2/3 complex of *Arabidopsis* cause abnormal actin organization and deformation of trichomes, leaf epidermal pavement cells and some hypocotyl epidermal cells, but overall plant morphology is scarcely affected. In these mutants, trichome branches are underdeveloped or twisted, the trichome stalk is unusually thick, and actin organization is compromised [30[•],32[•],34]. Along with the trichomes, leaf pavement cells and hypocotyl cells are affected: pavement cells are less lobed in the mutants than in wildtype, and some hypocotyl epidermal cells become separated at their end walls under rapid growth conditions [33,35].

Surprisingly, ARP2/3 mutations in *Arabidopsis* affect neither root hairs nor pollen tubes nor the majority of

diffuse growing cells in the plant, suggesting that only select cell types take advantage of the dendritic mode of assembly that is supported by ARP2/3. This is in contrast to animals, in which loss-of-function mutants have extreme, sometimes lethal, phenotypes (summarized by [32]). Even in the moss *Physcomitrella patens*, loss of one of the ARP2/3 subunits completely blocks the tip-growth of caulonemal cells and thus prevents bud formation and the development of leafy gametophores [36]. Several authors have noted that ARP2/3 mutants have less fine cortical actin and more thick endoplasmic actin cables, and have suggested that fine cortical actin plays a role in cell expansion [16,34,37]. It will presumably take ultrastructural investigation to reveal how fine actin that is established by ARP2/3 differs from that established by other regulators, such as formin.

WAVE complex

Another complex that is involved in actin regulation is the WAVE complex, members of which have also been recently identified in plants [29,37]. In animal cells, WAVE proteins regulate ARP2/3 activity and are also known to interact with the G-protein RAC. Hence, the identification of WAVE-like proteins in plants provides a putative link between G-protein-dependent signal transduction and cytoskeletal organization [37]. Mutations of the members of the plant WAVE complex that have been isolated to date result in phenotypes similar to those of ARP2/3 mutants [29,37], confirming that they act in the same actin-regulating pathway.

BRICK1 is a WAVE complex member that has been identified in maize and that is homologous to the human WAVE component hematopoietic stem progenitor clone 300 (HSPC300) [16]. BRICK1 revealed the involvement of actin in the formation of pavement cell lobes. Furthermore, lines that are mosaic for *Brick1* showed that wild-type BRICK1 can induce lobe formation in neighboring mutant cells, effective over a distance of two to three cells from the site of transcription. Intercellular interactions are presumably necessary to co-ordinate the interdigitating growth that is typical of pavement cells, and yet BRICK1 is the only cell-shaping protein known to act non cell-autonomously.

ROP-GTPase signaling pathway in plants

In animal cells, the *rho*-GTPase signaling pathway coordinates the cytoskeleton to perform various functions, including cell migration [38]. Previously, it has been established in plants that homologs of *rho*-GTPases, called *rho*-of-plants (ROP), participate in organizing both cortical actin and microtubules ([39,40]; Xu and Scheres, this issue). This year, Fu *et al.* [41] identified two CRIB-motif-containing ROP target proteins, RIC1 and RIC4, that mediate microtubule and actin organization via counteracting pathways. Furthermore, they proposed a model that accounts for the regulation of both actin and

microtubules to generate the convoluted shapes of pavement cells (Figure 1).

According to this model, RIC1 helps to organize microtubules into parallel arrays in the cell 'necks' and thus reins in expansion, whereas RIC4 mediates the polymerization of fine cortical actin filaments in the lobe tips and thus promotes outward growth. What's more, the RIC1 and RIC4 pathways negatively regulate each other to ensure well demarked lobe and neck regions. The discovery of these counterpoised RICs is fascinating because, although numerous microtubule- and actin-associated proteins are known, few cytoskeletal organizers, let alone pathways that coordinate the regulation of microtubules and actin, have been identified.

The interaction between the ROP G-proteins and the downstream RICs presumably depends on a guanine-nucleotide exchange facilitator (GEF) to activate the ROP. In plants, the identification of ROP-GEFs has been hindered by a lack of sequence similarity to known animal GEFs; only this year has their presence in plants been confirmed on the basis of biochemical assays [42]. ROP-GEFs are represented in *Arabidopsis* by a family of

Figure 1

Cartoon of pavement cell lobe formation that is based on the model by Fu *et al.* [41]. When ROP GTPase is turned on, it activates RIC4, inducing the formation of fine cortical actin filaments and lobe expansion and concomitantly suppressing RIC1 activity. ROP has limited mobility, so its activity is limited spatially. RIC1 (possibly activated by a signal from the neighboring cell) localizes to microtubules and organizes them into parallel bundles, restricting expansion locally. The polymerization status of microtubules in turn affects local ROP activation. RIC1-organized microtubules sequester and inactivate ROP, preventing its interaction with RIC4; whereas depolymerized microtubules release ROP. This ensures that active RIC4 and RIC1 are spatially separated, allowing the formation of distinct lobes and troughs at the pavement cell margin.

at least 14 highly related proteins, but specifically interacting pairs of ROP and GEF partners have yet to be found. The presence of more than 14 ROP-GEFs in *Arabidopsis* is suggested by the existence of a protein called SPIKE, which was identified from one of the many trichome mutants. SPIKE is also necessary for pavement cell lobe formation and shows characteristics of a *rho*-GEF [43]. In animal cells, GEF-H1 is believed to be activated by microtubule depolymerization [44]; if one of the plant GEFs behaves similarly, this would reinforce the model of Fu *et al.* [41^{••}] by enhancing ROP activity in the lobes, where microtubules are sparse.

Other examples of actin's role in directional cell expansion

In addition to cells that combine overall diffuse growth with localized expansion to create complex cell shapes, cells that grow principally along one axis also need actin. When actin is chemically disrupted, diffuse growth usually becomes less anisotropic [13^{••}]. Because actin and microtubules often have different arrangements at the expanding and non-expanding walls of a cell, Fu *et al.* [41^{••}] suggest that the ROP GTPase signaling pathway coordinates cytoskeletal organization in many, if not all, cell types. Consistently, transgenic constructs that render the ROP constitutively active disrupt actin organization and expansion, causing deformed organs, loss of pavement cell lobes, and roughly isotropic growth in mesophyll cells [45[•]].

Other examples of actin's involvement in cell wall synthesis and cell elongation come from members of the *fragile fiber* (*fra*) family of mutants [46–48]. Cell walls in the fiber cells of *fra4* (also known as *root hair defective 3*) are abnormally thin, have altered composition and reduced mechanical strength, and have reduced cell elongation in all organs [46]. Actin in *fra4* is clumped into thick cables and has less branching than the actin of wildtype plants. *FRA4* encodes a novel protein that has putative GTP-binding motifs [46]. *FRA3* encodes an inositol polyphosphate phosphatase and *FRA7* encodes a so-called 'suppressor of actin (SAC) domain' phosphatase, which is also active against inositol polyphosphates [47,48]. Both also appear to be necessary for normal actin organization and secondary cell wall formation, with *fra7* mutant plants having reduced expansion in most, if not all, organs [48]. The cell wall defects of these three *fra* mutants are plausibly caused by reduced vesicle delivery to the plasma membrane. In support of this, cytochalasin D, which disrupts actin, also causes reduced wall thickness in fiber cells [46]. Together, the *fra3*, *fra4* and *fra7* mutants link phosphoinositide metabolism, actin organization, and cell wall synthesis. Finally, the link between actin microfilaments, secretion, and directional expansion is underscored by *katamari1*, a mutant that has reduced elongation, and disrupted actin and endoplasmic reticulum. The protein that is defective in this mutant is a

Golgi-resident, glycosyl-transferase, previously identified as MUR3 [49].

Conclusions

Less than a decade ago, actin was all but ignored in treatments of the expansion of diffusely growing cells. It has now become clear that actin can promote outgrowth in plant cells, as seen in pavement cells, and that in cooperation with microtubules, it plays a major role in the construction of complex cell shapes and in expansion. We have only just begun, however, to discover the diverse roles of actin and the myriad signaling pathways that govern its organization and function. The recent work discussed here, focusing on signaling molecules and pathways, has opened up a field of study that we are confident will reveal mechanisms that control the plant cytoskeleton and directional cell expansion.

Acknowledgement

Work in the authors' laboratory on directional cell expansion is supported by a grant from US Department of Energy (award No. 03ER15421), which does not constitute endorsement of that department of the views expressed herein.

References and recommended reading

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Baskin TI: **On the alignment of cellulose microfibrils by cortical microtubules: a review and a model.** *Protoplasma* 2001, **215**:150-171.
2. Wasteneys GO: **Progress in understanding the role of microtubules in plant cells.** *Curr Opin Plant Biol* 2004, **7**:651-660.
3. Baskin TI, Bivens NJ: **Stimulation of radial expansion in *Arabidopsis* roots by inhibitors of actomyosin and vesicle secretion but not by various inhibitors of metabolism.** *Planta* 1995, **197**:514-521.
4. Ramachandran S, Christensen HEM, Ishimaru Y, Dong CH, Chao-Ming W, Cleary AL, Chua NH: **Profilin plays a role in cell elongation, cell shape maintenance, and flowering in *Arabidopsis*.** *Plant Physiol* 2000, **124**:1637-1647.
5. Baluska F, Jasik J, Edelmann HG, Salajova T, Volkmann D: **Latrunculin B-induced plant dwarfism: plant cell elongation is actin-dependent.** *Dev Biol* 2001, **231**:113-124.
6. Smith LG: **Cytoskeletal control of plant cell shape: getting the fine points.** *Curr Opin Plant Biol* 2003, **6**:63-73.
7. Hepler PK, Vidali L, Cheung AY: **Polarized cell growth in higher plants.** *Annu Rev Cell Dev Biol* 2001, **17**:159-187.
8. Ringli C, Baumberger N, Diet A, Frey B, Keller B: **ACTIN2 is essential for bulge site selection and tip growth during root hair development of *Arabidopsis*.** *Plant Physiol* 2002, **129**:1464-1472.
9. Samaj J, Ovecka M, Hlavacka A, Lecourieux F, Meskiene I, Lichtscheidl I, Lenart P, Slaj J, Volkmann D, Borge L *et al.*: **Involvement of the mitogen-activated protein kinase SIMK in regulation of root hair tip growth.** *EMBO J* 2002, **21**:3296-3306.
10. Ketelaar T, de Ruijter NCA, Emons AMC: **Unstable F-actin specifies the area and microtubule direction of cell expansion in *Arabidopsis* root hairs.** *Plant Cell* 2003, **15**:285-292.
11. Baskin TI: **Anisotropic expansion of the plant cell wall.** *Annu Rev Dev Cell Biol* 2005, **21**:203-222.

12. Wasteneys GO, Galway ME: **Remodeling the cytoskeleton for growth and form — an overview with some new views.** *Annu Rev Plant Biol* 2003, **54**:691-722.
13. Smith LG, Oppenheimer DG: **Spatial control of cell expansion • by the plant cytoskeleton.** *Annu Rev Cell Dev Biol* 2005, **21**:271-295.
An exhaustive review of the role of the cytoskeleton in plant cell shape formation, covering trichome and pavement cell morphogenesis, pollen tube growth, actin, microtubules and the signaling pathways that mediate their interaction.
14. Lovy-Wheeler A, Wilsen KL, Baskin TI, Hepler PK: **Enhanced fixation reveals the apical cortical fringe of actin filaments as a consistent feature of the pollen tube.** *Planta* 2005, **221**:95-104.
15. Kost B, Spielhofer P, Chua N-H: **A GFP-mouse talin fusion protein labels plant actin filaments *in vivo* and visualizes the actin cytoskeleton in growing pollen tubes.** *Plant J* 1998, **16**:393-401.
16. Frank MJ, Cartwright HN, Smith LG: **Three *BRICK* genes have distinct functions in a common pathway promoting polarized cell division and cell morphogenesis in the maize leaf epidermis.** *Development* 2003, **130**:753-762.
17. Ketelaar T, Anthony RG, Hussey PJ: **Green fluorescent protein-mTalin causes defects in actin organization and cell expansion in *Arabidopsis* and inhibits actin depolymerizing factor's actin depolymerizing activity *in vivo*.** *Plant Physiol* 2004, **136**:3990-3998.
18. Sheahan MB, Staiger CJ, Rose RJ, McCurdy DW: **A green fluorescent protein fusion to actin-binding domain 2 of *Arabidopsis* fimbrin highlights new features of a dynamic actin cytoskeleton in live plant cells.** *Plant Physiol* 2004, **136**:3968-3978.
The authors produced beautiful images of actin in root, hypocotyl, pavement, trichome and guard cells using *Arabidopsis* fimbrin binding domain 2 fused to GFP. They also compare several constructs and conclude that fABD2-GFP gives superior actin visualization with fewer morphological side effects when compared to GFP-mTalin.
19. Wang Y-S, Motes CM, Mohamawari DR, Blancaflor EB: **Green fluorescent protein fusions to *Arabidopsis* fimbrin 1 for spatio-temporal imaging of F-actin dynamics in roots.** *Cell Motil Cytoskeleton* 2004, **59**:79-93.
20. Collings DA, Allen NS: **Cortical actin interacts with the plasma membrane and microtubules.** In *Actin: a Dynamic Framework for Multiple Plant Cell Functions*. Edited by Staiger CJ, Baluska F, Volkmann D, Barlow PW. Kluwer Academic; 2000:145-163.
21. Chu B, Kerr GP, Carter JV: **Stabilizing microtubules with taxol increases microfilament stability during freezing of rye root tips.** *Plant Cell Environ* 1993, **16**:883-889.
22. Collings DA, Wasteneys GE, Williamson RE: **Actin-microtubule interactions in the alga *Nitella*: analysis of the mechanism by which microtubule depolymerisation potentiates cytochalasin's effect on streaming.** *Protoplasma* 1996, **191**:178-190.
23. Tominaga M, Morita K, Sonobe S, Yokota E, Shimmen T: **Microtubules regulate the organization of actin filaments at the cortical region in root hair cells of *Hydrocharis*.** *Protoplasma* 1997, **199**:83-92.
24. Blancaflor EB: **Cortical actin filaments potentially interact with cortical microtubules in regulating polarity of cell expansion in primary roots of maize (*Zea mays* L.).** *J Plant Growth Regul* 2000, **19**:406-414.
25. Yoneda A, Akatsuka M, Kumagai F, Hasezawa S: **Disruption of actin microfilaments causes cortical microtubule disorganization and extra phragmoplast formation at the M/G₁ interface in synchronized tobacco cells.** *Plant Cell Physiol* 2004, **45**:761-769.
26. Preuss ML, Kovar DR, Lee Y-RJ, Staiger CJ, Delmer DP, Liu B: **A plant-specific kinesin binds to actin microfilaments and interacts with cortical microtubules in cotton fibers.** *Plant Physiol* 2004, **136**:3945-3955.
The authors describe a novel cotton kinesin, unique to plants, that binds actin *in vitro* and decorates microtubules and, occasionally, fine cortical actin *in vivo*.
27. Schwab B, Mathur J, Saedler R, Schwartz H, Frey B, Scheidegger C, Hülskamp M: **Regulation of cell expansion by the *DISTORTED* genes in *Arabidopsis thaliana*: actin controls the spatial organization of microtubules.** *Mol Genet Genomics* 2003, **269**:350-360.
28. Panteris E, Galatis B: **The morphogenesis of lobed plant cells in the mesophyll and epidermis: organization and distinct roles of cortical microtubules and actin filaments.** *New Phytol* 2005, **167**:721-732.
A masterful review of lobed mesophyll and pavement cell morphogenesis, with an ultrastructural emphasis.
29. Zhang X, Dyachok J, Krishnakumar S, Smith LG, Oppenheimer DG: ***IRREGULAR TRICHOME BRANCH1* in *Arabidopsis* encodes a plant homolog of the actin-related protein2/3 complex activator Scar/WAVE that regulates actin and microtubule organization.** *Plant Cell* 2005, **17**:2314-2326.
30. Saedler R, Mathur N, Srinivas BP, Kernebeck B, Hülskamp M, Mathur J: **Actin control over microtubules suggested by *DISTORTED2* encoding the *Arabidopsis* ARPC2 subunit homolog.** *Plant Cell Physiol* 2004, **45**:813-822.
The work described in this paper provides evidence for actin-microtubule interaction and interdependence by showing that endoplasmic microtubules are abnormally organized and unusually stable in the ARP2/3 mutant *distorted2*. The authors propose that the abnormal microtubule organization is the probable cause of the distorted trichome shapes.
31. Rafelski SM, Theriot JA: **Crawling toward a unified model of cell mobility: spatial and temporal regulation of actin dynamics.** *Annu Rev Biochem* 2004, **73**:209-239.
32. Mathur J: **The ARP2/3 complex: giving plant cells a leading edge.** *Bioessays* 2005, **27**:377-387.
A detailed review of the ARP2/3 complex in plants and of mutants of the *distorted* family. The author mentions actin-microtubule interdependence and possible intercellular influences on the cytoskeleton and cell shape formation.
33. Mathur J, Mathur N, Kirik V, Kernebeck B, Srinivas BP, Hülskamp M: ***Arabidopsis* *CROOKED* encodes for the smallest subunit of the ARP2/3 complex and controls cell shape by region specific fine F-actin formation.** *Development* 2003, **130**:3137-3146.
34. Li S, Blanchoin L, Yang Z, Lord EM: **The putative *Arabidopsis* ARP2/3 complex controls leaf cell morphogenesis.** *Plant Physiol* 2003, **132**:2034-2044.
35. Mathur J, Mathur N, Kernebeck B, Hülskamp M: **Mutations in actin-related proteins 2 and 3 affect cell shape development in *Arabidopsis*.** *Plant Cell* 2003, **15**:1632-1645.
36. Harries PA, Pan A, Quatrano RS: **Actin-related protein2/3 complex component ARPC1 is required for proper cell morphogenesis and polarized cell growth in *Physcomitrella patens*.** *Plant Cell* 2005, **17**:2327-2339.
37. Brembu T, Winge P, Seem M, Bones AM: ***NAPP* and *PIRP* • encode subunits of a putative WAVE regulatory protein complex involved in plant cell morphogenesis.** *Plant Cell* 2004, **16**:2335-2349.
The authors show that the WAVE complex is at least partly conserved in plants, and that its subunits are important for cell shape formation in trichomes and pavement cells. The complex acts in the same pathway as the ARP2/3 complex to control actin branching and organization.
38. Etienne-Manneville S, Etienne-Manneville S: **Actin and microtubules in cell motility: which one is in control?** *Traffic* 2004, **5**:470-477.
39. Molendijk AJ, Bischoff F, Rajendrakumar CSV, Friml J, Gilroy S, Palme K: ***Arabidopsis thaliana* ROP GTPases are localized to tips of root hairs and control polar growth.** *EMBO J* 2001, **20**:2779-2788.
40. Fu Y, Li H, Yang Z: **The ROP2 GTPase controls the formation of cortical fine F-actin and the early phase of directional cell expansion during *Arabidopsis* organogenesis.** *Plant Cell* 2002, **14**:777-794.
41. Fu Y, Gu Y, Zheng Z, Wasteneys G, Yang Z: ***Arabidopsis* interdigitating cell growth requires two antagonistic pathways with opposing action on cell morphogenesis.** *Cell* 2005, **120**:687-700.

This fascinating paper demonstrates the existence of a conserved Rho-GTPase signaling pathway in plants that acts via RIC proteins to coordinate the actin and microtubule cytoskeletons. The two counteracting pathways negatively regulate each other locally, creating defined sub-cellular regions of expansion and restriction.

42. Berken A, Thomas C, Wittinghofer A: **A new family of RhoGEFs •• activates the ROP molecular switch in plants.** *Nature* 2005, **436**:1176-1180.

This paper describes the identification of the first confirmed ROP-specific GEFs in plants, which have no sequence homology to non-plant proteins. GEF activity was measured by the release of a fluorescent GDP analog from ROP4. The active region that is conserved in related proteins was named PRONE.

43. Qiu J-L, Jilk R, Marks MD, Szymanski DB: **The *Arabidopsis* SPIKE1 gene is required for normal cell shape control and tissue development.** *Plant Cell* 2002, **14**:101-118.
44. Krendel M, Zenke FT, Bokoch GM: **Nucleotide exchange factor GEF-H1 mediates cross-talk between microtubules and the actin cytoskeleton.** *Nat Cell Biol* 2002, **4**:294-301.
45. Bloch D, Lavy M, Efrat Y, Efroni I, Bracha-Drori K, Abu-Abied M, Sadot E, Yalovsky S: **Ectopic expression of an activated RAC in**

***Arabidopsis* disrupts membrane cycling.** *Mol Biol Cell* 2005, **16**:1913-1927.

The authors show that constitutively active ROP in *Arabidopsis* disrupts actin and causes deformed root hairs and pavement cells, as well as isotropic expansion of mesophyll cells.

46. Hu Y, Zhong R, Morrison WH, Ye Z-H: **The *Arabidopsis* RHD3 gene is required for cell wall biosynthesis and actin organization.** *Planta* 2003, **217**:912-921.
47. Zhong R, Burk DH, Morrison WH, Ye Z-H: **FRAGILE FIBRE3, an *Arabidopsis* gene encoding a type II inositol polyphosphate 5-phosphatase, is required for secondary wall synthesis and actin organization in fiber cells.** *Plant Cell* 2004, **16**:3242-3259.
48. Zhong R, Burk DH, Nairn CJ, Wood-Jones A, Morrison WH, Ye Z-H: **Mutation of AtSAC1, an *Arabidopsis* SAC domain phosphoinositide phosphatase, causes alterations in cell morphogenesis, cell wall synthesis and actin organization.** *Plant Cell* 2005, **17**:1449-1466.
49. Tamura K, Shimada T, Kondo M, Nishimura M, Hara-Nishimura I: **KATAMARI1/MURUS3 is a novel Golgi membrane protein that is required for endomembrane organization in *Arabidopsis*.** *Plant Cell* 2005, **17**:1764-1776.