

University of Massachusetts Amherst

ScholarWorks@UMass Amherst

Food Science Department Faculty Publication
Series

Food Science

2021

Genome Sequence of *Aspergillus aculeatinus* IC_8, Isolated from an Indoor Air Sample of an Urban Housing Complex in Abidjan, Ivory Coast

Shu Zhao

David Koffi

Jean-Paul Latge

Karidia Sylla

John G. Gibbons

Follow this and additional works at: https://scholarworks.umass.edu/foodsci_faculty_pubs



Part of the [Microbiology Commons](#)



Genome Sequence of *Aspergillus aculeatinus* IC_8, Isolated from an Indoor Air Sample of an Urban Housing Complex in Abidjan, Ivory Coast

Shu Zhao,^{a,b} David Koffi,^c Jean-Paul Latge,^d Karidia Sylla,^c  John G. Gibbons^{a,b,e}

^aDepartment of Food Science, University of Massachusetts, Amherst, Massachusetts, USA

^bMolecular and Cellular Biology Graduate Program, University of Massachusetts, Amherst, Massachusetts, USA

^cParasitology and Mycology Department, Institut Pasteur de Côte d'Ivoire, Abidjan, Ivory Coast

^dAspergillus Unit, Institut Pasteur, Paris, France

^eOrganismic & Evolutionary Biology Graduate Program, University of Massachusetts, Amherst, Massachusetts, USA

ABSTRACT *Aspergillus aculeatinus* is an industrially important species of *Aspergillus* section *Nigri* capable of producing bioactive, antibiotic, and antitumor compounds. We sequenced the genome of a strain of *A. aculeatinus* that was isolated from the interior of a housing complex in Abidjan, Ivory Coast.

A *Aspergillus* section *Nigri* (the black aspergilli) consists of species that cause food spoilage, cause plant disease, and produce industrially relevant compounds like lipases, amylase, citric acid, and gluconic acid (1). *Aspergillus aculeatinus* is a member of the black aspergilli and closely related to *Aspergillus aculeatus* (2). *A. aculeatinus* has the potential for industrial application, as it produces the bioactive compound neoxaline, the antifungal compound aculeacin, and the antitumor compound paclitaxel (originally named Taxol [Bristol-Myers Squibb]) (2, 3). To date, only one *A. aculeatinus* genome has been sequenced (4).

To provide additional genomic resources for *A. aculeatinus*, we sequenced the genome of *A. aculeatinus* IC_8 after isolating it from an indoor air sample of a 23-story urban housing complex in Abidjan, Ivory Coast, that houses ~2,000 residents. Specifically, petri dishes with Sabouraud chloramphenicol agar were left open for 24 hours and then incubated at 25°C for 3 days. We used the hyphal tipping approach followed by incubation and single spore isolation to retrieve pure culture. DNA extraction was carried out as previously described (5). Briefly, spores were plated onto potato dextrose agar (PDA) and incubated at 37°C for 96 hours. Spores were collected and directly used for DNA extraction using the MasterPure yeast DNA purification kit following the manufacturer's instructions, with several minor modifications.

Next, 150-bp paired-end libraries were constructed and sequenced on an Illumina NovaSeq 6000 sequencer by Novogene. Raw reads were first deduplicated using Tally v15-065 with the "--with-quality" and "--pair-by-offset" options (6). Trim_Galore v0.4.2 was then used to remove residual adaptor sequences and to trim low-quality sequences using the parameters "--paired," "--stringency 1," "--quality 30," and "--length 50" (http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) (7). The deduplicated and trimmed data set contained 14,017,719 paired reads with a total of 4.07 billion bp. Next, the data were error corrected, and the genome was assembled *de novo* using SPAdes v3.13.1 with the "--careful" mode and a *k*-mer range of 55, 77, and 99 (8).

The assembly consisted of 441 scaffolds, a cumulative assembly size of 36.47 Mb (nearly identical to that of the *A. aculeatinus* CBS 121060 genome [4]), an *N*₅₀ value

Citation Zhao S, Koffi D, Latge J-P, Sylla K, Gibbons JG. 2021. Genome sequence of *Aspergillus aculeatinus* IC_8, isolated from an indoor air sample of an urban housing complex in Abidjan, Ivory Coast. *Microbiol Resour Announc* 10:e00096-21. <https://doi.org/10.1128/MRA.00096-21>.

Editor Antonis Rokas, Vanderbilt University

Copyright © 2021 Zhao et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to John G. Gibbons, jggibbons@umass.edu.

Received 27 January 2021

Accepted 16 February 2021

Published 11 March 2021




FIG 1 Phylogenetic relationship of 25 *Aspergillus* section *Nigri* genomes, including IC_8. The phylogeny was inferred by the approximately maximum-likelihood approach in FastTree (8) from a concatenated protein alignment of 4,680 sequences. All bootstrap branch support values were 100%. IC_8 is monophyletic with *A. aculeatinus* CBS 121060, and both taxa have short branch lengths, indicating that the species identity of IC_8 is *A. aculeatinus*. The species used are as follows (with their GenBank accession numbers for the whole-genome sequences): *A. aculeatinus* (PSTE000000000), *A. aculeatus* (GCA_001890905.1), *A. brunneoviolaceus* (PSTC000000000), *A. costaricaensis* (PSTH000000000), *A. ellipticus* (PSSY000000000), *A. eucalypticola* (MSFU000000000), *A. fijiensis* (PSTG000000000), *A. heteromorphus* (MSFL000000000), *A. homomorphus* (PSTJ000000000), *A. ibericus* (PSTI000000000), *A. indologenus* (PSTB000000000), *A. japonicus* (PSTF000000000), *A. lacticoffeatus* (MSFR000000000), *A. neoniger* (MSFP000000000), *A. niger* ATCC 13157 (*A. phoenicis*) (QQUR000000000), *A. niger* ATCC 13496 (QQZP000000000), *A. piperis* (PSTD000000000), *A. saccharolyticus* (MSFQ000000000), *A. sclerotiicarbonarius* (PSSZ000000000), *A. sclerotioniger* (MSFK000000000), *A. uvarum* (MSFT000000000), *A. vadensis* (MSFS000000000), *A. violaceofuscus* (PSTA000000000), and *A. welwitschiae* (QQZQ000000000).

of 649,318 bp, and a GC content of 50.48%. Genome completeness was evaluated with BUSCO v3.1.0 using the “ascomycota_odb9” gene set (9). A total of 98.9% of BUSCO genes were recovered from the IC_8 genome, indicating a high-quality genome assembly.

To verify the species of IC_8, we conducted a phylogenetic analysis of IC_8 and 24 genomes from 22 *Aspergillus* section *Nigri* species, including *A. aculeatinus* CBS 121060 (4). For all genomes, we used the Funannotate v1.7.0 (10) pipeline to predict gene models. Next, we used Orthofinder v2.3.3 to identify orthologous genes across the 25 genomes (11). A concatenated amino acid sequence alignment was generated from 4,680 translated genes. FastTree v2.1.10 was used to infer the phylogenetic relationship of isolates from the concatenated sequence alignment, using the MLACC=3 and nearest-neighbor interchange (NNI) options, with 100 bootstraps (12, 13). IC_8 is monophyletic with *A. aculeatinus* CBS 121060, and both taxa have short branch lengths (Fig. 1), providing clear evidence that the species identity of IC_8 is *A. aculeatinus*.

Data availability. The whole-genome shotgun project for *A. aculeatinus* IC_8 has been deposited in GenBank under the accession number [JADPID000000000](https://www.ncbi.nlm.nih.gov/GenBank/JADPID000000000). Raw Illumina data have been deposited to the NCBI Sequence Read Archive under the BioProject accession number [PRJNA675076](https://www.ncbi.nlm.nih.gov/BioProject/PRJNA675076).

ACKNOWLEDGMENTS

This work was supported by the National Institutes of Health and National Institutes of Allergy and Infectious Diseases (R21AI137485).

REFERENCES

1. Varga J, Frisvad JC, Kocsubé S, Brankovics B, Tóth B, Szigeti G, Samson RA. 2011. New and revisited species in *Aspergillus* section *Nigri*. *Stud Mycol* 69:1–17. <https://doi.org/10.3114/sim.2011.69.01>.
2. Noonim P, Mahakarnchanakul W, Varga J, Frisvad JC, Samson RA. 2008. Two novel species of *Aspergillus* section *Nigri* from Thai coffee beans. *Int J Syst Evol Microbiol* 58:1727–1734. <https://doi.org/10.1099/ijs.0.65694-0>.
3. Qiao W, Tang T, Ling F. 2020. Comparative transcriptome analysis of a taxol-producing endophytic fungus, *Aspergillus aculeatinus* Tax-6, and its mutant strain. *Sci Rep* 10:10558. <https://doi.org/10.1038/s41598-020-67614-1>.
4. Vesth TC, Nybo JL, Theobald S, Frisvad JC, Larsen TO, Nielsen KF, Hoof JB, Brandl J, Salamov A, Riley R, Gladden JM, Phatale P, Nielsen MT, Lyhne EK, Kogle ME, Strasser K, McDonnell E, Barry K, Clum A, Chen C, LaButti K, Haridas S, Nolan M, Sandor L, Kuo A, Lipzen A, Hainaut M, Drula E, Tsang A, Magnuson JK, Henrissat B, Wiebenga A, Simmons BA, Mäkelä MR, de Vries RP, Grigoriev IV, Mortensen UH, Baker SE, Andersen MR. 2018. Investigation of inter- and intraspecies variation through genome sequencing of *Aspergillus* section *Nigri*. *Nat Genet* 50:1688–1695. <https://doi.org/10.1038/s41588-018-0246-1>.
5. Zhao S, Latgé JP, Gibbons JG. 2019. Genome sequences of two strains of the food spoilage mold *Aspergillus fischeri*. *Microbiol Resour Announc* 8:e01328-19. <https://doi.org/10.1128/MRA.01328-19>.
6. Davis MP, van Dongen S, Abreu-Goodger C, Bartonicek N, Enright AJ. 2013. Kraken: a set of tools for quality control and analysis of high-throughput sequence data. *Methods* 63:41–49. <https://doi.org/10.1016/j.ymeth.2013.06.027>.
7. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J* 17:10–12. <https://doi.org/10.14806/ej.17.1.200>.
8. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
9. Seppey M, Manni M, Zdobnov EM. 2019. BUSCO: assessing genome assembly and annotation completeness. *Methods Mol Biol* 1962:227–245. https://doi.org/10.1007/978-1-4939-9173-0_14.
10. Palmer J, Stajich J. 2017. Funannotate: eukaryotic genome annotation pipeline. *Zenodo*. <https://doi.org/10.5281/zenodo.3548120>.
11. Price MN, Dehal PS, Arkin AP. 2010. FastTree 2—approximately maximum-likelihood trees for large alignments. *PLoS One* 5:e9490. <https://doi.org/10.1371/journal.pone.0009490>.
12. Emms DM, Kelly S. 2015. OrthoFinder: solving fundamental biases in whole genome comparisons dramatically improves orthogroup inference accuracy. *Genome Biol* 16:157. <https://doi.org/10.1186/s13059-015-0721-2>.
13. Emms DM, Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative genomics. *Genome Biol* 20:238. <https://doi.org/10.1186/s13059-019-1832-y>.