

2007

## Observation of CP violation in $B \rightarrow \eta/K^0$ decays

BABAR Collaboration

Follow this and additional works at: [https://scholarworks.umass.edu/cee\\_faculty\\_pubs](https://scholarworks.umass.edu/cee_faculty_pubs)

---

### Recommended Citation

BABAR Collaboration, "Observation of CP violation in  $B \rightarrow \eta/K^0$  decays" (2007). *Physical Review Letters*. 67.

<https://doi.org/10.1103/PhysRevLett.98.031801>

This Article is brought to you for free and open access by the Civil and Environmental Engineering at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Civil and Environmental Engineering Faculty Publication Series by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact [scholarworks@library.umass.edu](mailto:scholarworks@library.umass.edu).



## Observation of $CP$ Violation in $B \rightarrow \eta/K^0$ Decays

B. Aubert,<sup>1</sup> M. Bona,<sup>1</sup> D. Boutigny,<sup>1</sup> F. Couderc,<sup>1</sup> Y. Karyotakis,<sup>1</sup> J. P. Lees,<sup>1</sup> V. Poireau,<sup>1</sup> V. Tisserand,<sup>1</sup> A. Zghiche,<sup>1</sup> E. Grauges,<sup>2</sup> A. Palano,<sup>3</sup> J. C. Chen,<sup>4</sup> N. D. Qi,<sup>4</sup> G. Rong,<sup>4</sup> P. Wang,<sup>4</sup> Y. S. Zhu,<sup>4</sup> G. Eigen,<sup>5</sup> I. Ofte,<sup>5</sup> B. Stugu,<sup>5</sup> G. S. Abrams,<sup>6</sup> M. Battaglia,<sup>6</sup> D. N. Brown,<sup>6</sup> J. Button-Shafer,<sup>6</sup> R. N. Cahn,<sup>6</sup> E. Charles,<sup>6</sup> M. S. Gill,<sup>6</sup> Y. Groyzman,<sup>6</sup> R. G. Jacobsen,<sup>6</sup> J. A. Kadyk,<sup>6</sup> L. T. Kerth,<sup>6</sup> Yu. G. Kolomensky,<sup>6</sup> G. Kukartsev,<sup>6</sup> D. Lopes Pegna,<sup>6</sup> G. Lynch,<sup>6</sup> L. M. Mir,<sup>6</sup> T. J. Orimoto,<sup>6</sup> M. Pripstein,<sup>6</sup> N. A. Roe,<sup>6</sup> M. T. Ronan,<sup>6</sup> W. A. Wenzel,<sup>6</sup> P. del Amo Sanchez,<sup>7</sup> M. Barrett,<sup>7</sup> K. E. Ford,<sup>7</sup> T. J. Harrison,<sup>7</sup> A. J. Hart,<sup>7</sup> C. M. Hawkes,<sup>7</sup> A. T. Watson,<sup>7</sup> T. Held,<sup>8</sup> H. Koch,<sup>8</sup> B. Lewandowski,<sup>8</sup> M. Pelizaeus,<sup>8</sup> K. Peters,<sup>8</sup> T. Schroeder,<sup>8</sup> M. Steinke,<sup>8</sup> J. T. Boyd,<sup>9</sup> J. P. Burke,<sup>9</sup> W. N. Cottingham,<sup>9</sup> D. Walker,<sup>9</sup> D. J. Asgeirsson,<sup>10</sup> T. Cuhadar-Donszelmann,<sup>10</sup> B. G. Fulsom,<sup>10</sup> C. Hearty,<sup>10</sup> N. S. Knecht,<sup>10</sup> T. S. Mattison,<sup>10</sup> J. A. McKenna,<sup>10</sup> A. Khan,<sup>11</sup> P. Kyberd,<sup>11</sup> M. Saleem,<sup>11</sup> D. J. Sherwood,<sup>11</sup> L. Teodorescu,<sup>11</sup> V. E. Blinov,<sup>12</sup> A. D. Bukin,<sup>12</sup> V. P. Druzhinin,<sup>12</sup> V. B. Golubev,<sup>12</sup> A. P. Onuchin,<sup>12</sup> S. I. Serednyakov,<sup>12</sup> Yu. I. Skovpen,<sup>12</sup> E. P. Solodov,<sup>12</sup> K. Yu Todyshev,<sup>12</sup> D. S. Best,<sup>13</sup> M. Bondioli,<sup>13</sup> M. Bruinsma,<sup>13</sup> M. Chao,<sup>13</sup> S. Curry,<sup>13</sup> I. Eschrich,<sup>13</sup> D. Kirkby,<sup>13</sup> A. J. Lankford,<sup>13</sup> P. Lund,<sup>13</sup> M. Mandelkern,<sup>13</sup> W. Roethel,<sup>13</sup> D. P. Stoker,<sup>13</sup> S. Abachi,<sup>14</sup> C. Buchanan,<sup>14</sup> S. D. Foulkes,<sup>15</sup> J. W. Gary,<sup>15</sup> O. Long,<sup>15</sup> B. C. Shen,<sup>15</sup> K. Wang,<sup>15</sup> L. Zhang,<sup>15</sup> H. K. Hadavand,<sup>16</sup> E. J. Hill,<sup>16</sup> H. P. Paar,<sup>16</sup> S. Rahatlou,<sup>16</sup> V. Sharma,<sup>16</sup> J. W. Berryhill,<sup>17</sup> C. Campagnari,<sup>17</sup> A. Cunha,<sup>17</sup> B. Dahmes,<sup>17</sup> T. M. Hong,<sup>17</sup> D. Kovalskiy,<sup>17</sup> J. D. Richman,<sup>17</sup> T. W. Beck,<sup>18</sup> A. M. Eisner,<sup>18</sup> C. J. Flacco,<sup>18</sup> C. A. Heusch,<sup>18</sup> J. Kroseberg,<sup>18</sup> W. S. Lockman,<sup>18</sup> G. Nesom,<sup>18</sup> T. Schalk,<sup>18</sup> B. A. Schumm,<sup>18</sup> A. Seiden,<sup>18</sup> P. Spradlin,<sup>18</sup> D. C. Williams,<sup>18</sup> M. G. Wilson,<sup>18</sup> J. Albert,<sup>19</sup> E. Chen,<sup>19</sup> C. H. Cheng,<sup>19</sup> A. Dvoretzskii,<sup>19</sup> F. Fang,<sup>19</sup> D. G. Hitlin,<sup>19</sup> I. Narsky,<sup>19</sup> T. Piatenko,<sup>19</sup> F. C. Porter,<sup>19</sup> G. Mancinelli,<sup>20</sup> B. T. Meadows,<sup>20</sup> K. Mishra,<sup>20</sup> M. D. Sokoloff,<sup>20</sup> F. Blanc,<sup>21</sup> P. C. Bloom,<sup>21</sup> S. Chen,<sup>21</sup> W. T. Ford,<sup>21</sup> J. F. Hirschauer,<sup>21</sup> A. Kreisel,<sup>21</sup> M. Nagel,<sup>21</sup> U. Nauenberg,<sup>21</sup> A. Olivas,<sup>21</sup> W. O. Ruddick,<sup>21</sup> J. G. Smith,<sup>21</sup> K. A. Ulmer,<sup>21</sup> S. R. Wagner,<sup>21</sup> J. Zhang,<sup>21</sup> A. Chen,<sup>22</sup> E. A. Eckhart,<sup>22</sup> A. Soffer,<sup>22</sup> W. H. Toki,<sup>22</sup> R. J. Wilson,<sup>22</sup> F. Winklmeier,<sup>22</sup> Q. Zeng,<sup>22</sup> D. D. Altenburg,<sup>23</sup> E. Feltresi,<sup>23</sup> A. Hauke,<sup>23</sup> H. Jasper,<sup>23</sup> J. Merkel,<sup>23</sup> A. Petzold,<sup>23</sup> B. Spaan,<sup>23</sup> T. Brandt,<sup>24</sup> V. Klose,<sup>24</sup> H. M. Lacker,<sup>24</sup> W. F. Mader,<sup>24</sup> R. Nogowski,<sup>24</sup> J. Schubert,<sup>24</sup> K. R. Schubert,<sup>24</sup> R. Schwierz,<sup>24</sup> J. E. Sundermann,<sup>24</sup> A. Volk,<sup>24</sup> D. Bernard,<sup>25</sup> G. R. Bonneaud,<sup>25</sup> E. Latour,<sup>25</sup> Ch. Thiebaux,<sup>25</sup> M. Verderi,<sup>25</sup> P. J. Clark,<sup>26</sup> W. Gradl,<sup>26</sup> F. Muheim,<sup>26</sup> S. Playfer,<sup>26</sup> A. I. Robertson,<sup>26</sup> Y. Xie,<sup>26</sup> M. Andreotti,<sup>27</sup> D. Bettoni,<sup>27</sup> C. Bozzi,<sup>27</sup> R. Calabrese,<sup>27</sup> G. Cibinetto,<sup>27</sup> E. Luppi,<sup>27</sup> M. Negrini,<sup>27</sup> A. Petrella,<sup>27</sup> L. Piemontese,<sup>27</sup> E. Prencipe,<sup>27</sup> F. Anulli,<sup>28</sup> R. Baldini-Ferroli,<sup>28</sup> A. Calcaterra,<sup>28</sup> R. de Sangro,<sup>28</sup> G. Finocchiaro,<sup>28</sup> S. Pacetti,<sup>28</sup> P. Patteri,<sup>28</sup> I. M. Peruzzi,<sup>28,\*</sup> M. Piccolo,<sup>28</sup> M. Rama,<sup>28</sup> A. Zallo,<sup>28</sup> A. Buzzo,<sup>29</sup> R. Contri,<sup>29</sup> M. Lo Vetere,<sup>29</sup> M. M. Macri,<sup>29</sup> M. R. Monge,<sup>29</sup> S. Passaggio,<sup>29</sup> C. Patrignani,<sup>29</sup> E. Robutti,<sup>29</sup> A. Santroni,<sup>29</sup> S. Tosi,<sup>29</sup> G. Brandenburg,<sup>30</sup> K. S. Chaisanguanthum,<sup>30</sup> C. L. Lee,<sup>30</sup> M. Morii,<sup>30</sup> J. Wu,<sup>30</sup> R. S. Dubitzky,<sup>31</sup> J. Marks,<sup>31</sup> S. Schenk,<sup>31</sup> U. Uwer,<sup>31</sup> D. J. Bard,<sup>32</sup> W. Bhimji,<sup>32</sup> D. A. Bowerman,<sup>32</sup> P. D. Dauncey,<sup>32</sup> U. Egede,<sup>32</sup> R. L. Flack,<sup>32</sup> J. A. Nash,<sup>32</sup> M. B. Nikolich,<sup>32</sup> W. Panduro Vazquez,<sup>32</sup> P. K. Behera,<sup>33</sup> X. Chai,<sup>33</sup> M. J. Charles,<sup>33</sup> U. Mallik,<sup>33</sup> N. T. Meyer,<sup>33</sup> V. Ziegler,<sup>33</sup> J. Cochran,<sup>34</sup> H. B. Crawley,<sup>34</sup> L. Dong,<sup>34</sup> V. Eyges,<sup>34</sup> W. T. Meyer,<sup>34</sup> S. Prell,<sup>34</sup> E. I. Rosenberg,<sup>34</sup> A. E. Rubin,<sup>34</sup> A. V. Gritsan,<sup>35</sup> A. G. Denig,<sup>36</sup> M. Fritsch,<sup>36</sup> G. Schott,<sup>36</sup> N. Arnaud,<sup>37</sup> M. Davier,<sup>37</sup> G. Grosdidier,<sup>37</sup> A. Höcker,<sup>37</sup> V. Lepeltier,<sup>37</sup> F. Le Diberder,<sup>37</sup> A. M. Lutz,<sup>37</sup> A. Oyanguren,<sup>37</sup> S. Pruvot,<sup>37</sup> S. Rodier,<sup>37</sup> P. Roudeau,<sup>37</sup> M. H. Schune,<sup>37</sup> J. Serrano,<sup>37</sup> A. Stocchi,<sup>37</sup> W. F. Wang,<sup>37</sup> G. Wormser,<sup>37</sup> D. J. Lange,<sup>38</sup> D. M. Wright,<sup>38</sup> C. A. Chavez,<sup>39</sup> I. J. Forster,<sup>39</sup> J. R. Fry,<sup>39</sup> E. Gabathuler,<sup>39</sup> R. Gamet,<sup>39</sup> K. A. George,<sup>39</sup> D. E. Hutchcroft,<sup>39</sup> D. J. Payne,<sup>39</sup> K. C. Schofield,<sup>39</sup> C. Touramanis,<sup>39</sup> A. J. Bevan,<sup>40</sup> C. K. Clarke,<sup>40</sup> F. Di Lodovico,<sup>40</sup> W. Menges,<sup>40</sup> R. Sacco,<sup>40</sup> G. Cowan,<sup>41</sup> H. U. Flaecher,<sup>41</sup> D. A. Hopkins,<sup>41</sup> P. S. Jackson,<sup>41</sup> T. R. McMahon,<sup>41</sup> F. Salvatore,<sup>41</sup> A. C. Wren,<sup>41</sup> D. N. Brown,<sup>42</sup> C. L. Davis,<sup>42</sup> J. Allison,<sup>43</sup> N. R. Barlow,<sup>43</sup> R. J. Barlow,<sup>43</sup> Y. M. Chia,<sup>43</sup> C. L. Edgar,<sup>43</sup> G. D. Lafferty,<sup>43</sup> M. T. Naisbit,<sup>43</sup> J. C. Williams,<sup>43</sup> J. I. Yi,<sup>43</sup> C. Chen,<sup>44</sup> W. D. Hulsbergen,<sup>44</sup> A. Jawahery,<sup>44</sup> C. K. Lae,<sup>44</sup> D. A. Roberts,<sup>44</sup> G. Simi,<sup>44</sup> G. Blaylock,<sup>45</sup> C. Dallapiccola,<sup>45</sup> S. S. Hertzbach,<sup>45</sup> X. Li,<sup>45</sup> T. B. Moore,<sup>45</sup> S. Saremi,<sup>45</sup> H. Staengle,<sup>45</sup> R. Cowan,<sup>46</sup> G. Sciolla,<sup>46</sup> S. J. Sekula,<sup>46</sup> M. Spitznagel,<sup>46</sup> F. Taylor,<sup>46</sup> R. K. Yamamoto,<sup>46</sup> H. Kim,<sup>47</sup> S. E. Mclachlin,<sup>47</sup> P. M. Patel,<sup>47</sup> S. H. Robertson,<sup>47</sup> A. Lazzaro,<sup>48</sup> V. Lombardo,<sup>48</sup> F. Palombo,<sup>48</sup> J. M. Bauer,<sup>49</sup> L. Cremaldi,<sup>49</sup> V. Eschenburg,<sup>49</sup> R. Godang,<sup>49</sup> R. Kroeger,<sup>49</sup> D. A. Sanders,<sup>49</sup> D. J. Summers,<sup>49</sup> H. W. Zhao,<sup>49</sup> S. Brunet,<sup>50</sup> D. Côté,<sup>50</sup> M. Simard,<sup>50</sup> P. Taras,<sup>50</sup> F. B. Viard,<sup>50</sup> H. Nicholson,<sup>51</sup> N. Cavallo,<sup>52,†</sup> G. De Nardo,<sup>52</sup> F. Fabozzi,<sup>52,†</sup> C. Gatto,<sup>52</sup> L. Lista,<sup>52</sup> D. Monorchio,<sup>52</sup> P. Paolucci,<sup>52</sup> D. Piccolo,<sup>52</sup> C. Sciacca,<sup>52</sup> M. A. Baak,<sup>53</sup> G. Raven,<sup>53</sup> H. L. Snoek,<sup>53</sup> C. P. Jessop,<sup>54</sup> J. M. LoSecco,<sup>54</sup> G. Benelli,<sup>55</sup> L. A. Corwin,<sup>55</sup> K. K. Gan,<sup>55</sup> K. Honscheid,<sup>55</sup> D. Hufnagel,<sup>55</sup> P. D. Jackson,<sup>55</sup> H. Kagan,<sup>55</sup> R. Kass,<sup>55</sup> A. M. Rahimi,<sup>55</sup> J. J. Regensburger,<sup>55</sup> R. Ter-Antonyan,<sup>55</sup> Q. K. Wong,<sup>55</sup> N. L. Blount,<sup>56</sup> J. Brau,<sup>56</sup> R. Frey,<sup>56</sup> O. Igonkina,<sup>56</sup> J. A. Kolb,<sup>56</sup> M. Lu,<sup>56</sup> C. T. Potter,<sup>56</sup> R. Rahmat,<sup>56</sup> N. B. Sinev,<sup>56</sup> D. Strom,<sup>56</sup> J. Strube,<sup>56</sup> E. Torrence,<sup>56</sup>

A. Gaz,<sup>57</sup> M. Margoni,<sup>57</sup> M. Morandin,<sup>57</sup> A. Pompili,<sup>57</sup> M. Posocco,<sup>57</sup> M. Rotondo,<sup>57</sup> F. Simonetto,<sup>57</sup> R. Stroili,<sup>57</sup> C. Voci,<sup>57</sup> M. Benayoun,<sup>58</sup> H. Briand,<sup>58</sup> J. Chauveau,<sup>58</sup> P. David,<sup>58</sup> L. Del Buono,<sup>58</sup> Ch. de la Vaissière,<sup>58</sup> O. Hamon,<sup>58</sup> B. L. Hartfiel,<sup>58</sup> Ph. Leruste,<sup>58</sup> J. Malclès,<sup>58</sup> J. Ocariz,<sup>58</sup> L. Roos,<sup>58</sup> G. Therin,<sup>58</sup> L. Gladney,<sup>59</sup> M. Biasini,<sup>60</sup> R. Covarelli,<sup>60</sup> C. Angelini,<sup>61</sup> G. Batignani,<sup>61</sup> S. Bettarini,<sup>61</sup> F. Bucci,<sup>61</sup> G. Calderini,<sup>61</sup> M. Carpinelli,<sup>61</sup> R. Cenci,<sup>61</sup> F. Forti,<sup>61</sup> M. A. Giorgi,<sup>61</sup> A. Lusiani,<sup>61</sup> G. Marchiori,<sup>61</sup> M. A. Mazur,<sup>61</sup> M. Morganti,<sup>61</sup> N. Neri,<sup>61</sup> E. Paoloni,<sup>61</sup> G. Rizzo,<sup>61</sup> J. J. Walsh,<sup>61</sup> M. Haire,<sup>62</sup> D. Judd,<sup>62</sup> D. E. Wagoner,<sup>62</sup> J. Biesiada,<sup>63</sup> N. Danielson,<sup>63</sup> P. Elmer,<sup>63</sup> Y. P. Lau,<sup>63</sup> C. Lu,<sup>63</sup> J. Olsen,<sup>63</sup> A. J. S. Smith,<sup>63</sup> A. V. Telnov,<sup>63</sup> F. Bellini,<sup>64</sup> G. Cavoto,<sup>64</sup> A. D’Orazio,<sup>64</sup> D. del Re,<sup>64</sup> E. Di Marco,<sup>64</sup> R. Faccini,<sup>64</sup> F. Ferrarotto,<sup>64</sup> F. Ferroni,<sup>64</sup> M. Gaspero,<sup>64</sup> L. Li Gioi,<sup>64</sup> M. A. Mazzoni,<sup>64</sup> S. Morganti,<sup>64</sup> G. Piredda,<sup>64</sup> F. Polci,<sup>64</sup> F. Safai Tehrani,<sup>64</sup> C. Voena,<sup>64</sup> M. Ebert,<sup>65</sup> H. Schröder,<sup>65</sup> R. Waldi,<sup>65</sup> T. Adye,<sup>66</sup> B. Franek,<sup>66</sup> E. O. Olaiya,<sup>66</sup> S. Ricciardi,<sup>66</sup> F. F. Wilson,<sup>66</sup> R. Aleksan,<sup>67</sup> S. Emery,<sup>67</sup> A. Gaidot,<sup>67</sup> S. F. Ganzhur,<sup>67</sup> G. Hamel de Monchenault,<sup>67</sup> W. Kozanecki,<sup>67</sup> M. Legendre,<sup>67</sup> G. Vasseur,<sup>67</sup> Ch. Yèche,<sup>67</sup> M. Zito,<sup>67</sup> X. R. Chen,<sup>68</sup> H. Liu,<sup>68</sup> W. Park,<sup>68</sup> M. V. Purohit,<sup>68</sup> J. R. Wilson,<sup>68</sup> M. T. Allen,<sup>69</sup> D. Aston,<sup>69</sup> R. Bartoldus,<sup>69</sup> P. Bechtel,<sup>69</sup> N. Berger,<sup>69</sup> R. Claus,<sup>69</sup> J. P. Coleman,<sup>69</sup> M. R. Convery,<sup>69</sup> J. C. Dingfelder,<sup>69</sup> J. Dorfan,<sup>69</sup> G. P. Dubois-Felsmann,<sup>69</sup> D. Dujmic,<sup>69</sup> W. Dunwoodie,<sup>69</sup> R. C. Field,<sup>69</sup> T. Glanzman,<sup>69</sup> S. J. Gowdy,<sup>69</sup> M. T. Graham,<sup>69</sup> P. Grenier,<sup>69</sup> V. Halyo,<sup>69</sup> C. Hast,<sup>69</sup> T. Hryn’ova,<sup>69</sup> W. R. Innes,<sup>69</sup> M. H. Kelsey,<sup>69</sup> P. Kim,<sup>69</sup> D. W. G. S. Leith,<sup>69</sup> S. Li,<sup>69</sup> S. Luitz,<sup>69</sup> V. Luth,<sup>69</sup> H. L. Lynch,<sup>69</sup> D. B. MacFarlane,<sup>69</sup> H. Marsiske,<sup>69</sup> R. Messner,<sup>69</sup> D. R. Muller,<sup>69</sup> C. P. O’Grady,<sup>69</sup> V. E. Ozcan,<sup>69</sup> A. Perazzo,<sup>69</sup> M. Perl,<sup>69</sup> T. Pulliam,<sup>69</sup> B. N. Ratcliff,<sup>69</sup> A. Roodman,<sup>69</sup> A. A. Salnikov,<sup>69</sup> R. H. Schindler,<sup>69</sup> J. Schwiening,<sup>69</sup> A. Snyder,<sup>69</sup> J. Stelzer,<sup>69</sup> D. Su,<sup>69</sup> M. K. Sullivan,<sup>69</sup> K. Suzuki,<sup>69</sup> S. K. Swain,<sup>69</sup> J. M. Thompson,<sup>69</sup> J. Va’vra,<sup>69</sup> N. van Bakel,<sup>69</sup> A. P. Wagner,<sup>69</sup> M. Weaver,<sup>69</sup> A. J. R. Weinstein,<sup>69</sup> W. J. Wisniewski,<sup>69</sup> M. Wittgen,<sup>69</sup> D. H. Wright,<sup>69</sup> H. W. Wulsin,<sup>69</sup> A. K. Yarritu,<sup>69</sup> K. Yi,<sup>69</sup> C. C. Young,<sup>69</sup> P. R. Burchat,<sup>70</sup> A. J. Edwards,<sup>70</sup> S. A. Majewski,<sup>70</sup> B. A. Petersen,<sup>70</sup> L. Wilden,<sup>70</sup> S. Ahmed,<sup>71</sup> M. S. Alam,<sup>71</sup> R. Bula,<sup>71</sup> J. A. Ernst,<sup>71</sup> V. Jain,<sup>71</sup> B. Pan,<sup>71</sup> M. A. Saeed,<sup>71</sup> F. R. Wappler,<sup>71</sup> S. B. Zain,<sup>71</sup> W. Bugg,<sup>72</sup> M. Krishnamurthy,<sup>72</sup> S. M. Spanier,<sup>72</sup> R. Eckmann,<sup>73</sup> J. L. Ritchie,<sup>73</sup> A. Satpathy,<sup>73</sup> C. J. Schilling,<sup>73</sup> R. F. Schwitters,<sup>73</sup> J. M. Izen,<sup>74</sup> X. C. Lou,<sup>74</sup> S. Ye,<sup>74</sup> F. Bianchi,<sup>75</sup> F. Gallo,<sup>75</sup> D. Gamba,<sup>75</sup> M. Bomben,<sup>76</sup> L. Bosisio,<sup>76</sup> C. Cartaro,<sup>76</sup> F. Cossutti,<sup>76</sup> G. Della Ricca,<sup>76</sup> S. Dittongo,<sup>76</sup> L. Lanceri,<sup>76</sup> L. Vitale,<sup>76</sup> V. Azzolini,<sup>77</sup> N. Lopez-March,<sup>77</sup> F. Martinez-Vidal,<sup>77</sup> Sw. Banerjee,<sup>78</sup> B. Bhuyan,<sup>78</sup> C. M. Brown,<sup>78</sup> D. Fortin,<sup>78</sup> K. Hamano,<sup>78</sup> R. Kowalewski,<sup>78</sup> I. M. Nugent,<sup>78</sup> J. M. Roney,<sup>78</sup> R. J. Sobie,<sup>78</sup> J. J. Back,<sup>79</sup> P. F. Harrison,<sup>79</sup> T. E. Latham,<sup>79</sup> G. B. Mohanty,<sup>79</sup> M. Pappagallo,<sup>79,‡</sup> H. R. Band,<sup>80</sup> X. Chen,<sup>80</sup> B. Cheng,<sup>80</sup> S. Dasu,<sup>80</sup> M. Datta,<sup>80</sup> K. T. Flood,<sup>80</sup> J. J. Hollar,<sup>80</sup> P. E. Kutter,<sup>80</sup> B. Mellado,<sup>80</sup> A. Mihalyi,<sup>80</sup> Y. Pan,<sup>80</sup> M. Pierini,<sup>80</sup> R. Prepost,<sup>80</sup> S. L. Wu,<sup>80</sup> Z. Yu,<sup>80</sup> and H. Neal<sup>81</sup>

(BABAR Collaboration)

<sup>1</sup>Laboratoire de Physique des Particules, IN2P3/CNRS et Université de Savoie, F-74941 Annecy-Le-Vieux, France

<sup>2</sup>Universitat de Barcelona, Facultat de Física, Departament ECM, E-08028 Barcelona, Spain

<sup>3</sup>Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

<sup>4</sup>Institute of High Energy Physics, Beijing 100039, China

<sup>5</sup>University of Bergen, Institute of Physics, N-5007 Bergen, Norway

<sup>6</sup>Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

<sup>7</sup>University of Birmingham, Birmingham, B15 2TT, United Kingdom

<sup>8</sup>Ruhr Universität Bochum, Institut für Experimentalphysik I, D-44780 Bochum, Germany

<sup>9</sup>University of Bristol, Bristol BS8 1TL, United Kingdom

<sup>10</sup>University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

<sup>11</sup>Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

<sup>12</sup>Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

<sup>13</sup>University of California at Irvine, Irvine, California 92697, USA

<sup>14</sup>University of California at Los Angeles, Los Angeles, California 90024, USA

<sup>15</sup>University of California at Riverside, Riverside, California 92521, USA

<sup>16</sup>University of California at San Diego, La Jolla, California 92093, USA

<sup>17</sup>University of California at Santa Barbara, Santa Barbara, California 93106, USA

<sup>18</sup>University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

<sup>19</sup>California Institute of Technology, Pasadena, California 91125, USA

<sup>20</sup>University of Cincinnati, Cincinnati, Ohio 45221, USA

<sup>21</sup>University of Colorado, Boulder, Colorado 80309, USA

<sup>22</sup>Colorado State University, Fort Collins, Colorado 80523, USA

<sup>23</sup>Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany

<sup>24</sup>Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany

- <sup>25</sup>Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, F-91128 Palaiseau, France
- <sup>26</sup>University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
- <sup>27</sup>Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
- <sup>28</sup>Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
- <sup>29</sup>Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
- <sup>30</sup>Harvard University, Cambridge, Massachusetts 02138, USA
- <sup>31</sup>Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
- <sup>32</sup>Imperial College London, London, SW7 2AZ, United Kingdom
- <sup>33</sup>University of Iowa, Iowa City, Iowa 52242, USA
- <sup>34</sup>Iowa State University, Ames, Iowa 50011-3160, USA
- <sup>35</sup>Johns Hopkins University, Baltimore, Maryland 21218, USA
- <sup>36</sup>Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany
- <sup>37</sup>Laboratoire de l'Accélérateur Linéaire, IN2P3/CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B.P. 34, F-91898 ORSAY Cedex, France
- <sup>38</sup>Lawrence Livermore National Laboratory, Livermore, California 94550, USA
- <sup>39</sup>University of Liverpool, Liverpool L69 7ZE, United Kingdom
- <sup>40</sup>Queen Mary, University of London, E1 4NS, United Kingdom
- <sup>41</sup>University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
- <sup>42</sup>University of Louisville, Louisville, Kentucky 40292, USA
- <sup>43</sup>University of Manchester, Manchester M13 9PL, United Kingdom
- <sup>44</sup>University of Maryland, College Park, Maryland 20742, USA
- <sup>45</sup>University of Massachusetts, Amherst, Massachusetts 01003, USA
- <sup>46</sup>Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
- <sup>47</sup>McGill University, Montréal, Québec, Canada H3A 2T8
- <sup>48</sup>Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
- <sup>49</sup>University of Mississippi, University, Mississippi 38677, USA
- <sup>50</sup>Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7
- <sup>51</sup>Mount Holyoke College, South Hadley, Massachusetts 01075, USA
- <sup>52</sup>Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
- <sup>53</sup>NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
- <sup>54</sup>University of Notre Dame, Notre Dame, Indiana 46556, USA
- <sup>55</sup>The Ohio State University, Columbus, Ohio 43210, USA
- <sup>56</sup>University of Oregon, Eugene, Oregon 97403, USA
- <sup>57</sup>Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
- <sup>58</sup>Laboratoire de Physique Nucléaire et de Hautes Energies, IN2P3/CNRS, Université Pierre et Marie Curie-Paris6, Université Denis Diderot-Paris7, F-75252 Paris, France
- <sup>59</sup>University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
- <sup>60</sup>Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
- <sup>61</sup>Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
- <sup>62</sup>Prairie View A&M University, Prairie View, Texas 77446, USA
- <sup>63</sup>Princeton University, Princeton, New Jersey 08544, USA
- <sup>64</sup>Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
- <sup>65</sup>Universität Rostock, D-18051 Rostock, Germany
- <sup>66</sup>Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
- <sup>67</sup>DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
- <sup>68</sup>University of South Carolina, Columbia, South Carolina 29208, USA
- <sup>69</sup>Stanford Linear Accelerator Center, Stanford, California 94309, USA
- <sup>70</sup>Stanford University, Stanford, California 94305-4060, USA
- <sup>71</sup>State University of New York, Albany, New York 12222, USA
- <sup>72</sup>University of Tennessee, Knoxville, Tennessee 37996, USA
- <sup>73</sup>University of Texas at Austin, Austin, Texas 78712, USA
- <sup>74</sup>University of Texas at Dallas, Richardson, Texas 75083, USA
- <sup>75</sup>Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
- <sup>76</sup>Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
- <sup>77</sup>IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain
- <sup>78</sup>University of Victoria, Victoria, British Columbia, Canada V8W 3P6
- <sup>79</sup>Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom
- <sup>80</sup>University of Wisconsin, Madison, Wisconsin 53706, USA
- <sup>81</sup>Yale University, New Haven, Connecticut 06511, USA

(Received 22 September 2006; published 16 January 2007)

We present measurements of the time-dependent  $CP$ -violation parameters  $S$  and  $C$  in  $B^0 \rightarrow \eta' K^0$  decays. The data sample corresponds to  $384 \times 10^6 B\bar{B}$  pairs produced by  $e^+e^-$  annihilation at the  $Y(4S)$ . The results are  $S = 0.58 \pm 0.10 \pm 0.03$  and  $C = -0.16 \pm 0.07 \pm 0.03$ . We observe mixing-induced  $CP$  violation with a significance of 5.5 standard deviations in this  $b \rightarrow s$  penguin dominated mode.

DOI: 10.1103/PhysRevLett.98.031801

PACS numbers: 13.25.Hw, 11.30.Er, 12.15.Hh

Measurements of time-dependent  $CP$  asymmetries in  $B^0$  meson decays through Cabibbo-Kobayashi-Maskawa (CKM) favored  $b \rightarrow c\bar{c}s$  amplitudes [1] have provided crucial tests of the mechanism of  $CP$  violation in the standard model (SM) [2]. Decays of  $B^0$  mesons to charmless hadronic final states such as  $\eta' K^0$  proceed mostly via a single loop (penguin) amplitude. In the SM the penguin amplitude has approximately the same weak phase as the  $b \rightarrow c\bar{c}s$  transition, but it is sensitive to the possible presence of new heavy particles in the loop [3]. The measurement of  $CP$  asymmetries in  $B^0 \rightarrow \eta' K^0$  thus provides an important test for such effects.

Within the SM, CKM-suppressed amplitudes and multiple particles in the loop introduce additional weak phases whose contribution may not be negligible [4–7]. The time-dependent  $CP$ -violation parameter  $S$  [defined in Eq. (1) below] measured in the decay  $B^0 \rightarrow \eta' K^0$  is compared with the value of  $\sin 2\beta$  from measurements of time-dependent  $CP$  violation in  $B$  decays to states containing charmonium and a neutral kaon. The deviation  $\Delta S = S - \sin 2\beta$  has been estimated in several theoretical approaches: QCD factorization (QCDF) [6,8], QCDF with modeled rescattering [9], soft collinear effective theory [10], and SU(3) symmetry [4,5,11]. These models estimate  $|\Delta S|$  to be of the order 0.01, and with uncertainties give bounds  $|\Delta S| \leq 0.05$ .

The time-dependent  $CP$  asymmetry in the decay  $B^0 \rightarrow \eta' K_S^0$  has been measured previously by the BABAR [12] and Belle [13] Collaborations. In this Letter we update our previous measurements using an integrated luminosity of  $349 \text{ fb}^{-1}$ , corresponding to  $(384 \pm 4) \times 10^6 B\bar{B}$  pairs, recorded at the  $Y(4S)$  resonance (center-of-mass energy  $\sqrt{s} = 10.58 \text{ GeV}$ ). Belle has since updated their results [14]. Our data were collected with the BABAR detector [15] at the PEP-II asymmetric-energy  $e^+e^-$  collider. In addition to the  $B^0 \rightarrow \eta' K_S^0$  decays used previously, we now also include the decay  $B^0 \rightarrow \eta' K_L^0$ .

Charged particles from  $e^+e^-$  interactions are detected, and their momenta measured, by a combination of five layers of double-sided silicon microstrip detectors and a 40-layer drift chamber, both operating in the 1.5 T magnetic field of a superconducting solenoid. Photons and electrons are identified with a CsI(Tl) electromagnetic calorimeter (EMC). Charged particle identification is provided by the average energy loss in the tracking devices and by an internally reflecting ring imaging Cherenkov detector covering the central region. The instrumented flux return (IFR) of the magnet allows the identification of muons and  $K_L^0$  mesons.

We reconstruct a  $B^0$  decaying into the  $CP$  eigenstate  $\eta' K_S^0$  or  $\eta' K_L^0$  ( $B_{CP}$ ). From the remaining particles in the event we also reconstruct the decay vertex of the other  $B$  meson ( $B_{\text{tag}}$ ) and identify its flavor. The difference  $\Delta t \equiv t_{CP} - t_{\text{tag}}$  of the proper decay times  $t_{CP}$  and  $t_{\text{tag}}$  of the  $CP$  and tag  $B$  mesons, respectively, is obtained from the measured distance between the  $B_{CP}$  and  $B_{\text{tag}}$  decay vertices and from the boost ( $\beta\gamma = 0.56$ ) of the  $e^+e^-$  system. The  $\Delta t$  distribution is given by

$$F(\Delta t) = \frac{e^{-|\Delta t|/\tau}}{4\tau} \{1 \mp \Delta w \pm (1 - 2w) \times [-\eta S \sin(\Delta m_d \Delta t) - C \cos(\Delta m_d \Delta t)]\} \quad (1)$$

where  $\eta$  is the  $CP$  eigenvalue of the final state ( $-1$  for  $\eta' K_S^0$ ,  $+1$  for  $\eta' K_L^0$ ). The upper (lower) sign denotes a decay accompanied by a  $B^0(\bar{B}^0)$  tag,  $\tau$  is the mean  $B^0$  lifetime,  $\Delta m_d$  is the mixing frequency, and the mistag parameters  $w$  and  $\Delta w$  are the average and difference, respectively, of the probabilities that a true  $B^0$  is incorrectly tagged as a  $\bar{B}^0$  or vice versa. The tagging algorithm has six mutually exclusive tagging categories and a measured analyzing power of  $(30.4 \pm 0.3)\%$  [16]. A nonzero value of the parameter  $C$  would indicate direct  $CP$  violation.

We establish the event selection criteria with the aid of a detailed Monte Carlo (MC) simulation of the  $B$  production and decay sequences, and of the detector response [17]. These criteria are designed to retain signal events with high efficiency while removing most of the background.

The  $B$ -daughter candidates are reconstructed through their decays  $\pi^0 \rightarrow \gamma\gamma$ ,  $\eta \rightarrow \gamma\gamma$  ( $\eta_{\gamma\gamma}$ ),  $\eta \rightarrow \pi^+\pi^-\pi^0$  ( $\eta_{3\pi}$ ),  $\eta' \rightarrow \eta_{\gamma\gamma}\pi^+\pi^-$  ( $\eta'_{\eta(\gamma\gamma)\pi\pi}$ ),  $\eta' \rightarrow \eta_{3\pi}\pi^+\pi^-$  ( $\eta'_{\eta(3\pi)\pi\pi}$ ),  $\eta' \rightarrow \rho^0\gamma$  ( $\eta'_{\rho\gamma}$ ), where  $\rho^0 \rightarrow \pi^+\pi^-$ ,  $K_S^0 \rightarrow \pi^+\pi^-$  ( $K_{\pi^+\pi^-}^0$ ) or  $\pi^0\pi^0$  ( $K_{\pi^0\pi^0}^0$ ). Only the  $\eta'_{\eta(\gamma\gamma)\pi\pi}$  mode is used for the  $\eta' K_L^0$  sample. The requirements on the invariant masses of these particle combinations are the same as in our previous analysis [12]. The list of all decay modes used in the current analysis can be seen in Table I. Signal  $K_L^0$  candidates are reconstructed from clusters of energy deposited in the EMC or from hits in the IFR not associated with any charged track in the event [18]. From the cluster centroid and the  $B_0$  decay vertex we determine the direction (but not the magnitude) of the  $K_L^0$  momentum  $\mathbf{p}_{K_L^0}$ .

For  $\eta' K_S^0$  decays we reconstruct the  $B$ -meson candidate by combining the four-momenta of the  $K_S^0$  and  $\eta'$  with a vertex constraint. We also constrain the  $\eta$ ,  $\eta'$ , and  $\pi^0$  masses to world-average values [19]. From the kinematics

TABLE I. Results of the fits. Subscripts for  $\eta'$  decay modes denote  $\eta'_{\eta(\gamma\gamma)\pi\pi}$  (1),  $\eta'_{\rho\gamma}$  (2), and  $\eta'_{\eta(3\pi)\pi\pi}$  (3).

Mode	# events	Signal yield	$S$	$C$
$\eta'_1 K_{\pi^+\pi^-}^0$	664	$224 \pm 16$	$0.61 \pm 0.23$	$-0.26 \pm 0.14$
$\eta'_2 K_{\pi^+\pi^-}^0$	11 943	$566 \pm 30$	$0.56 \pm 0.14$	$-0.24 \pm 0.10$
$\eta'_3 K_{\pi^+\pi^-}^0$	177	$73 \pm 9$	$0.89 \pm 0.35$	$0.14 \pm 0.25$
$\eta'_1 K_{\pi^0\pi^0}^0$	490	$52 \pm 9$	$0.84 \pm 0.42$	$-0.26 \pm 0.36$
$\eta'_2 K_{\pi^0\pi^0}^0$	13 915	$133 \pm 24$	$0.56 \pm 0.41$	$0.15 \pm 0.27$
$\eta'_1 K_S^0$			$0.62 \pm 0.11$	$-0.18 \pm 0.07$
$\eta'_1 K_L^0$	4199	$204 \pm 24$	$0.32 \pm 0.28$	$0.08 \pm 0.23$
$\eta' K^0$			$0.58 \pm 0.10$	$-0.16 \pm 0.07$

of  $Y(4S)$  decays we determine the energy-substituted mass  $m_{\text{ES}} \equiv \sqrt{(\frac{1}{2}s + \mathbf{p}_0 \cdot \mathbf{p}_B)^2/E_0^2 - \mathbf{p}_B^2}$  and the energy difference  $\Delta E \equiv E_B^* - \frac{1}{2}\sqrt{s}$ , where  $(E_0, \mathbf{p}_0)$  and  $(E_B, \mathbf{p}_B)$  are the laboratory four-momenta of the  $Y(4S)$  and the  $B$  candidate, respectively, and the asterisk denotes the  $Y(4S)$  rest frame. The resolution is 3 MeV in  $m_{\text{ES}}$  and 20–50 MeV in  $\Delta E$ , depending on the decay mode.

For  $\eta' K_L^0$  candidates we obtain  $\Delta E$  and  $\mathbf{p}_{K_L^0}$  from a fit with the  $B^0$  and  $K_L^0$  masses constrained to world-average values [19]. To make a match with the measured  $K_L^0$  direction we construct the missing momentum  $\mathbf{p}_{\text{miss}}$  from  $\mathbf{p}_0$  and all charged tracks and neutral clusters other than the  $K_L^0$  candidate. We then project  $\mathbf{p}_{\text{miss}}$  onto  $\mathbf{p}_{K_L^0}$ , and require the component perpendicular to the beam line,  $p_{\text{miss}\perp}^{\text{proj}}$ , to satisfy  $p_{\text{miss}\perp}^{\text{proj}} - p_{K_L^0\perp} > -0.5$  GeV. This value was chosen to minimize the yield uncertainty in the presence of background.

For  $\eta' K_S^0$  we require  $5.25 < m_{\text{ES}} < 5.29$  GeV and  $|\Delta E| < 0.2$  GeV, for  $\eta' K_L^0$  we require  $-0.01 < \Delta E < 0.04$  GeV, and for all decays  $|\Delta t| < 20$  ps, and, for the error on  $\Delta t$ ,  $\sigma_{\Delta t} < 2.5$  ps.

Background events arise primarily from random combinations of particles in continuum  $e^+e^- \rightarrow q\bar{q}$  events ( $q = u, d, s, c$ ). We reduce these with requirements on the angle  $\theta_T$  between the thrust axis of the  $B$  candidate in the  $Y(4S)$  frame and that of the rest of the charged tracks and neutral calorimeter clusters in the event. In the fit we discriminate further against  $q\bar{q}$  background with a Fisher discriminant  $\mathcal{F}$  that combines several variables that characterize the production dynamics and energy flow in the event [20]. For the  $\eta'_\rho\gamma$  decays we require  $|\cos\theta_{\text{dec}}^\rho| < 0.9$  to reduce the combinatorial background. Here  $\theta_{\text{dec}}^\rho$  is the angle between the momenta of the  $\rho^0$  daughter  $\pi^-$  and of the  $\eta'$ , measured in the  $\rho^0$  rest frame.

For  $B^0 \rightarrow \eta' K_L^0$  candidates we require that the cosine of the polar angle of the total missing momentum in the laboratory system be less than 0.95, to reject very forward  $q\bar{q}$  jets. The purity of the  $K_L^0$  candidates reconstructed in

the EMC is further improved by a requirement on the output of a neural network (NN) that takes cluster-shape variables as inputs. The NN was trained on MC signal events and data events in the region  $0.02 < \Delta E < 0.04$  GeV. We check the performance of the NN on data with  $K_L^0$  candidates in the larger  $B^0 \rightarrow J/\psi K_L^0$  data sample.

The average number of candidates found per selected event is between 1.08 and 1.32, depending on the final state. In the case of events with multiple candidates we choose the candidate with the smallest value of a  $\chi^2$  constructed from the deviations from expected values of one or more of the daughter resonance masses, or with the best decay vertex probability for the  $B$ , depending on the decay channel. Furthermore, in the  $\eta' K_L^0$  sample, if several  $B$  candidates have the same vertex probability, we choose the candidate with the  $K_L^0$  information taken from, in order, EMC and IFR, EMC only, or IFR only. From the simulation we find that this algorithm selects the correct-combination candidate in about two-thirds of the events containing multiple candidates.

We obtain the common  $CP$ -violation parameters and signal yields for each channel from a maximum likelihood fit with the input observables  $\Delta E$ ,  $m_{\text{ES}}$ ,  $\mathcal{F}$ , and  $\Delta t$ . The selected sample sizes are given in the first column of Table I. We estimate from the simulation a contribution to the input sample of less than 1.1% of background from other charmless  $B$  decay modes. These events have final states different from the signal, but similar kinematics, and exhibit broad peaks in the signal regions of some observables. We find that the  $B\bar{B}$  background component is needed only for the channels with  $\eta'_\rho\gamma$ . We account for these with a separate component in the probability density function (PDF). For each component  $j$  (signal,  $q\bar{q}$  combinatorial background, or  $B\bar{B}$  background) and tagging category  $c$ , we define a total probability density function for event  $i$  as

$$\mathcal{P}_{j,c}^i \equiv \mathcal{P}_j(m_{\text{ES}}^i) \cdot \mathcal{P}_j(\Delta E^i) \cdot \mathcal{P}_j(\mathcal{F}^i) \cdot \mathcal{P}_j(\Delta t^i, \sigma_{\Delta t}^i; c), \quad (2)$$

except for  $\eta' K_L^0$  for which  $\mathcal{P}_j(m_{\text{ES}}^i)$  is omitted. The factored form of the PDF is a good approximation since linear correlations are small.

We write the extended likelihood function for all events of the decay mode  $d$  as

$$\mathcal{L}_d = \prod_c \exp(-n_c) \prod_i^{N_c} \left[ \sum_j n_j f_{j,c} \mathcal{P}_{j,c}^i \right], \quad (3)$$

where  $n_j$  is the yield of events of component  $j$ ,  $f_{j,c}$  is the fraction of events of component  $j$  for each category  $c$ ,  $n_c = n_{\text{sig}} f_{\text{sig},c} + n_{q\bar{q}} f_{q\bar{q},c} + n_{B\bar{B}} f_{B\bar{B},c}$  is the number of events found by the fitter for category  $c$ , and  $N_c$  is the number of events of category  $c$  in the sample. When combining decay modes we form the grand likelihood  $\mathcal{L} = \prod \mathcal{L}_d$ . We fix both  $f_{\text{sig},c}$  and  $f_{B\bar{B},c}$  to  $f_{B_{\text{nav}},c}$ , the values measured with

the large sample of fully reconstructed  $B^0$  decays into flavor eigenstates ( $B_{\text{flav}}$  sample) [18].

The PDF  $\mathcal{P}_{\text{sig}}(\Delta t, \sigma_{\Delta t}; c)$ , for each category  $c$ , is the convolution of  $F(\Delta t; c)$  [Eq. (1)] with the signal resolution function (sum of three Gaussians) determined from the  $B_{\text{flav}}$  sample. The other PDF forms are the following: the sum of two Gaussians for  $\mathcal{P}_{\text{sig}}(m_{\text{ES}})$  and  $\mathcal{P}_{\text{sig}}(\Delta E)$ ; the sum of three Gaussians for  $\mathcal{P}_{q\bar{q}}(\Delta t; c)$  and  $\mathcal{P}_{B\bar{B}}(\Delta t; c)$ ; an asymmetric Gaussian with different widths below and above the peak for  $\mathcal{P}_j(\mathcal{F})$  [a small “tail” Gaussian is added for  $\mathcal{P}_{q\bar{q}}(\mathcal{F})$ ]; a linear dependence for  $\mathcal{P}_{q\bar{q}}(\Delta E)$  and a fourth-order polynomial for  $\mathcal{P}_{B\bar{B}}(\Delta E)$ ; for  $\mathcal{P}_{q\bar{q}}(m_{\text{ES}})$  and  $\mathcal{P}_{B\bar{B}}(m_{\text{ES}})$  the function  $x\sqrt{1-x^2}\exp[-\xi(1-x^2)]$ , with  $x \equiv 2m_{\text{ES}}/\sqrt{s}$  and  $\xi$  a free parameter [21] and the same function plus a Gaussian, respectively.

For the signal and  $B\bar{B}$  background components we determine the PDF parameters from simulation. We study large control samples of  $B$  decays to charm final states of similar topology to verify the simulated resolutions in  $\Delta E$  and  $m_{\text{ES}}$ , adjusting the PDFs to account for any differences found. The  $q\bar{q}$  background parameters are free to vary in the final fit. Thus, for the six channels listed in Table I, we perform a single fit with 93 free parameters:  $S$ ,  $C$ , signal yields (6),  $\eta'_{\rho\gamma}K^0$   $B\bar{B}$  background yields (2), continuum background yields (6) and fractions (30), background  $\Delta t$ ,  $m_{\text{ES}}$ ,  $\Delta E$ ,  $\mathcal{F}$  PDF parameters (47). The parameters  $\tau$  and  $\Delta m_d$  are fixed to world-average values [19].

We test and calibrate the fitting procedure by applying it to ensembles of simulated experiments with  $q\bar{q}$  events drawn from the PDF into which we have embedded the expected number of signal and  $B\bar{B}$  background events randomly extracted from the fully simulated MC samples. We find negligible bias for  $C$ . For  $S$  we find and apply multiplicative correction factors for bias from dilution due to cross-feed from  $B\bar{B}$  background to signal events equal to 1.03 in the final states  $\eta'_{\rho\gamma}K^0_{\pi^+\pi^-}$ ,  $\eta'_{\eta(\gamma\gamma)\pi\pi}K^0_L$ , and  $\eta'_{\rho\gamma}K^0_{\pi^0\pi^0}$ .

Results from the fit for the signal yields and the  $CP$  parameters  $S$  and  $C$  are presented in Table I. In Fig. 1 we show the projections onto  $m_{\text{ES}}$  and  $\Delta E$  for a subset of the




FIG. 1 (color online). Distributions projected (see text) onto (a)  $m_{\text{ES}}$  and (b)  $\Delta E$  for  $\eta'K_S^0$  candidates, and (c)  $\Delta E$  for  $\eta'K_L^0$  candidates. The solid lines show the full fit result and the dashed lines show the background contributions.

data for which the ratio between the likelihood of signal events and the sum of likelihoods of signal and background events (computed without the variable plotted) exceeds a mode-dependent threshold that optimizes the sensitivity. In Fig. 2 we give the  $\Delta t$  and asymmetry projections of the events selected as for Fig. 1. We measure a correlation of 3.2% between  $S$  and  $C$  in the fit.

We perform several crosschecks of our analysis technique including time-dependent fits for  $B^+$  decays to the charged final states  $\eta'_{\eta(\gamma\gamma)\pi\pi}K^+$ ,  $\eta'_{\rho\gamma}K^+$ , and  $\eta'_{\eta(3\pi)\pi\pi}K^+$ ; fits removing one fit variable at a time; fits without  $B\bar{B}$  PDFs; fits with multiple  $B\bar{B}$  components; fits allowing for nonzero  $CP$  information in  $B\bar{B}$  events; fits with  $C = 0$ . In all cases, we find results consistent with expectation. The value  $S = 0.62 \pm 0.11$  for  $\eta'K_S^0$  differs from our previous measurement  $S = 0.30 \pm 0.14$  [12] due to the improved event reconstruction (with a contribution of +0.08) and selection (+0.12) and to the additional data collected (+0.12). With a model of the data sample changes introduced by our revised event reconstruction and new data, we find that our current result has a statistical probability of 35% (50%) for an assumed true value of  $S$  of 0.61 (0.70).

We have studied the systematic uncertainties arising from several sources (in decreasing order of magnitude): variation of the signal PDF shape parameters within their errors, modeling of the signal  $\Delta t$  distribution, use of  $\Delta t$  signal parameters from the  $B_{\text{flav}}$  sample, interference between the CKM-suppressed  $\bar{b} \rightarrow \bar{u}cd$  amplitude and the favored  $b \rightarrow c\bar{u}d$  amplitude for some tag-side  $B$  decays [22],  $B\bar{B}$  background, SVT alignment, and position and size of the beam spot. The  $B_{\text{flav}}$  sample is used to determine the errors associated with the signal  $\Delta t$  resolutions, tagging efficiencies, and mistag rates. We take the uncertainties in  $\tau_B$  and  $\Delta m_d$  from published measurements [19]. Summing




FIG. 2 (color online). Projections (see text) onto  $\Delta t$  for (a)  $\eta'K_S^0$  and (c)  $\eta'K_L^0$  of the data (points with error bars for  $B^0$  tags in solid circles and  $\bar{B}^0$  tags in empty rectangles), fit function (solid and dashed lines for  $B^0$  and  $\bar{B}^0$  tagged events, respectively), and background function (shaded regions). We show the asymmetry between  $B^0$  and  $\bar{B}^0$  tags for (b)  $\eta'K_S^0$  and (d)  $\eta'K_L^0$ ; the lines represent the fit functions.

all systematic errors in quadrature, we obtain 0.03 for  $S$  and 0.03 for  $C$ .

In conclusion, we have used a sample containing  $1252 \pm 50$  flavor-tagged  $\eta'K^0$  events to measure the time-dependent  $CP$  violation parameters,  $S = 0.58 \pm 0.10 \pm 0.03$  and  $C = -0.16 \pm 0.07 \pm 0.03$ . This sample is 2.1 times as large as that of our previous measurement [12]. Our result for  $S$  is consistent with the world average of  $\sin 2\beta$  measured in  $B^0 \rightarrow J/\psi K_S^0$  [19]. We observe mixing-induced  $CP$  violation in  $B^0$  decays to  $\eta'K^0$  with a significance (systematic uncertainties included) of 5.5 standard deviations. Our result for direct- $CP$  violation is 2.1 standard deviations from zero.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (U.S.A.), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), MEC (Spain), and PPARC (United Kingdom). Individuals have received support from the Marie Curie EIF (European Union) and the A. P. Sloan Foundation.

---

\*Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

†Also with Università della Basilicata, Potenza, Italy

‡Also with IPPP, Physics Department, Durham University, Durham DH1 3LE, United Kingdom

- [1] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **89**, 201802 (2002); K. Abe *et al.* (Belle Collaboration), Phys. Rev. D **66**, 071102(R) (2002).
- [2] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [3] Y. Grossman and M.P. Worah, Phys. Lett. B **395**, 241 (1997); D. Atwood and A. Soni, Phys. Lett. B **405**, 150 (1997); M. Ciuchini *et al.*, Phys. Rev. Lett. **79**, 978 (1997).
- [4] Y. Grossman *et al.*, Phys. Rev. D **68**, 015004 (2003).
- [5] C.-W. Chiang *et al.*, Phys. Rev. D **68**, 074012 (2003); M. Gronau, J. L. Rosner, and J. Zupan, Phys. Lett. B **596**, 107 (2004).
- [6] M. Beneke and M. Neubert, Nucl. Phys. **B675**, 333 (2003).
- [7] D. London and A. Soni, Phys. Lett. B **407**, 61 (1997).
- [8] M. Beneke, Phys. Lett. B **620**, 143 (2005); G. Buchalla *et al.*, J. High Energy Phys. 09 (2005) 074.
- [9] H. Y. Cheng *et al.*, Phys. Rev. D **72**, 014006 (2005); **71**, 014030 (2005); S. Fajfer *et al.*, Phys. Rev. D **72**, 114001 (2005).
- [10] A. R. Williamson and J. Zupan, Phys. Rev. D **74**, 014003 (2006).
- [11] M. Gronau, J. L. Rosner, and J. Zupan, Phys. Rev. D **74**, 093003 (2006).
- [12] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **94**, 191802 (2005).
- [13] K. Abe *et al.* (Belle Collaboration), Phys. Rev. D **72**, 012004 (2005).
- [14] K. F. Chen *et al.* (Belle Collaboration), following Letter, Phys. Rev. Lett. **98**, 031802 (2007).
- [15] B. Aubert *et al.* (*BABAR* Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [16] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **94**, 161803 (2005).
- [17] S. Agostinelli *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **506**, 250 (2003).
- [18] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. D **66**, 032003 (2002).
- [19] Y.-M. Yao *et al.* (Particle Data Group), J. Phys. G **33**, 1 (2006).
- [20] B. Aubert *et al.* (*BABAR* Collaboration), Phys. Rev. Lett. **92**, 061801 (2004).
- [21] H. Albrecht *et al.* (ARGUS Collaboration), Phys. Lett. B **241**, 278 (1990).
- [22] O. Long *et al.*, Phys. Rev. D **68**, 034010 (2003).