

Journal of Medicinally Active Plants

Volume 3
Issue 1 Vol 3 Issues 1-4

January 2014

Chemotyping and Determination of Antimicrobial, Insecticidal, and Cytotoxic Properties of Wild-Grown Cannabis sativa from Nepal

Prabodh Satyal
University of Alabama in Huntsville, prabodhsatyal@gmail.com

William N. Setzer
University of Alabama - Huntsville, wsetzer@chemistry.uah.edu

Follow this and additional works at: <https://scholarworks.umass.edu/jmap>



Part of the [Plant Sciences Commons](#)

Recommended Citation

Satyal, Prabodh and William N. Setzer. 2014. "Chemotyping and Determination of Antimicrobial, Insecticidal, and Cytotoxic Properties of Wild-Grown Cannabis sativa from Nepal." *Journal of Medicinally Active Plants* 3, (1):9-16.

DOI: <https://doi.org/10.7275/R58W3B8V>

<https://scholarworks.umass.edu/jmap/vol3/iss1/3>

This Article is brought to you for free and open access by ScholarWorks@UMass Amherst. It has been accepted for inclusion in *Journal of Medicinally Active Plants* by an authorized editor of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

Journal of Medicinally Active Plants

Volume 3

Issue 1 Vol 3 Issues 1-4

December 2014

Chemotyping and Determination of Antimicrobial, Insecticidal, and Cytotoxic Properties of Wild-Grown *Cannabis sativa* from Nepal

Prabodh Satyal

University of Alabama in Huntsville, prabodhsatyal@gmail.com

William N. Setzer

University of Alabama - Huntsville, wsetzer@chemistry.uah.edu

Follow this and additional works at: <http://scholarworks.umass.edu/jmap>

Recommended Citation

Satyal, Prabodh, William N. Setzer. 2014. "Chemotyping and Determination of Antimicrobial, Insecticidal, and Cytotoxic Properties of Wild-Grown *Cannabis sativa* from Nepal," *Journal of Medicinally Active Plants* 3 (Vol 3 Issues 1-4):9-16.

DOI: <https://doi.org/10.7275/R58W3B8V>

Available at: <http://scholarworks.umass.edu/jmap/vol3/iss1/3>

This Article is brought to you for free and open access by ScholarWorks@UMass Amherst. It has been accepted for inclusion in Journal of Medicinally Active Plants by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

Chemotyping and Determination of Antimicrobial, Insecticidal and Cytotoxic Properties of Wild-Grown *Cannabis sativa* from Nepal

Prabodh Satyal and William N. Setzer*

Department of Chemistry, University of Alabama in Huntsville, Huntsville, AL 35899, USA

*Corresponding author: wsetzer@chemistry.uah.edu

Manuscript received: September 21, 2014

Keywords: Caryophyllene, cannabidiol, *Cannabis indica*, cluster analysis, humulene, tetrahydrocannabinol.

ABSTRACT

Cannabis sativa was collected from a wild-growing population in Biratnagar, Nepal. The essential oil was obtained by hydrodistillation and analyzed by gas chromatography – mass spectrometry. A total of 107 constituents were identified in the oil accounting for 94.2% of the composition. This Nepalese chemotype is characterized by a predominance of sesquiterpenoids (68.1%) dominated by (*E*)-caryophyllene (20.4%), α -humulene (7.0%), and α -bisabolol (5.8%), but a paucity of monoterpene hydrocarbons (0.9%). In particular, neither myrcene nor terpinolene were detected. The oil in the Nepalese *Cannabis* plants did contain small amounts of cannabidiol (1.6%), cannabichromene (0.2%) and Δ^9 -tetrahydrocannabinol (0.4%). The essential oil from Nepalese *Cannabis sativa* was screened for antimicrobial, cytotoxic, larvicidal, and insecticidal activity, and it appears as though *C. sativa* is relatively non-toxic.

INTRODUCTION

Cannabis sativa L. is one of the three species of genus *Cannabis* that includes *C. sativa*, *C. indica*, and *C. ruderalis* (Guy *et al.*, 2004). *C. sativa*, commonly known as “hemp” or “marijuana” in English and “bhang” or “ganja” in Nepali language, is an annual herbaceous, dioecious plant (Kriese *et al.*, 2004). The origin of wild *Cannabis* is not known with certainty, but the plant, which appears to have been cultivated in northern China since 4000 BC

(Mabberley, 2008), is widely distributed and cultivated throughout the world (Nigam *et al.*, 1965).

Some confusion and controversy regarding the taxonomy of *Cannabis* does exist. Small and Cronquist (1976) had proposed a monotypic genus, *C. sativa* with two subspecies, *sativa* and *indica*, while several others have recognized three different species, *C. sativa*, *C. indica*, and *C. ruderalis* (Schultes *et al.*, 1974; Anderson, 1974, 1980; Emboden, 1974). Recent, genetic, morphological, and biochemical investigations have helped to delineate the three species of *Cannabis* (Hillig and Mahlberg, 2004; Hillig, 2004, 2005). *C. sativa* generally has lower levels of Δ^9 -tetrahydrocannabinol (THC) than cannabidiol (CBD), while *C. indica* has less CBD than THC (Hillig and Mahlberg, 2004). Wide-leaf varieties of *C. indica* have shown relatively high ratio of guaiol, γ -eudesmol, and β -eudesmol, while the narrow-leaf varieties of *C. indica* have a relatively high ratio of (*E*)- β -farnesene (Hillig, 2004).

The essential oil of *Cannabis* has demonstrated antimicrobial activity (Novak *et al.*, 2001; Nissen *et al.*, 2010) and mosquito larvicidal activities (Pavela, 2009). In the current study, the composition, chemotype, and bioactivities of the essential oil from a wild-growing *Cannabis* plant, tentatively identified as *C. sativa*, from Biratnagar, Nepal, were investigated.

MATERIALS AND METHODS

Plant material. Flowering *Cannabis* plants collected from city of Biratnagar (26°28' N, 87°16' E,

72 m above sea level) in Morang district in Koshi Zone in Nepal on 18 May 2011 were used in this study. The plant material was identified by Tilak Gautam as male, narrow-leafed *Cannabis sativa* L. plants based on the plant morphology. A voucher specimen (4304) has been deposited in the herbarium of the Tribhuvan University, Post-Graduate Campus, Botany Department, Biratnagar.

The essential oil was extracted from fresh leaf samples (100 g) that were crushed and hydrodistilled using a Clevenger type apparatus for 4 h. The clear pale yellow essential oil (0.5 g) produced was stored at 4°C until analyzed.

Constituent identification. The essential oil from the *Cannabis* plant was analyzed by GC-MS using an Agilent 6890 GC with Agilent 5973 mass selective detector [MSD, operated in the EI mode (electron energy = 70 eV), with scan range = 40-400 amu, and scan rate = 3.99 scans/sec], and an Agilent ChemStation data system. The GC column was an HP-5ms fused silica capillary with a (5% phenyl)-polymethylsiloxane stationary phase, a film thickness of 0.25 µm, a length of 30 m, and an internal diameter of 0.25 mm. The carrier gas was helium with a column head pressure of 48.7 kPa and a flow rate of 1.0 mL/min. The injector temperature was 200°C, and detector temperature was 280°C. The GC oven temperature was programmed with an initial temperature of 40°C, hold for 10 min, followed by an increase in temperature at 3°C min to 200°C, and then increased at 2°/min to 220°C. A 1% w/v solution of each sample in CH₂Cl₂ was prepared, and 1 µL of the sample was injected using a 10:1 split ratio.

Identification of the oil components was based on the retention indices determined by reference to a homologous series of *n*-alkanes and by comparison of the mass spectral fragmentation patterns with those reported in the literature (Adams, 2007) and stored on the MS library [NIST database (G1036A, revision D.01.00)/ChemStation data system (G1701CA, version C.00.01.080)]. The percentages of each component were determined as raw percentages based on total ion current without standardization.

Antimicrobial activity. The essential oil was screened for anti-microbial activity against Gram-positive bacteria, *Bacillus cereus* (ATCC No. 14579), and *Staphylococcus aureus* (ATCC No. 29213), Gram-negative bacteria, *Pseudomonas aeruginosa* (ATCC No. 27853) and *Escherichia coli* (ATCC No. 10798), and the fungi *Candida albicans* (ATCC No.10231) and *Aspergillus niger* (ATCC No. 16888). Minimum inhibitory concentrations (MICs) were determined using the microbroth dilution technique as previously described (Satyal *et al.*, 2012).

Toxicity. The essential oil was screened for cytotoxicity using human MCF-7 breast adenocarcinoma cells (ATCC No. HTB-22), along with tests of lethality using brine shrimp (*Artemia salina*), nematicidal activity using the nematode (*Caenorhabditis elegans*), larvicidal activity using glass worm (*Chaoborus plumicornis*), and insecticidal activity using termites (*Reticulitermes virginicus*) and fruit flies (*Drosophila melanogaster*) as previously described (Satyal *et al.*, 2012). LC₅₀ values were determined using the Reed-Muench method (Reed and Muench, 1938).

Hierarchical cluster analysis. To develop a hierarchical cluster analysis, a selection of 68 *Cannabis* essential oil compositions, taken from published literature (Ross and ElSohly, 1996; Mediavilla and Steinemann, 1997; Novak *et al.*, 2001; Rothschild *et al.*, 2005; Bertoli *et al.*, 2010; Nissen *et al.*, 2010; Fishedick *et al.*, 2010), were treated as operational taxonomic units (OTUs). The percentage composition of 16 terpenoid essential oil components (α -pinene, β -pinene, δ -3-carene, myrcene, limonene, β -phellandrene, (*Z*)- β -ocimene, (*E*)- β -ocimene, terpinolene, α -bergamotene, (*E*)-caryophyllene, α -humulene, (*E*)- β -farnesene, β -selinene, selina-3,7(11)-diene, and caryophyllene oxide) were used to determine the chemical relationship among the *Cannabis* essential oil samples by the agglomerative hierarchical cluster (AHC) analysis using the XLSTAT software, version 2014.4.09. Pearson's correlation was selected as a measure of similarity, and the unweighted, pair-group method with arithmetic average (UPGMA) was used for cluster definition and to develop a dendrogram for the *Cannabis* selections.

RESULTS

The essential oil of *C. sativa* from Nepal yielded a total of 112 components of which 107 were identified, accounting for 94.2% of the total composition (Table 1). The oil was rich in sesquiterpene hydrocarbons (46.8%) and oxygenated sesquiterpenoids (21.3%), dominated by (*E*)-caryophyllene (20.4%), α -humulene (7.0%), α -bisabolol (5.8%), and caryophyllene oxide (3.8%).

In the screening of the Nepalese *C. sativa* essential oil for antimicrobial activity and toxicity, the oil showed notable activity in the brine shrimp lethality test with a $LC_{50} = 13.6 \mu\text{g/mL}$ (Table 2). Tests for antimicrobial activity indicated the essential oil had essentially no effect on the tested microorganisms.

The hierarchical cluster analysis of the *C. sativa* essential oil compositions indicated the essential oil from the Nepalese plants was chemically distinct from the essential oil of other *Cannabis* plants (Figure 1). Nepalese *Cannabis* oil was high in caryo-phyllene and deficient in myrcene and terpinolene as compared with other *Cannabis sativa* essential oils.

DISCUSSION

The lack of appreciable antimicrobial activity exhibited by Nepalese *C. sativa* is consistent with previous studies that show only marginal activity of *Cannabis* oils against Gram-positive bacteria (Novak *et al.*, 2001; Nissen *et al.*, 2010). In addition, the marginal toxicity of Nepalese *C. sativa* against nematodes, glass worms, fruit flies, and termites is consistent with the reported marginal activity of Canadian *C. sativa* against mosquito (*Culex quinquefasciatus*) larvae ($LC_{50} = 127 \mu\text{g/mL}$) (Pavela, 2009). Apparently, *C. sativa* essential oil is relatively non-toxic and thus the lack of insecticidal activity of *C. sativa* oil is not surprising. The major component in the oil, (*E*)-caryophyllene, has shown only limited insecticidal activity against *Aedes aegypti* (Silva *et al.*, 2008), *Rhyzopertha dominica*, *Sitophilus oryzae*, *Tribolium castaneum* (Zeng *et al.*,

2010) or *Lycoriella ingenua* (Park *et al.*, 2008). Likewise, (*E*)-caryophyllene is neither notably anti-bacterial (Palazzo *et al.*, 2009) nor cytotoxic (Schmidt *et al.*, 2006; Wright *et al.*, 2007).

The activity of the *C. sativa* essential oil in the brine shrimp lethality test is comparable to many other essential oils (Werka *et al.*, 2007; Setzer *et al.*, 2008) and may be attributed to the major sesquiterpene components (*E*)-caryophyllene ($LC_{50} = 0.23 \mu\text{g/mL}$) and α -humulene ($LC_{50} = 0.75 \mu\text{g/mL}$).

In contrast to previously reported *Cannabis* essential oils (Ross and ElSohly, 1996; Mediavilla and Steinemann, 1997; Novak *et al.*, 2001; Rothschild *et al.*, 2005; Bertoli *et al.*, 2010; Nissen *et al.*, 2010), the oil from Nepal was poor in monoterpene hydrocarbons. In particular, the Nepalese *C. sativa* oil had no detectable quantities of myrcene or terpinolene, normally abundant in *C. sativa* (Ross and ElSohly, 1996; Mediavilla and Steinemann, 1997; Novak *et al.*, 2001; Rothschild *et al.*, 2005; Bertoli *et al.*, 2010; Nissen *et al.*, 2010). The sample collected from Biratnagar, Nepal, was identified as *C. sativa* based on plant and leaf morphology, and the cannabidiol (CBD) to Δ^9 -tetrahydrocannabinol (THC) ratio is consistent with *C. sativa* (CBD/THC = 3.9) (Hillig and Mahlberg, 2004). The absence of myrcene and terpinolene, the low α -pinene concentration, and the relatively high concentration of (*E*)- β -farnesene in the Nepalese sourced essential oil, however, is consistent with *C. indica*. *C. sativa* essential oils generally contain high concentrations of myrcene and α -pinene (Hillig, 2004).

The hierarchical cluster analysis of essential oil compositions revealed the essential oil from Nepal is chemically distinct from other *Cannabis* essential oils and represents a new chemotype, rich in caryo-phyllene, but deficient in myrcene and terpinolene. Of the essential oils examined, two other samples of *C. indica*, also showed no terpinolene and small concentrations of myrcene (samples Bedropuur and AN) (Fischedick *et al.*, 2010).

Table 1. Composition of the essential oil of wild *Cannabis sativa* from Biratnagar, Nepal.

RI	Compound	%	RI	Compound	%
777	(2Z)-Pentenol	0.1	1346	4-Hydroxybenzyl alcohol	0.2
799	Hexanal	0.2	1349	α -Cubebene	tr
830	2-Furaldehyde	0.2	1357	Eugenol	0.2
851	(2E)-Hexenal	0.9	1366	Cyclosativene	0.1
853	(3Z)-Hexenol	4.7	1375	α -Copaene	0.1
866	(2E)-Hexenol	0.6	1391	7- <i>epi</i> -Sesquithujene	0.2
869	1-Hexanol	0.6	1408	(Z)-Caryophyllene	0.3
893	2-Heptanone	0.1	1410	Sesquithujene	0.2
897	Cyclohexanone	0.4	1422	(E)-Caryophyllene	20.4
905	Heptanal	0.1	1430	β -Copaene	0.1
940	α -Pinene	0.1	1438	α - <i>trans</i> -Bergamotene	3.0
954	Camphene	tr	1445	6,9-Guaiadiene	0.2
964	Benzaldehyde	0.1	1450	Unidentified	0.8
974	Sabinene	tr	1456	α -Humulene	7.0
979	β -Pinene	0.1	1460	(E)- β -Farnesene	3.3
989	6-Methyl-5-hepten-2-one	0.1	1470	4,5-di- <i>epi</i> -Aristolochene	0.1
1009	(3Z)-Hexenyl acetate	0.2	1478	γ -Muurolole	0.6
1029	Limonene	0.3	1481	γ -Curcumene	0.2
1032	1,8-Cineole	3.8	1485	γ -Selinene	0.4
1033	Benzyl alcohol	0.7	1488	β -Selinene	2.3
1039	(Z)- β -Ocimene	tr	1496	α -Selinene	2.3
1043	Phenylacetaldehyde	0.1	1501	α -Muurolole	0.2
1049	(E)- β -Ocimene	0.3	1504	(E,E)- α -Farnesene	0.1
1058	γ -Terpinene	0.1	1511	β -Bisabolene	2.6
1068	<i>cis</i> -Sabinene hydrate	1.5	1526	β -Sesquiphellandrene	2.8
1072	1-Octanol	0.1	1529	(E)- <i>iso</i> - γ -Bisabolene	0.3
1088	Fenchone	0.3	1531	10- <i>epi</i> -Cubebol	0.2
1095	Methyl benzoate	0.1	1538	α -Cadinene	0.1
1098	<i>trans</i> -Sabinene hydrate	0.9	1545	Unidentified	3.2
1101	Linalool	0.2	1554	Unidentified	0.6
1105	Nonanal	0.1	1564	(E)-Nerolidol	0.5
1113	2-Phenylethyl alcohol	0.4	1569	Caryophyllenyl alcohol	0.2
1120	<i>trans</i> -Pinene hydrate	0.4	1580	<i>trans</i> -Sesquisabinene hydrate	0.5
1138	<i>trans</i> -Pinocarveol	0.1	1584	Caryophyllene oxide	3.8
1141	<i>cis</i> -Pinene hydrate	0.2	1600	Guaiol	1.4
1143	4-Ketoisophorone	0.1	1607	Humulene epoxide II	1.3
1147	Ipsdienol	0.4	1621	10- <i>epi</i> - γ -Eudesmol	0.5
1149	Myrcenone	0.2	1632	(E)-Sesquilandulol	2.0
1155	(3Z)-Nonenol	0.1	1637	Caryophylla-4(12),8(13)-dien-5-ol	0.7
1158	β -Pinene oxide	0.2	1646	α -Muurolol (= Torreyol)	0.1
1162	Pinocarvone	0.2	1652	β -Eudesmol	0.4
1165	Borneol	0.2	1655	α -Eudesmol	0.8
1166	δ -Terpineol	0.1	1657	Selin-11-en-4 α -ol	0.4
1177	Terpinen-4-ol	0.6	1660	Intermedeol	0.7
1185	<i>p</i> -Cymen-8-ol	tr	1669	Bulnesol	1.2
1187	(3Z)-Hexenyl butanoate	0.2	1672	14-Hydroxy-9- <i>epi</i> -(E)-caryophyllene	0.8
1190	α -Terpineol	0.5	1686	α -Bisabolol	5.8
1193	Hexyl butanoate	0.2	1689	Unidentified	0.9
1208	Verbenone	0.1	2100	(E)-Phytol	1.0
1232	(3Z)-Hexenyl 2-methylbutanoate	tr	2419	Cannabidiol	1.6
1238	Neral	0.1	2427	Cannabichromene	0.2
1241	Carvone	tr	2518	Δ^9 -Tetrahydrocannabinol	0.4
1249	<i>p</i> -Anisaldehyde	0.1	-	Total Identified	94.2
1252	Geraniol	tr	-	Monoterpene hydrocarbons	0.9
1260	2-Hydroxybenzenemethanol	0.2	-	Oxygenated monoterpenoids	10.1
1269	Unidentified	0.3	-	Sesquiterpene hydrocarbons	46.8
1296	Perilla alcohol	tr	-	Oxygenated sesquiterpenoids	21.3
1312	<i>p</i> -Vinylguaiaicol	1.1	-	Benzenoid aromatics	3.1
1325	(3Z)-Hexenyl tiglate	tr	-	Cannabinoids	2.2
1339	Piperitenone	0.1	-	Others	9.7

References

- Adams, R.P. 2007. *Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry*, 4th Ed. Allured Publishing Corporation. Carol Stream, IL, USA.
- Anderson, L.C. 1974. A study of systematic wood anatomy in *Cannabis*. Harvard University Botanical Museum Leaflets 24:29-36.
- Anderson, L.C. 1980. Leaf variation among *Cannabis* species from a controlled garden. Harvard University Botanical Museum Leaflets 28:61-69.
- Bertoli, A., S. Tozzi, L. Pistelli, and L.G. Angelini. 2010. Fibre hemp inflorescences: From crop-residues to essential oil production. *Industrial Crops and Products* 32:329-337.
- Emboden, W.A. 1974. *Cannabis* – a polytypic genus. *Economic Botany* 28: 304-310.
- Fischedick, J.T., A. Hazenkamp, T. Erkelens, Y.H. Choi, and R. Verpoorte. 2010. Metabolic fingerprinting of *Cannabis sativa* L., cannabinoids and terpenoids for chemotaxonomic and drug standardization purposes. *Phytochemistry* 71:2058-2073.
- Guy, G.W., B.A. Whittle, and P. Robson. 2004. *The Medicinal Uses of Cannabis and Cannabinoids*. Pharmaceutical Press, London, UK.
- Hillig, K.W. 2004. A chemotaxonomic analysis of terpenoid variation in *Cannabis*. *Biochemical Systematics and Ecology* 32:875-891.
- Hillig, K.W. 2005. Genetic evidence for speciation in *Cannabis* (Cannabaceae). *Genetic Resources and Crop Evolution* 52:161-180.
- Hillig, K.W. and P.G. Mahlberg. 2004. A chemotaxonomic analysis of cannabinoid variation in *Cannabis* (Cannabaceae). *American Journal of Botany* 91:966-975.
- Kriese, U., E. Schumann, W. Weber, M. Beyer, L. Bruhl, and B. Matthaus. 2004. Oil content, tocopherol composition and fatty acid patterns of the seeds of 51 *Cannabis sativa* L. genotypes. *Euphytica* 137:339-351.
- Mabberley, D.J. 2008. *Mabberley's Plant Book*, 3rd Ed. Cambridge University Press, UK.
- Mediavilla, V. and S. Steinemann. 1997. Essential oil of some hemp genotypes. *Journal of the International Hemp Association*, 4(2):80-82.
- Nigam, M., K. Handa, I. Nigam, and L. Levi. 1965. The essential oils of marijuana composition of genuine Indian *Cannabis sativa* L. *Canadian Journal of Chemistry* 43:3372-3376.
- Nissen, L., A. Zatta, I. Stefanini, S. Grandi, B. Sgorbati, B. Biavati, and A. Monti. 2010. Characterization and antimicrobial activity of essential oils of industrial hemp varieties (*Cannabis sativa* L.). *Fitoterapia* 81:413-419.
- Novak, J., K. Zitterl-Eglseer, S.G. Deans, and C.M. Franz. 2001. Essential oils of different cultivars of *Cannabis sativa* L. and their antimicrobial activity. *Flavour and Fragrance Journal* 16:259-262.
- Palazzo, M.C., H.L. Wright, B.R. Agius, B.S. Wright, Moriarity, D.M. Haber, W.A. and W.N. Setzer. 2009. Chemical compositions and biological activities of leaf essential oils of six species of Annonaceae from Monteverde, Costa Rica. *Records of Natural Products* 3:153-160.
- Park, I.K, J.N. Kim, Y.S. Lee, S.G. Lee, Y.J. Ahn, and S.C. Shin. 2008. Toxicity of plant essential oils and their components against *Lycoriella ingenua* (Diptera: Sciaridae). *Horticultural Entomology* 101:139-144.
- Pavela, R. 2009. Larvicidal property of essential oils against *Culex quinquefasciatus* Say (Diptera: Culicidae). *Industrial Crops and Products* 30:311-315.
- Reed, L.J. and H. Muench. 1938. A simple method of estimating fifty percent endpoints. *American Journal of Hygiene* 27:493-497.
- Ross, S.A., and M.A. El Sohly. 1996. The volatile oil composition of fresh and air-dried buds of *Cannabis sativa*. *Journal of Natural Products* 59:49-51.
- Rothschild, M., G. Bergström, and S. Wängberg. 2005. *Cannabis sativa*: Volatile compounds from pollen and entire male and female plants of two variants, Northern Lights and Hawaiian Indica. *Botanical Journal of the Linnean Society* 147:387-397.
- Satyal, P., K.E. Woods, N.S. Dosoky, S. Neupane, W.N. Setzer. 2012. Biological activities and volatile constituents of *Aegle marmelos* (L.) Corrêa from Nepal. *Journal of Medicinally Active Plants* 1(3):114-122.

- Schmidt, J.M., J.A. Noletto, B. Vogler, and W.N. Setzer. 2006. Abaco bush medicine: Chemical composition of the essential oils of four aromatic medicinal plants from Abaco Island, Bahamas. *Journal of Herbs, Spices & Medicinal Plants* 12(3):43-65.
- Schultes, R.E., W.M. Klein, T. Plowman, and T.E. Lockwood. 1974. *Cannabis*: an example of taxonomic neglect. *Harvard University Botanical Museum Leaflets* 23:337-367.
- Setzer, W.N., G. Park, B.R. Agius, S.L. Stokes, T.M. Walker, and W.A. Haber. 2008. Chemical compositions and biological activities of leaf essential oils of twelve species of *Piper* from Monteverde, Costa Rica. *Natural Product Communications* 3:1367-1374.
- Silva, W.J., G.A.A. Dória, R.T., Maia, R.S. Nunes, G.A. Carvalho, A.F. Blank, P.B. Alves, R.M. Marçal, and S.C.H. Cavalcanti. 2008. Effects of essential oils on *Aedes aegypti* larvae: Alternatives to environmentally safe insecticides. *Bioresource Technology* 99:3251-3255.
- Small, E. and Cronquist, A. 1976. A practical and natural taxonomy for *Cannabis*. *Taxon* 25:405-435.
- Werka, J.S., A.K. Boehme, and W.N. Setzer. 2007. Biological activities of essential oils from Monteverde, Costa Rica. *Natural Product Communications* 2:1215-1219.
- Wright, B.S., A. Bansal, D.M. Moriarity, S. Takakuand, W.N. Setzer. 2007. Cytotoxic leaf essential oils from Neotropical Lauraceae: Synergistic effects of essential oil components. *Natural Product Communications* 2:1241-1244.
- Zeng, L., C.Z. Lao, Y.J. Cen, and G.W. Liang, 2010. Study on the insecticidal activity compounds of the essential oil from *Syzygium aromaticum* against stored grain insect pests. *Julius-Kühn-Archiv* 425:766-771.

