2005

Isotropy of the early universe from CMB anisotropies

EP Donoghue
donoghue@physics.umass.edu

JF Donoghue
donoghue@physics.umass.edu

Follow this and additional works at: https://scholarworks.umass.edu/physics_faculty_pubs

Part of the [Physics Commons](#)

Recommended Citation

Donoghue, EP and Donoghue, JF, "Isotropy of the early universe from CMB anisotropies" (2005). *PHYSICAL REVIEW D*. 98.
Retrieved from https://scholarworks.umass.edu/physics_faculty_pubs/98

This Article is brought to you for free and open access by the Physics at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Physics Department Faculty Publication Series by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

Isotropy of the early universe from CMB anisotropies

Evan P. Donoghue ^{a,b} and John F. Donoghue^a

^a Department of Physics
University of Massachusetts
Amherst, MA 01003

^b University of Notre Dame,
Notre Dame, IN 46556

February 2, 2008

Abstract

The acoustic peak in the CMB power spectrum is sensitive to causal processes and cosmological parameters in the early universe up to the time of last scattering. We provide limits on correlated spatial variations of the peak height and peak position and interpret these as constraints on the spatial variation of the cosmological parameters (baryon density, cold dark matter density and cosmological constant as well as the amplitude and tilt of the original fluctuations). We utilize recent work of Hansen, Banday and Górski (HBG) who have studied the spatial isotropy of the power spectrum as measured by WMAP by performing the power spectrum analysis on smaller patches of the sky. We find that there is no statistically significant correlated asymmetry of the peak. HBG have also provided preliminary indications of a preferred direction in the lower angular momentum range ($\ell \sim 2-40$) and we show how possible explanations of this asymmetry are severely constrained by the data on the acoustic peak. Finally we show a possible non-gaussian feature in the data, associated with a difference in the northern and southern galactic hemispheres.

1 Introduction

One of the foundations of our description of the universe is the principle of isotropy and homogeneity - the statement that the universe is the same in all directions and locations. The isotropy of the universe can be explained by the theory of inflation, in which all of the observed universe descends from a single causally connected patch of the early universe. However, inflation also contains the ingredients for generating anisotropies on various scales. This is manifest most importantly in the temperature anisotropies of the cosmic microwave background (CMB). The temperature anisotropies are generated by the quantum fluctuations of a scalar field during the inflationary epoch. Other scalar fields could also have quantum fluctuations and these could in principle influence the distribution of matter in different portions of the universe. It is possible that these could leave a residual “tilt” to the universe[1].

Recently Hansen, Banday and Górski (HBG) [2, 3, 4] have explored novel tests of isotropy by studying the WMAP[5] power spectrum extracted from the CMB using patches in different directions of the sky. This is an important test because the CMB provides the largest lever arm for the study of spatial asymmetries. In fact, in the lower angular range of $\ell \sim 2-40$ HBG find preliminary indications (at about the 2-3 sigma level) for non-zero differences in the power spectrum in hemispheres oriented along galactic colatitude and longitude (80° , 57°), close to the ecliptic pole. This directional asymmetry seems to be connected with the orientation of the quadrupole and octopole [6]. If this signal proves to be real, it could be an important indication of a fundamental tilt in the properties of the universe. In addition, a lack of isotropy has been seen in tests for non-gaussian behavior. Eriksen et al [7] and Park [8] were the first to find a difference between the northern and southern galactic hemispheres in tests for non-gaussianity, and this hemisphere difference has subsequently been confirmed by other measures [9, 10, 11, 12].

HBG also studied the height and the location of the first acoustic peak on different patches of the sky, finding no obvious anisotropy within the expected cosmic variance. We use the HBG data to study the properties of the acoustic peak in more detail. The region of the acoustic peak is special because it is sensitive to causal processes in the early universe up to the time of last scattering. These processes depend on the cosmological parameters and any gradient of the parameters would yield a specific pattern between the peak height and peak position, i.e. both the height and the position would change in a correlated way. These correlations could reveal a sig-

nal of anisotropy that is not visible when looking at the height or position individually. We explore the HBG data for the possible presence of these correlated variations. Interpreting the results as variations in the cosmological parameters, we provide upper bounds on their possible variation. When we allow for an arbitrary correlated variation, we will find a weak correlation to a directions on the sky, but we will argue that it is not statistically significant. We also use the properties of the acoustic peak as a probe of the possible physics that could produce the low ℓ signal found by HBG. The isotropy of the peak is a strong constraint on the physics that could produce the low ℓ signal. If one studies only the variation of a single cosmological parameter, only a difference of the spectral tilt could explain the low- ℓ anomaly while being consistent with the properties of the acoustic peak. Finally we describe an unusual behavior in the distribution of the properties of the acoustic peak, associated with a difference between the northern and southern galactic hemispheres. This may possibly be related to the north-south difference which has been found in tests for non-gaussianity.

2 Characteristic patterns of individual variations

The acoustic peak is categorized by its location in ℓ and by its height. However, these are not fully independent variables in terms of their dependence on the cosmological parameters. If one were modify the baryon density, for example, there would be a correlated variation in both the height and location of the peak. In this section we describe the nature of these correlations.

Consider the variation of the cosmological parameters. In all cases, we adopt as the central value that found by the WMAP team. In addition, we define a parameter

$$\epsilon_i = \frac{p_i - p_i^{\text{WMAP}}}{p_i^{\text{WMAP}}} \quad (1)$$

where p_i is any of the cosmological parameters and the superscript WMAP denotes its value measured by WMAP.

The dependence of the power spectrum on all the cosmological parameters can be calculated using CMBFAST[13]. Here we focus on the properties of the acoustic peak. It is important to use the version of CMBFAST that does not implement the COBE normalization. Otherwise, changes in the power at low values of ℓ would have an extra impact on the amplitude of the acoustic peak by rescaling the overall power spectrum. This is especially relevant for variations of the spectral tilt and the cosmological constant, which appreciably modify the power at lower values of ℓ . Taking small variations of

the cosmological parameters with modify the location of the acoustic peak, ℓ_p , and the height of the peak, h_p , from the values found by WMAP, called ℓ_0 and h_0 . In each case we keep $\Omega_{\text{tot}} = 1$ as we vary the parameter. For small variations the changes will be linear in ϵ_i , so that we parameterize the results by

$$\begin{aligned}\ell_p &= \ell_0 + a_i \epsilon_i \\ h_p &= h_0 + b_i \epsilon_i\end{aligned}\tag{2}$$

These can be combined to describe the correlation characteristic of each parameter, using

$$\ell_p - \ell_0 = r_i(h_p - h_0)\tag{3}$$

where

$$r_i = \frac{a_i}{b_i}\tag{4}$$

The value of the different coefficients are given in Table 1. The correlated variations in the (h, ℓ) plane are shown in Figure 1 for the different parameters.

parameter	Baryon	CDM	Λ	amplitude	tilt
a	10	0	-22	-16	37
b	1875	-2381	0	5600	-5491

Table 1: Parameters describing the variation of the height and location of the acoustic peak when one changes the energy density of baryons, the energy density of cold dark matter, the value of the cosmological constant or the value of the spectral tilt.

3 Measures of Isotropy

Hansen et. al. (HBG)[2] have studied the WMAP power spectrum on 34 patches of the sky, each with an opening angle of 19° . These cover most of the sky, avoiding the galactic plane where foreground contaminations could distort the signal. The HBG data on the location and height of the acoustic peak is displayed in Fig 1. The correlation with position on the sky is given in Table 2. The expected spread in these values has been estimated by HBG through the variance found in 512 simulated skies with gaussian distributed temperature fluctuations using the WMAP power spectrum. This yields a variance in the ℓ direction of $\sigma_\ell = 6.5$ and in the height of $\sigma_h = 300$.

Figure 1: The lines show the correlated variation of the height and location in ℓ of the acoustic peak as one varies each of the indicated quantities. Also shown are the HBG data for the height and location of the peak for 34 patches of the sky.

A comparison of the average values of the position and height of the peak over opposite hemispheres is subject to large fluctuations as individual patches near the borders of the hemispheres are counted first in one hemisphere and then in the other as the orientation of the hemisphere changes slightly in direction. Therefore we seek a measure comparing directions that is not as sensitive to the border between the hemispheres. To this end we consider a possible dipole asymmetry. Theoretically a dipole is often the leading multipole for a spatial asymmetry [1, 14, 15, 16, 17, 19, 20, 21]. A dipole tilt in the Universe can also be hidden beneath the overall dipole seen in the CMB temperature, which is usually attributed to our motion with respect to the CMB, but which can also have other components[1, 15].

Consider a reference direction given by polar angles θ , ϕ and let γ be the angle between this reference direction and a point in the sky. For a given parameter p_i , a correlated dipole variation centered on this reference direction has the form

$$\begin{aligned}\ell(\theta, \phi) &= \ell_0 + a_i \eta \cos \gamma \\ h(\theta, \phi) &= h_0 + b_i \eta \cos \gamma\end{aligned}\tag{5}$$

where a_i , b_i are the constants determined in the previous section and η is the amplitude of a possible variation.

For each of the cosmological parameters, we allow the reference direction to vary and search for a direction that gives the maximum possible dipole asymmetry.¹ While in each case a preferred direction can be found, the resulting asymmetries are not statistically significant, as we discuss below. The amplitude and standard deviation for each variation is shown in Table 3. For each of the parameters, we can convert our lack of a signal into a bound. For the direction that maximizes the asymmetry we take the 2σ upper bound on the amplitude η . The results are given in Table 3.

We also searched for a dipole anisotropy with an arbitrary correlation between the height and the peak location. We do this by allowing the relative size of a_i and b_i to be arbitrary. The result was an anisotropy centered on the direction ($\theta = 107^\circ \pm 34^\circ$, $\phi = 30^\circ \pm 42^\circ$) with a small amplitude that is nominally 1.6 standard deviations from zero and a correlation

$$r = \frac{a}{b} = -0.3 \pm 2.0 \quad (6)$$

The ratio r has a large uncertainty but favors a signal which is mostly in the overall height of the amplitude. Numerically, the amplitude corresponds to $b\eta = 148 \pm 93$, which is a signal of less than 3% and a nominal statistical likelihood of 1.6 standard deviations. Although this likelihood is of marginal significance, the true significance is weaker because we have allowed an arbitrary correlation by accepting any ratio r . To demonstrate this, we performed a similar search on 100 sets of simulated data, generated randomly with a gaussian weight and the same standard deviations given by HBG. If we allow the signal to have an arbitrary value of r and an arbitrary direction, we find that 85% of the simulated data sets also have a non-zero amplitude of at least 1.6 standard deviations. We conclude that the signal is not statistically significant.

4 Constraints on the possible asymmetry at low ℓ

In the previous section, we studied the constraints imposed by the properties of the acoustic peak. However, some of the parameters can also modify the power spectrum at lower values of ℓ . Indeed a study at lower values is especially interesting because HBG have identified a potentially significant asymmetric signal for the values $\ell \sim 2 - 40$, with the hemisphere oriented in

¹These studies were performed with MINUIT.

the directions (80° , 57°) having about 20% more power than in the opposite hemisphere. Subsequent work by Hansen et.al. (HBBG)[3] has explored the possibility that this signal could be due to correlated changes in the combination of the optical depth and the initial amplitude of fluctuations, or alternatively in the combination of the amplitude and tilt of the initial fluctuations. In this section, we explore how the constraints from the acoustic peak impact on possible explanations of the asymmetry found by HBG.

Of the cosmological parameters that we study, the ones that influence the power spectrum at lower values of ℓ are the overall amplitude, the spectral tilt and the cosmological constant. The latter is by far the easiest to rule out as the source of the HBG asymmetry. A variation in the cosmological constant would lead to changes at low values of ℓ through the late time integrated Sachs-Wolfe effect and also lead to changes in the acoustic peak². However, in order to produce a 20% change in the power spectrum at low ℓ , one would require over 100% variation of the cosmological constant. Our constraints given above firmly rule this out.

The simplest variation that would explain the HBG signal is that of the overall amplitude, requiring a 20% larger amplitude in one hemisphere than the other. Since the whole power spectrum is proportional to this amplitude, the same effect would show up at the acoustic peak. This is not seen, as is clear from our overall constraint of the previous section. Moreover, when we study the amplitude variation centered on the HBG direction, we find an even tighter constraint, with a best fit amplitude of $\eta = 0.02 \pm 0.02$ and a 2σ bound of $\eta \leq 0.06$.

Finally a variation in the tilt of the CMB spectrum would also influence the relative height of both the low and high ℓ regions. Here we are guided by the final result of HBBG in the section where they consider the possibility of correlated variations of both the tilt and the amplitude. Their central values are consistent with the same amplitude but a tilt that is different in the two hemispheres by 8%, a $2\text{-}3\sigma$ effect. We test this hypothesis by considering the properties of the acoustic peak under such variation. The spectral tilt has the property that it influences the power spectrum more strongly at lower values of ℓ than it does at the acoustic peak. Our studies with CMBFAST reveal that a $20 \pm 8\%$ variation in the power at low values can be fit with a variation of the spectral index of 0.07 ± 0.03 , and that this would be expected to produce a variation in the height of the acoustic peak of $7 \pm 3\%$, with a corresponding change in the peak location. We

²The cosmological constant does not influence the causal physics up to the time of last scatter, but it influences the angular scale of the acoustic peak.

can search for such a variation, finding a difference of $3 \pm 2\%$, i.e. a small but not statistically compelling signal. However, within the error bars, this observation is compatible with the expected signal.

We also can consider the combined variation of both the amplitude and the spectral tilt, using the information at lower values of ℓ as well as information at the peak. Performing a simultaneous fit to both regions, we find consistency for

$$\begin{aligned} n_N - n_S &= 0.083 \pm 0.041 \\ \frac{A_N}{A_S} &= 1.05 \pm 0.025 \end{aligned} \tag{7}$$

This result is consistent with that of HBBG within the error bars.

5 A hint of non-gaussianity

We have found no significant signal of a correlated variation of the height and location of the acoustic peak. However the data does appear to contain a more unusual asymmetry, which we discuss in this section. The northern and southern galactic hemispheres appear to have a different variance in the spread of the heights and locations of the peak, with the northern hemisphere having more variation. This is visible to the eye in Fig. 2. To probe this effect we define a variable

$$\delta_i = \sqrt{\left(\frac{\ell_i - \ell_0}{\sigma_\ell}\right)^2 + \left(\frac{h_i - h_0}{\sigma_h}\right)^2} \tag{8}$$

The distribution of the values of δ for the northern and southern hemispheres is shown in Fig. 3. It is easy to see visually that there is a significant difference between the hemispheres, with the northern hemisphere appearing to be anomalous. We refer to this as a non-gaussian feature since it would imply that there is not a uniform gaussian distribution in the properties of the peak.

We search for the direction that has the greatest non-gaussianity by identifying the direction that maximizes the asymmetry

$$f(\theta, \phi) = \sum_i \delta_i \cos \gamma_i \tag{9}$$

This search identifies $\theta \sim \pi$ as the direction that maximizes the difference, i.e. there is an asymmetry between the northern and southern galactic hemispheres. The net difference corresponds to $f(\theta = \pi, \phi) = -7.5$.

Figure 2: The data points on the acoustic peak are here separated into those of the northern (open circles) and southern (filled circles) galactic hemispheres. One can see visually that the data for the northern hemisphere is more widely scattered than that of the southern hemisphere.

We study the significance of this signal by simulating 256 data sets chosen randomly with a gaussian weight and the standard deviations given by HBG. In each case we search for the direction that maximizes the asymmetry $f(\theta, \phi)$ and find how large the asymmetry is when centered on this direction. The simulations reveal a median of $\bar{f}(\theta, \phi) = -3.3$ and a standard deviation of $\sigma_f = 1.3$. Consistently, we find that only one of the 256 simulated data sets has an asymmetry as large as is seen in the real data. We see that our signal is unlikely to be purely a statistical fluctuation, with an exclusion at the 3σ level.

If this signal is real physics, it appears as an extra random ingredient in the temperature fluctuations - dominantly in the northern hemisphere. Since it involves regions that were out of causal contact at the time of last scatter, it would either correspond to local foreground effects in the present universe or to physics very early in the period of inflation when these regions were causally connected. While the signal that we have identified appears to be statistically significant, it is not clear how it is related to other measures of non-gaussian behavior. The use of other measures has produced ample evidence for non-gaussianity in the WMAP data[7, 8, 9, 10, 11, 12, 22, 23,

Figure 3: The distribution of the parameter δ in the northern (light) and southern hemispheres (dark). The distribution for the southern hemisphere is consistent with the expected gaussian distribution within statistics. As described in the text, the distribution for the northern hemisphere appears more dispersed than expected.

24, 25, 26, 27, 28, 29]. Moreover, as mentioned in the introduction, for several of these signals there seems to be a correlation with the northern and southern hemispheres[7, 8, 9, 10, 11, 12]. For example, using a measure of the frequency of hills, lakes or saddles in the temperature distribution, [12] found that the northern hemisphere is non-gaussian at the 2-3 sigma level, while no significant non-gaussianity was found in the southern hemisphere. Our signal shares this directionality, even including the possibility that it is the northern hemisphere that is anomalous.

6 Conclusion

We have studied the properties of the acoustic peak in the CMB power spectrum and have provided measures of isotropy for the types of correlated variations of the height and location of the peak. Within the statistical uncertainty, no significant asymmetric signal was found.

The properties of the acoustic peak also constrain the possible explanations of the tentative signal of an asymmetry at low ℓ found by HBG. If

one attempts to explain this by the variation of a single parameter from the set considered in this paper, only the variation of the spectral tilt is able to accommodate the constraints of the acoustic peak. This is in agreement with the overall fit performed by HBBG[3].

We have also identified a curious difference between the data in the northern and southern hemispheres, with the northern hemisphere being more widely scattered than expected from the cosmic variance.

Acknowledgements

We would like to thank F. Hansen for helpful comments and for providing the data listed in Table 2 and K. Górski for useful communication. We also thank K. Dutta and A. Ross for comments and advice during the course of this work. This work was supported in part by the U.S. National Science Foundation.

Acknowledgement

References

- [1] M. S. Turner, “A Tilted universe (and other remnants of the preinflationary universe),” *Phys. Rev. D* **44**, 3737 (1991).
- [2] F. K. Hansen, A. J. Banday and K. M. Górski, “Testing the cosmological principle of isotropy: local power spectrum estimates of the WMAP data,” *arXiv:astro-ph/0404206*.
- [3] F. K. Hansen, A. Balbi, A. J. Banday and K. M. Górski, “Cosmological parameters and the WMAP data revisited,” *arXiv:astro-ph/0406232*.
- [4] F. K. Hansen, “Testing the isotropy of the Cosmic Microwave Background,” *arXiv:astro-ph/0406393*.
- [5] D. N. Spergel *et al.* [WMAP Collaboration], “First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Determination of Cosmological Parameters,” *Astrophys. J. Suppl.* **148**, 175 (2003) [*arXiv:astro-ph/0302209*].
C. L. Bennett *et al.*, “First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Preliminary Maps and Basic Results,” *Astrophys. J. Suppl.* **148**, 1 (2003) [*arXiv:astro-ph/0302207*].

- [6] D. J. Schwarz, G. D. Starkman, D. Huterer and C. J. Copi, “Is the low- l microwave background cosmic?,” arXiv:astro-ph/0403353.
- [7] H. K. Eriksen, F. K. Hansen, A. J. Banday, K. M. Górski and P. B. Lilje, “Asymmetries in the CMB anisotropy field,” *Astrophys. J.* **605**, 14 (2004) [arXiv:astro-ph/0307507].
- [8] C. G. Park, “Non-Gaussian Signatures in the Temperature Fluctuation Observed by the WMAP,” *Mon. Not. Roy. Astron. Soc.* **349**, 313 (2004) [arXiv:astro-ph/0307469].
- [9] H. K. Eriksen, D. I. Novikov, P. B. Lilje, A. J. Banday and K. M. Górski, “Testing for non-Gaussianity in the WMAP data: Minkowski functionals and the length of the skeleton,” *Astrophys. J.* **612**, 64 (2004) [arXiv:astro-ph/0401276].
- [10] H. K. Eriksen, A. J. Banday, K. M. Górski and P. B. Lilje, “The N-point correlation functions of the first-year Wilkinson Microwave Anisotropy Probe sky maps,” arXiv:astro-ph/0407271.
- [11] K. Land and J. Magueijo, “Cubic anomalies in WMAP,” arXiv:astro-ph/0405519.
- [12] F. K. Hansen, P. Cabella, D. Marinucci and N. Vittorio, “Asymmetries in the local curvature of the WMAP data,” *Astrophys. J.* **607**, L67 (2004) [arXiv:astro-ph/0402396].
- [13] U. Seljak and M. Zaldarriaga, “A Line of Sight Approach to Cosmic Microwave Background Anisotropies,” *Astrophys. J.* **469**, 437 (1996) [arXiv:astro-ph/9603033] and <http://www.cmbfast.org/>.
- [14] P. G. Castro, M. Douspis and P. G. Ferreira, “Scale of Homogeneity of the Universe from WMAP,” *Phys. Rev. D* **68**, 127301 (2003) [arXiv:astro-ph/0309320].
- [15] J. F. Donoghue, “Spatial gradients in the cosmological constant,” *JHEP* **0303**, 052 (2003) [arXiv:hep-ph/0101130].
- [16] T. Clifton, D. F. Mota and J. D. Barrow, “Inhomogeneous gravity,” arXiv:gr-qc/0406001.
- [17] V. Boucher, J. M. Gerard, P. Vandergheynst and Y. Wiaux, “Primordial constraint on the spatial dependence of the Newton constant,” arXiv:astro-ph/0407508.

- [18] S. M. Carroll and E. A. Lim, “Lorentz-violating vector fields slow the universe down,” arXiv:hep-th/0407149.
- [19] N. Kaloper and M. Kaplinghat, “Primeval corrections to the CMB anisotropies,” Phys. Rev. D **68**, 123522 (2003) [arXiv:hep-th/0307016].
- [20] H. Kurki-Suonio, P. Laguna and R. A. Matzner, “Inhomogeneous inflation: Numerical evolution,” Phys. Rev. D **48**, 3611 (1993) [arXiv:astro-ph/9306009].
- [21] D. Langlois and T. Piran, “Dipole anisotropy from an entropy gradient,” Phys. Rev. D **53**, 2908 (1996) [arXiv:astro-ph/9507094].
- [22] K. Land and J. Magueijo, “Multipole invariants and non-Gaussianity,” arXiv:astro-ph/0407081.
- [23] D. L. Larson and B. D. Wandelt, “The Hot and Cold Spots in the WMAP Data are Not Hot and Cold Enough,” arXiv:astro-ph/0404037.
- [24] P. Vielva, E. Martinez-Gonzalez, R. B. Barreiro, J. L. Sanz and L. Cayon, “Detection of non-Gaussianity in the WMAP 1-year data using spherical wavelets,” arXiv:astro-ph/0310273.
- [25] M. Cruz, E. Martinez-Gonzalez, P. Vielva and L. Cayon, “Detection of a non-Gaussian Spot in WMAP,” arXiv:astro-ph/0405341.
- [26] C. J. Copi, D. Huterer and G. D. Starkman, “Multipole Vectors—a new representation of the CMB sky and evidence for statistical anisotropy or non-Gaussianity at $2 \leq \ell \leq 8$,” Phys. Rev. D **70**, 043515 (2004) [arXiv:astro-ph/0310511].
- [27] P. Mukherjee and Y. Wang, “Wavelets and WMAP non-Gaussianity,” arXiv:astro-ph/0402602.
- [28] J. D. McEwen, M. P. Hobson, A. N. Lasenby and D. J. Mortlock, “A 6 sigma detection of non-Gaussianity in the WMAP 1-year data using directional spherical wavelets,” arXiv:astro-ph/0406604.
- [29] P. Cabella, M. Liguori, F. K. Hansen, D. Marinucci, S. Matarrese, L. Moscardini and N. Vittorio, “Primordial non-Gaussianity: local curvature method and statistical significance of constraints on $f_N L$ from WMAP data,” arXiv:astro-ph/0406026.

co – latitude	longitude	ℓ location	amplitude
0	0	223.3	5551
35	0	217.0	5235
35	60	215.8	5515
35	120	218.1	5909
35	180	207.5	6211
35	240	231.8	5231
35	300	232.7	5758
65	0	212.3	5957
65	36	217.9	4640
65	72	230.9	5410
65	108	224.5	5099
65	144	223.5	5506
65	180	221.5	5888
65	216	213.1	5506
65	252	217.9	5986
65	288	222.3	5927
65	324	224.6	5335
115	0	219.5	5633
115	36	222.3	5415
115	72	221.3	5510
115	108	219.5	5615
115	144	224.8	5804
115	180	216.3	5693
115	216	230.5	5627
115	252	220.8	5113
115	288	225.9	5395
115	324	219.4	5759
145	0	218.8	5315
145	60	219.3	5683
145	120	221.9	5792
145	180	216.6	5302
145	240	217.1	5362
145	300	221.4	5636
180	0	218.7	5513

Table 2: HBG data on the height and location of the acoustic peak on patches centered on different locations on the sky. We thank the authors of [2] for permission to include this data.

parameter	Baryon	CDM	Λ	amplitude	tilt
η fit	0.076 ± 0.048	0.004 ± 0.11	0.055 ± 0.035	0.027 ± 0.017	0.03 ± 0.02
2σ bound	0.17	0.28	0.125	0.061	0.057

Table 3: Isotropy bounds when interpreted as variations of the underlying parameters. The first row of results gives the fit value of the fractional variation of each parameter, while the second row reports the 2σ upper limit on the possible variation.