

3-2014

## 2014 Chart Book: Disease Management

Frank L. Caruso

*University of Massachusetts Amherst Cranberry Station, fcaruso@umass.edu*

Martha M. Sylvia

*University of Massachusetts Amherst Cranberry Station, martys@umass.edu*

Follow this and additional works at: <https://scholarworks.umass.edu/cranchart>



Part of the [Agriculture Commons](#), and the [Plant Sciences Commons](#)

---

Caruso, Frank L. and Sylvia, Martha M., "2014 Chart Book: Disease Management" (2014). *Cranberry Chart Book - Management Guide*. 189.

Retrieved from <https://scholarworks.umass.edu/cranchart/189>

This Public Service and Outreach is brought to you for free and open access by the Cranberry Station Outreach and Public Service Activities at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Cranberry Chart Book - Management Guide by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact [scholarworks@library.umass.edu](mailto:scholarworks@library.umass.edu).

**DISEASE MANAGEMENT 2014**

Prepared by Frank L. Caruso and Martha M. Sylvia

**UPRIGHT DIEBACK**

Cranberry plants affected by this disease typically have individual uprights that die back from the growing point toward the runner. Every upright may be infected on some runners, while other runners may only have one or a few infected uprights. In severe cases, the entire runner will be stressed or dying. Scattered uprights may be infected in the bed or whole patches of dieback may show up, particularly in younger beds. There are three phases during the season when symptoms appear: one shortly after the winter flood has been withdrawn, another in June and early July, and another phase in late August and September. Damage caused by this disease appears to be worse during growing seasons that have prolonged periods of drought or heat stress. This weakens the vines making them more susceptible to infection by fungal pathogens.

Most damage is probably caused by three different fungi. *Phomopsis vaccinii* (the fungus that causes viscid rot in fruit) is routinely isolated from symptomatic uprights and has been proven to cause symptoms by artificial inoculations. *Fusicoccum putrefaciens* (often) and *Synchronoblastia crypta* (rarely) are also isolated from infected uprights and have been shown to cause symptoms in artificial inoculations. Other fungi that may possibly infect uprights are *Aureobasidium*, *Colletotrichum*, and *Pestalotia*. Infection by these fungi in the field probably occurs at or shortly after bud break when the tissue is particularly susceptible. Infection may also occur during the entire growing season. Symptoms do not appear, however, until weather-related stresses weaken the plants. At this point, the tips are killed at the growing point and the symptoms progress downward on the upright.

The disease can be partially controlled through avoidance of stress on the plants through the hottest (and potentially the driest) portion of the growing season. Early-season fungicide applications at bud break and/or early bud expansion have given excellent control of the disease. Spores of the primary causal agent *Phomopsis* begin to be produced from overwintering cranberry tissue in April and May and the emerging buds are particularly susceptible to the infection. Fungicides targeted for fruit rot control also provide a degree of protection against this disease during early and mid-season infection periods.

<b>DISEASE/ TIMING</b>	<b>PESTICIDE/ FORMULATION</b>	<b>RATE (amt/A)</b>	<b>COMMENTS/RESTRICTIONS</b>
APRIL 25 THROUGH MAY 15	Champ DP Dry Prill Champ Formula 2 Flowable Champ WG	5.3 lb 5.33 pt 4.2 lb	Must be applied pre-bloom. 48-hour restricted entry interval.
<b><u>CHLOROTHALONIL FORMULATIONS</u></b>			
	Bravo Ultrex	3.8 – 6 lb	12-hour restricted entry interval. One pre-bloom application should be applied after the terminal bud has broken dormancy and begun to swell or has begun new growth. Exact timing will depend on whether the variety is early or late-season.
	Bravo Weather Stik	4 – 6.5 pt	
	Chlorothalonil 720 SC	4 – 6.5 pt	
	Chloronil 720	4 – 6.5 pt	
	Echo 720	4 - 7 pt	
	Echo 90DF	3.25 - 5.75 lb	
	Equus DF	3.8 – 6 lb	
	Equus 500 ZN	5.75 – 9.25 pt	
	Equus 720 SST	4 – 6.5 pt	
	Initiate ZN	5.75 – 9.25 pt	

For all above chlorothalonil formulations: Hold water for 3 days after application. When chlorothalonil formulations are to be used in a bed subject to Zone II regulations, growers must follow the required process to determine if these products may be used. See Zone II Section. The maximum allowable number of chlorothalonil applications is 3. If a chlorothalonil application is used for upright dieback control, only 2 fruit rot applications are allowed.

## 2 Diseases

### PHYTOPHTHORA ROOT ROT

Control of the disease can be achieved only through several integrated strategies. It is essential that drainage be improved in low areas of the bed. Tile, stones or other materials can be utilized, and new ditches can be dug. Existing ditches should be maintained to the proper depth as well. Areas of dieback should receive a uniform addition of sand to get the areas up to grade with the remainder of the bed. Stressed plants on the margin of dieback areas should be given an extra dose of fertilizer to stimulate root growth. Use of a soil fumigant should be employed on renovations. After the drainage has been improved, fungicides should be applied several times per season until the vines have completely filled in the bare spots. Once this has been achieved, only a single spring application should be necessary.

Precautions should be taken to avoid spread of the pathogen from infested beds to uninfested beds. Machinery, equipment, footwear, etc., should be sterilized using steam, bleach (10% Clorox solution), or alcohol. If possible, the sequence of flooding the beds during water harvest should be adjusted to flood heavily infected beds last. When vines are purchased from other growers, the grower should be certain that they have not come from infected beds. Though very few roots are present in such vine deliveries, it only takes a few infected roots to initiate infection in a new location.

**Three fungicide applications** per season are recommended for newly diagnosed instances. As areas of dieback recover, consult with the Extension Plant Pathologist regarding the fungicide schedule. The first application should go on between April 25 and May 15<sup>th</sup>. The second application should occur 60-90 days after the first but 45 days before harvest if using mefenoxam (e.g., Ridomil). A third application can be done after harvest, preferably prior to November 15.

The drainage should be improved BEFORE applying any fungicide to the affected bed.

DISEASE/ TIMING	PESTICIDE/ FORMULATION	RATE (amt/A)	COMMENTS/RESTRICTIONS
APRIL 25 THROUGH MAY 15 (1 <sup>st</sup> app)	Aliette WDG	5 lb	Rates are for chemigation application!  Do not tank mix! Read label of product used!
<b>Foliar applications</b>	Fosphite	2-4 qt	
	Fungi-Phite	1-2 qt	
	Helena ProPhyt	4 pt	
	K-Phite 7LP AG	2-4 qt	
	Phostrol	5-6 pt	
	Rampart	2-4 qt	
<b>Soil applications</b>	Metastar 2EC	4-7 pt	Ground or chemigation only. No air!
	Ridomil Gold SL	1-1.75 pt	
	Ultra Flourish	2-3.5 pt	
Water fungicide in!	Ridomil Gold GR	20-35 lb	Ground or air only. No Chemigation!
*Any formulation can be used for spot treatment if the affected area(s) are small.			

**Soil applications** must be watered in after application. Run the sprinklers for 3 hours after application to water the fungicide into the root zone. Too much water, however, may push the chemical past the root zone. Therefore, do not apply if more than 0.5 inch of rainfall is forecast or if the sprinklers will need to be run for more than 5 hours during the first few days after application.

## FRUIT ROT

This is the most prevalent disease problem that cranberry growers face from season to season. Cranberry fruit can be infected by a large number of different fungal pathogens.

**Fruit rot, various pathogens include:**

*Allantophomopsis lycopodina* and *A. cystisporae* (black rot)

*Botryosphaeria vaccinii* (Botryosphaeria fruit rot)

*Coleophoma empetri* (ripe rot)

*Glomerella cingulata* (bitter rot)

*Fusicoccum putrefaciens* (end rot)

*Penicillium* sp. (Penicillium rot)

*Phomopsis vaccinii* (viscid rot)

*Phyllosticta vaccinii* (early rot or bull's eye rot)

*Phyalospora vaccinii* (blotch rot)

The degree of fruit rot that occurs in different beds during different growing seasons is dependent on many factors. Weather is probably the most important variable that affects the amount of rot that occurs. Temperature, humidity, rainfall, as well as special events such as hailstorms, are important on a regional basis. Local microclimatic variations can also lead to particular sections of a bog that are more prone to the development of rot. The density of vine growth and drainage are important because they have a direct bearing on the length of time vines remain wet from rain, fog, dew or irrigation. Excessive vine growth and poor drainage prevent rapid drying and favor the infection by rot fungi. Practices that promote rank (excessive) vine growth, such as excessive fertilizer, frequent late water, too frequent sanding or too heavy sand on muck bottom, holding water high in ditches, and too frequent irrigation should be avoided to minimize conditions that are favorable for fruit rot development.

Spores of the fungi are dispersed from overwintering sources by wind or wind-driven rain. These spores land on the blossom or small developing fruit and, if there is a suitable layer of moisture present for 6-8 hours, the fungi will be allowed to penetrate into the plant tissue. If conditions are favorable and the cultivar is susceptible, development of the fungus may be rapid and berries showing symptoms typical of one of the rots may be apparent within a week. If conditions are unfavorable and if the cultivar is less susceptible, development of the infection will be slow. Many of the fruit rot fungal pathogens infect the berry but remain latent in the plant until the berry reaches a certain maturation or physiological stage, at which point they will further infect the tissue, resulting in visible symptoms. Symptoms may not be visible until later in the growing season, or they may not be visible until after the berries have been harvested and held in storage.

Fungicides are an important strategy for the control of fruit rot. Most berry infection occurs during the bloom period. Applications typically begin during early bloom (mid-June). Once the fruit has set and begun to increase in size (mid-late July), further fungicides are no longer necessary. The choice of fungicides, the rate of the fungicide, and the time interval between applications are dependent on the individual bed and its past incidence of fruit rot.

Fungicide decisions should be based on the Keeping Quality Forecast (KQF) found on the Station's website (<http://www.umass.edu/cranberry/cropinfo/keepqf.html>). If the forecast is good to excellent, consider fewer applications and/or lower rates of the fungicides. History of the cranberry bed is also important: if a bed is prone to fruit rot, one will need to be more conservative in the decision to reduce the total amount of active ingredient. Four fungicide applications are usually necessary for a bed prone to fruit rot. One or two fungicide applications may be adequate for a bed with very little fruit rot in previous growing seasons. Fungicide applications are more important when the berries will be harvested for fresh fruit, as this fruit will be held in storage for extended periods. Storage rot is not a concern for berries that are water harvested, as these berries will immediately be frozen in most cases. Normally, 3-5% fruit rot at delivery is considered acceptable. If in doubt, call the Extension Plant Pathologist.

## 4 Diseases

DISEASE/ TIMING	PESTICIDE/ FORMULATION	RATE (amt/A)	COMMENTS/RESTRICTIONS
--------------------	---------------------------	-----------------	-----------------------

### FRUIT ROT

EARLY BLOOM (10-20%), THEN AT 10-14 DAY INTERVALS

#### NEWER CHEMISTRIES

Abound	6.0-15.5 fl oz	Although 6 applications are allowed, no more than 2 applications should be used. See Resistance Management section page iii. Use this fungicide for the earlier applications spaced 7-10 days. Hold water for 14 days.
Indar 2F	6-12 fl oz	Do not use more than 2 apps. Do not use prior to bloom. 7-14 days between apps. 30 day PHI.
Proline 480SC	5 fl oz	No more than 2 apps. 45 day PHI.

For Abound Indar, and Proline: Resistance development to Abound, Indar, and Proline by the fruit rot fungi is a very real and serious threat. Applications of the fungicide should be made pre-infection rather than post-infection to minimize resistance development. One application of each fungicide is recommended per growing season unless the Keeping Quality Forecast (KQF) predicts poor or worse keeping quality. Use Abound very carefully and avoid drift if the bed is next to a McIntosh apple orchard, as the fungicide is highly phytotoxic to this cultivar.

#### CHLOROTHALONIL FORMULATIONS

Bravo Ultrex	3.8-6.0 lb	Use the maximum rate in beds with high rot incidence on a 10-day schedule. Zone II restricted, 3 applications/season, 12 hr REI, hold water for 3 days.
Bravo Weather Stik	4-6.5 pt	
Chloronil 720, Initiate 720	4-6.5 pt	
Chlorothalonil 720 SC	4-6.5 pt	
Echo 720	4-7 pt	
Echo 90DF	3.25-5.75 lb	
Equus DF	3.8-6.0 lb	
Equus 500 ZN	5.75-9.25 pt	
Equus 720 SST	4-6.5 pt	
Initiate ZN	5.75-9.25 pt	

For all chlorothalonil formulations: When chlorothalonil formulations are to be used in a bed subject to Zone II regulations, growers must follow the required process to determine if these products may be used. See Zone II Section. The maximum allowable number of chlorothalonil applications is 3. If 1 Bravo application was used for upright dieback control, only 2 fruit rot applications are allowed. Do not mix with Dipel. 12-hour restricted entry, note 6.5 day eye irritant! Do not release irrigation water for at least 3 days following application. Existing product may be used according to the label.

Ferbam Granuflo	6 lb	Do not apply more than 5 times. Apply at 14-day intervals. Using rates below recommended rate will be ineffective. 50-day PHI, 24-hr REI.
-----------------	------	---

DISEASE/ TIMING	PESTICIDE/ FORMULATION	RATE (amt/A)	COMMENTS/RESTRICTIONS
--------------------	---------------------------	-----------------	-----------------------

**FRUIT ROT (CONTINUED)**

EARLY BLOOM (10-20%), THEN AT 7-10 DAY INTERVALS

MANCOZEBS

Dithane F-45 Rainshield	2.4-4.8 qt	Addition of spray adjuvants will improve distribution and deposition for all of the mancozeb compounds.  May delay color development!
Dithane M-45	3-6 lb	
Penncozeb 75DF	3-6 lb	
Penncozeb 80WP	3-6 lb	
Koverall	3-6 lb	
Roper DF Rainshield	3-6 lb	

MID-BLOOM,  
THEN AT 7-10 DAY INTERVALSCOPPERS

Cuprofix Ultra 40 Disperss	5 lb
Top Cop with Sulfur	2 qt

MANCOZEBS

ManKocide	7 lb	Addition of spray adjuvants will improve distribution and deposition for all of the mancozeb compounds.  May delay color development
Manzate Flowable	2.4-4.8 qt	
Manzate Max	2.4-4.8 qt	
Manzate Pro-Stick	3-6 lb	

LATE BLOOM,  
THEN 1 OR 2 APPLICATIONS AT 10-14 DAY INTERVALSCOPPERS

Badge X2	3.5 lb	Note: Do not mix copper fungicides with insecticides. Do not tank mix with Aliette or phosphites unless the spray solution has been buffered first.
Badge SC	7 pt	
Champ DP Dry Prill	5.6 lb	
Champ Formula 2 flowable	5.33 pt	
Champ WG	4.2 lb	
Copper-Count-N	4-8 qt	
Kentan DF	4-5.25 lb	
Kocide 2000	6 lb	
Kocide 3000	3.5 lb	
Master Cop	3 pt	
Nordox	8 lb	
Nordox 75 WG	5 lb	
Nu-Cop HB	4.2 lb	
Nu-Cop 3L	4.33-5.67 pt	
Nu-Cop 50 DF	4.2 lb	
3lb copper flowable	10.67 pt	

## 6 Diseases

### FRUIT ROTS - CULTURAL CONSIDERATIONS

**Late Water:** Holding late water (flooding from mid-April to mid-May) will improve berry quality by disrupting the life cycles of rot-inducing fungi. In late water years, fungicide rates and/or the number of total applications can be reduced with no sacrifice in fruit quality. The fungicide program should not be eliminated completely or vine diseases may be a problem the following growing season. The number of fungicide applications can be reduced and the lowered rates can also be used during the first year after late water. Fungal inoculum will begin to build up during the second year after late water.

**Lush Vines:** Where fertilizer applications have been heavy, vines will tend to become very overgrown. This will lead to poor air circulation, retention of high humidity, and slow drying-out of heavy dew. These conditions encourage infection by the fungi that cause fruit rot and red leaf spot. When growth is excessive, pruning is recommended to promote air circulation in the vine canopy.

**Trash Removal:** Cranberry leaves, stems, and fruits left behind after harvest are colonized by several fungi that cause field and storage rot. This trash can serve as an inoculum source for fungal infections of the uprights, blossoms, or fruits in subsequent growing seasons. If the bed was dry-harvested, trash should be removed from the bed with a post-harvest flood in the fall or from the winter flood before it is withdrawn in February or March. Remove trash from water-harvested beds during harvest or as soon after as possible. Trash piles should not be left next to the bed. Trash should be deposited at least a quarter mile from the bed if possible. Self-pollinated seeds in berries left behind on the bog may germinate in the soil and possibly produce plants that are the typical "mongrels". These genotypes may produce much vegetation but few berries, and in worst case scenarios, may take over the productive vines in the bed.

**Irrigation:** When irrigation is necessary, sprinkler systems should be run for up to 4-5 hours in the early morning, and not in the early evening. Vines can get watered with minimal evaporation, and the surface of the vines can dry out in the sun's heat. When watering is done in the early evening, the vines are kept wet for an extended time period, thus creating favorable conditions for infection by the fruit rot fungi. On days with excessive temperatures (>100°F on the bed), particularly in newly planted or recently sanded beds, sprinklers should be run for 1-2 hours in the late morning or early afternoon to cool the vines and berries and to prevent injury. Sprinklers should be run to prevent scalding of the fruit when all of the following conditions persist: (1) dewpoints of 55°F or less during midday and afternoon hours, (2) high temperatures of 80°F or more, (3) clear or scattered sky conditions during the day, (4) bed soil moisture is low, (5) wind speeds average greater than 11 mph, and (6) no rainfall has occurred during the last 48 hours. This "forecast" is based on research performed in New Jersey. Scalded berries are typically browned on one side, with a clear demarcation between the brown area and the green (usually) area of the fruit. The rotted area in a berry affected with fruit rot typically has an area of anthocyanin production (reddish border) adjacent to the affected area. After seven days, a scalded berry will be hard to discern from a totally rotted berry, particularly since fungi will colonize the stressed scalded berry.

**Resistant Varieties:** When replanting bogs or planting new bogs, certain varieties with proven field rot resistance should be considered: Black Veil, Foxboro Howes, Matthews, Shaw's Success, and Wilcox. We are waiting to see how Crimson Queen, Demoranville, Grygleski, HyRed, Mullica Queen and Scarlet Knight rate as far as fruit rot susceptibility goes.

Research is still needed on the consistency of these cultivars to produce good crops. Small plantings are encouraged initially. Howes and Stevens also have good fruit rot resistance.

One or two fungicide applications during the first two years after planting will help reduce fungal inoculum and may reduce fruit rot in subsequent years.

---

<b>DISEASE/ TIMING</b>	<b>PESTICIDE/ FORMULATION</b>	<b>RATE (amt/A)</b>	<b>COMMENTS/RESTRICTIONS</b>
<b>FAIRY RING</b>			
This disease is sporadic in occurrence and the severity of symptoms varies from year to year. It can be spread from one bed to another through uprooted vines during wet or dry harvest and their subsequent dislodgment in the next harvested bed. Picking machines should be freed of vines before moving to the next bed. Damage is usually worst during periods of drought; keep vines well-irrigated. Applications of lime during the growing season give limited benefits to the vines and have no effect on the fungal pathogen(s).			
MAY	Abound Indar 2F	15.5 fl oz/30-100 gal 12 fl oz/30-100 gal	Make first application at bud break. Measure ring diameter and add 10 feet to the diameter. Irrigate for 1-2 hours before and following application. Repeat 2-4 weeks later if necessary.
JUNE - JULY	Ferbam Granuflo	9 lb/100 gal	Apply 1 gal of this mixture to 1 sq ft area. Treat the area 3 feet beyond the advancing line of dying vines and 2 feet within the line. Do not apply after July 31. Only 1 application!
MID-AUGUST THROUGH OCTOBER	Sul-Po-Mag or K-Mag 0-0-22	4000 lb/A or 1.5 oz/sq. ft.	Use alone or after lime to help vines recover. Follow-up applications may be necessary. This may help vines out-compete the fungus.

## 8 Diseases

### DISEASE MANAGEMENT NOTES

1. Read and follow label instructions. Always check label for variations in restricted entry interval and worker protection standards.
2. Make all pesticide applications in a manner to prevent contamination of streams, ponds and public ways. Impound water (as per label) for as long as possible after applying.
3. RESISTANCE DEVELOPMENT to Abound, Indar, and Proline by the fruit rot fungi is a very real and serious threat. Applications of the fungicide should be made pre-infection rather than post-infection to minimize resistance development. One application of each fungicide is recommended per growing season unless the KQF predicts poor or worse keeping quality. Use Abound very carefully and avoid drift if the bed is next to a McIntosh apple orchard, as the fungicide is highly phytotoxic to this cultivar. See Resistance Management section.
4. REFLOODING for black-headed fireworm control in June is likely to INCREASE FRUIT ROT and MAY SERIOUSLY REDUCE THE CROP.
5. Holding late water will most likely reduce the incidence of fruit rot.
6. PRE-MIX fungicides with a small amount of water until a smooth suspension is obtained before final dilution. Use immediately. Blossom injury may occur with concentrate sprays especially when sprayed by air when the temperature in the bog is above 85°F. Do not combine any copper fungicide with an insecticide. Do not tank mix copper compounds with Aliette or any of the phosphites for Phytophthora unless appropriate precautions have been taken to buffer the spray solution or severe phytotoxicity will result.
7. Consider delaying harvest to obtain acceptable color in thick vines or when Mancozeb is used.
8. SANDING and FERTILIZING. Frequent resanding and fertilizing helps reclaim beds infected with false blossom disease. These are accepted IPM practices. Regular uniform sanding most likely helps to reduce inoculum of the fungi that cause fruit rot. Sanding should not be done during the same year late water is to be held.
9. SPREADER STICKERS are contained in most fungicides. The additions of wetting agents or spreader stickers to Bravo, Echo, or Equus may cause phytotoxicity damage. Please check the fungicide label.
10. STORED PESTICIDES may deteriorate. Avoid freezing liquid formulations. It is not advisable to use old materials in opened containers. Follow Pesticide Bureau regulations for disposing of pesticides and their containers.
11. Even if half-rates are used, the maximum number of applications (not material applied) must not be exceeded. It is not recommended that any fungicide application be lower than the lowest recommended rate.
12. Review the Disease Management BMP in the UMass Best Management Practices Guide: (<http://www.umass.edu/cranberry/pubs/bmps.html>).
13. Organic Options. Some fungicides are certified organic for disease management in cranberry. They include many of the coppers, OxiDate (128 fl oz/100 gallons of water; apply 25-100 gal/solution/treated acre), and Serenade ASO, MAX, or Optimum; a biofungicide labeled for mummy berry, botrytis, and bacterial canker at 2-6 qts/A.