

Spring 3-2016

## 2016 Chart Book: Resistance Management

Martha Sylvia

*University of Massachusetts Amherst Cranberry Station, martys@umass.edu*

Follow this and additional works at: <https://scholarworks.umass.edu/cranchart>



Part of the [Agriculture Commons](#), and the [Plant Sciences Commons](#)

---

Sylvia, Martha, "2016 Chart Book: Resistance Management" (2016). *Cranberry Chart Book - Management Guide*. 217.  
Retrieved from <https://scholarworks.umass.edu/cranchart/217>

This Public Service and Outreach is brought to you for free and open access by the Cranberry Station Outreach and Public Service Activities at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Cranberry Chart Book - Management Guide by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact [scholarworks@library.umass.edu](mailto:scholarworks@library.umass.edu).

## 4 RESISTANCE MANAGEMENT

### RESISTANCE MANAGEMENT 2016

Prepared by Martha M. Sylvia

In an effort to manage resistance with our pesticides, many labels now come with a “group” number assigned to them. The group ID is specific among insecticides, herbicides and fungicides. Many of our cranberry pesticides are in their own group. The largest groups with the same ID are the organophosphates and neonicotinoids. The following 3 pages show the groupings for our cranberry pesticides. The goal in resistance management is to not repeatedly use compounds that fall within the same group. Resistance management may include alternating products with different modes of action or limiting the total number of applications per season.

#### Fungicide Resistance Action Committee (FRAC) (<http://www.frac.info/home>)

The group that advises for fungicide resistance is the Fungicide Resistance Action Committee (FRAC). Their goal is to prolong the effectiveness of fungicides that are likely to encounter resistance problems and to limit crop losses should resistance appear. For cranberry, Ridomil and Abound are fungicides that are at high risk for resistance development, while Indar and Proline are at medium risk. They should not be used repeatedly and should be carefully alternated with other fungicides from other groupings. See grouping of cranberry fungicides on page 6. A box like this would appear on the front of the label:

GROUP      **11**      FUNGICIDE

#### Herbicide Resistance Action Committee (HRAC) (<http://www.hracglobal.com/pages/Home.aspx>)

The Herbicide Resistance Action Committee developed a classification of herbicides according to their mode of action. A similar system to FRAC has been developed by the Weed Science Society of America (WSSA) using numbers instead of letters to designate the categories. This classification is found on a few herbicide labels, for example Callisto labels have this marking:

GROUP      **27**      HERBICIDE

Herbicide resistance is a world wide phenomenon with more than 215 documented cases. Selection of herbicide-resistant weed populations is often the result of the continuous use of the same herbicide or herbicides with the same mode of action. A key step in resistance management is to minimize the continuous use of herbicides with the same mode of action through rotations and combinations of products. One of the purposes of these classification systems is to make it easier for farmers and farm advisors to understand which herbicides share the same mode of action without having to actually know the biochemical basis.

In cranberry, our biggest concern is our new reliance on Callisto. Be sure to rotate other compounds into your herbicide schedule. Do not treat the same bog with Callisto year after year. See table of cranberry herbicides by grouping on page 7.

#### Insecticide Resistance Action Committee (IRAC) (<http://www.iraconline.org/>)

An Insecticide Resistance Action Committee (IRAC) has been formed to assemble the information for insecticides. Their goal is to manage resistance to keep agriculture sustainable. For cranberry, organophosphates and neonicotinoids have the most compounds within their group. We are reliant on several compounds in these groupings. As long as growers remember to alternate between groupings and not repeat same mode-of-action compounds over and over, we should be able to keep newer compounds viable for decades. See Cranberry Insecticides by grouping on the next page. Insecticides are grouped clearly by chemical makeup and most insecticide labels now included markings such as this:

GROUP      **5**      INSECTICIDE

## Insecticide Resistance Action Committee (IRAC) Grouping for cranberry insecticides

IRAC GROUP	TRADE NAME	ACTIVE INGREDIENT	MODE OF ACTION	CHEMICAL FAMILY
1	<b>Diazinon</b>	diazinon	Acetylcholine esterase inhibitor	Organophosphates and carbamates
	<b>Imidan</b>	phosmet		
	<b>Lorsban</b>	chlorpyrifos		
	<b>Orthene</b>	acephate		
	<b>Sevin</b>	carbaryl		
3	<b>Pyreth-It Pyganic</b>	pyrethrin	Sodium channel modulators	Pyrethrins
4, 4A	<b>Actara</b>	thiamethoxam	Nicotinic Acetylcholine receptor agonists	Neonicotinoids
	<b>Admire</b>	imidacloprid		
	<b>Assail</b>	acetamiprid		
	<b>Belay</b>	clothianidin		
	<b>Scorpion</b>	dinotefuran		
5	<b>Delegate</b>	spinetoram	Nicotinic Acetylcholine receptor allosteric activators	Spinosyns
	<b>Entrust</b>	spinosad		
11	<b>Dipel, Xentari Biobit</b>	<i>Bacillus thuringiensis</i>	Microbial disruptors of insect midgut membranes	<i>Bacillus thuringiensis</i>
15	<b>Rimon</b>	novaluron	Inhibitors of chitin biosynthesis	Benzoylureas
18	<b>Confirm</b>	tebufenozide	Ecdysone agonists / molting disruptors	Diacylhydrazines
	<b>Intrepid</b>	methoxyfenozide		
21	<b>Nexter</b>	pyridaben	Mitochondrial complex / electron transport inhibitor	Meti acaracides
22	<b>Avaunt</b>	indoxacarb	Voltage-dependent sodium channel blockers	Oxadiazines
23	<b>Oberon</b>	spiromesifen	Inhibitors of acetyl CoA carboxylase	Tetramic acid derivatives
28	<b>Altacor</b>	chlorantraniliprole	Ryanodine receptor modulators	Diamides

## 6 RESISTANCE MANAGEMENT

### Fungicide Resistance Action Committee (FRAC) Grouping for cranberry fungicides

FRAC GROUP	TRADE NAME	COMMON NAME	MODE OF ACTION	GROUP NAME	CHEMICAL GROUP	Resistance Development Risk
4	<b>Metastar</b>	mefenoxam	<b>A1:</b> RNA polymerase I	PA – fungicides (PhenylAmides)	acylalanines	<b>High Risk</b>
	<b>Ridomil</b>	metalaxyl				
	<b>Ultra</b>					
	<b>Flourish</b>					
11	<b>Abound</b>	azoxystrobin	<b>C3:</b> cytochrome bc1 at Qo site	QoI-fungicides	methoxy-acrylates	<b>High Risk</b>
	<b>Aftershock</b>	fluoxastrobin		Strobilurins	dihydro-dioxazines	
	<b>Evito</b>					
3	<b>Indar</b>	fenbuconazole	<b>G1:</b> c14-demethylase in sterol biosynthesis	DMI-fungicides (DeMethylation Inhibitors)	triazoles	<b>Medium Risk</b>
	<b>Proline</b>	prothioconazole				
19	<b>OSO</b>	polyoxin	<b>H4:</b> chitin synthase	polyoxins	peptidyl pyrimidine nucleoside	<b>Medium Risk</b>
	<b>Ph-D</b>					
33	<b>Aliette</b>	fosetyl-Al	Unknown	phosphonates	ethyl phosphonates	Low Risk
	<b>Legion</b>	aluminum-tris				
	<b>Fosphite</b>	phosphorous acids and salts				
	<b>Fungi-Phite</b>					
<b>K-Phite, Phostrol, ProPhyt, Rampart</b>						
M1	<b>Champ</b>	copper (salts)	<b>M1:</b> Multi-site contact activity	inorganic	inorganic	Low Risk
	<b>Kocide</b>					
M3	<b>Ferbam</b>	ferbam	<b>M3:</b> Multi-site contact activity	dithiocarbamates	dithiocarbamates	Low Risk
	<b>Manzate</b>	mancozebs		EBDC's (Ethylene bis dithio carbamate)		
	<b>Dithane</b>					
	<b>Penncozeb</b>					
M5	<b>Bravo</b>	chlorothalonil	<b>M5:</b> Multi-site contact activity	chloronitriles	chloronitriles	Low Risk
	<b>Chloronil</b>					
	<b>Echo, Equus</b>					
	<b>Initiate</b>					

Herbicide Resistance Action Committee (HRAC) Grouping for cranberry herbicides  
Group numbering from Weed Science Society of America (WSSA) at right

HRAC GROUP	TRADE NAME	ACTIVE INGREDIENT	MODE OF ACTION	CHEMICAL FAMILY	WSSA GROUP
<b>A</b>	<b>Fusilade</b>	fluazifop-P-butyl	Inhibition of acetyl CoA carboxylase (ACCase)	Aryloxyphenoxy-propionate 'FOPs'	<b>1</b>
	<b>Select</b>	clethodim		Cyclohexanedione	
	<b>Poast</b>	sethoxydim		'DIMs'	
<b>F1</b>	<b>Evital</b>	norflurazon	Bleaching: Inhibition of carotenoid biosynthesis at the phytoene desaturase step (PDS)	Pyridazinone	<b>12</b>
<b>F2</b>	<b>Callisto</b>	mesotrione	Bleaching: Inhibition of 4-hydroxyphenyl-pyruvate-dioxygenase (4-HPPD)	Triketone	<b>27</b>
<b>G</b>	<b>Roundup</b>	glyphosate	Inhibition of EPSP synthase	Glycine	<b>9</b>
<b>K3</b>	<b>Devrinol</b>	napropramide	Inhibition of VLCFAs (Inhibition of cell division)	Acetamide	<b>15</b>
<b>L</b>	<b>Casoron</b>	dichlobenil	Inhibition of cell wall (cellulose) synthesis	Nitrile	<b>20</b>
	<b>Quinstar</b>	quinclorac		Quinoline carboxylic acid	<b>26</b>
<b>O</b>	<b>Quinstar</b>	quinclorac	Action like indole acetic acid (synthetic auxins)	Quinoline carboxylic acid	<b>4</b>
	<b>2,4-D Weedar 64</b>	2,4-D		Phenoxy-carboxylic acid	
	<b>Stinger</b>	clopyralid		Pyridine carboxylic acid	