

2018

Genome Sequence of Verrucomicrobium sp. Strain GAS474, a Novel Bacterium Isolated from Soil

Grace Pold

University of Massachusetts Amherst

Erin M. Conlon

University of Massachusetts Amherst

Marcel Huntemann

U.S. Department of Energy Joint Genome Institute

Manoj Pillay

U.S. Department of Energy Joint Genome Institute

Natalia Mikhailova

U.S. Department of Energy Joint Genome Institute

See next page for additional authors

Follow this and additional works at: https://scholarworks.umass.edu/micro_faculty_pubs

 Part of the [Microbiology Commons](#)

Recommended Citation

Pold, Grace; Conlon, Erin M.; Huntemann, Marcel; Pillay, Manoj; Mikhailova, Natalia; Stamatis, Dimitrios; Reddy, T.B.K.; Daum, Chris; Shapiro, Nicole; Kyripides, Nikos; Woyke, Tanja; and DeAngelis, Kristen, "Genome Sequence of Verrucomicrobium sp. Strain GAS474, a Novel Bacterium Isolated from Soil" (2018). *Genome Announcements*. 314.
[10.1128/genomeA.01451-17](https://doi.org/10.1128/genomeA.01451-17)

This Article is brought to you for free and open access by the Microbiology at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Microbiology Department Faculty Publication Series by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

Authors

Grace Pold, Erin M. Conlon, Marcel Huntemann, Manoj Pillay, Natalia Mikhailova, Dimitrios Stamatis, T.B.K. Reddy, Chris Daum, Nicole Shapiro, Nikos Kyrpides, Tanja Woyke, and Kristen DeAngelis



Genome Sequence of *Verrucomicrobium* sp. Strain GAS474, a Novel Bacterium Isolated from Soil

Grace Pold,^a Erin M. Conlon,^b Marcel Huntemann,^c Manoj Pillay,^c Natalia Mikhailova,^c Dimitrios Stamatis,^c T. B. K. Reddy,^c Chris Daum,^c Nicole Shapiro,^c Nikos Kyrpides,^c Tanja Woyke,^c Kristen M. DeAngelis^d

^aGraduate Program in Organismic and Evolutionary Biology, University of Massachusetts–Amherst, Amherst, Massachusetts, USA

^bDepartment of Mathematics and Statistics, University of Massachusetts–Amherst, Amherst, Massachusetts, USA

^cU.S. Department of Energy Joint Genome Institute, Walnut Creek, California, USA

^dDepartment of Microbiology, University of Massachusetts–Amherst, Amherst, Massachusetts, USA

ABSTRACT *Verrucomicrobium* sp. strain GAS474 was isolated from the mineral soil of a temperate deciduous forest in central Massachusetts. Here, we present the complete genome sequence of this phylogenetically novel organism, which consists of a total of 3,763,444 bp on a single scaffold, with a 65.8% GC content and 3,273 predicted open reading frames.

Verrucomicrobium sp. strain GAS474 is a novel slow-growing Gram-negative bacterium isolated from the upper mineral soil of a long-term warming experiment (1). *Verrucomicrobia* represent 23% of the bacteria present in soil (2), but fewer than 50 ecotypes have been cultured and sequenced. Therefore, the characterization and sequencing of members of this group can improve our understanding of vital soil processes, such as carbon cycling.

Soil was collected from the top 10 cm of mineral soil at the Prospect Hill warming experiment at the Harvard Forest Long Term Ecological Research site (42.54°N, 72.18°W) and subsequently dispersed for 30 min in a phosphate buffer-dithiothreitol-sodium pyrophosphate solution. After settling, the solution was spread onto VL55 medium (3) solidified with gellan gum and amended with 0.05 g/liter xylan from birchwood. Plates were incubated aerobically at 22 to 25°C, and GAS474 appeared after 8 weeks.

Cells were prepared for sequencing by growing on an R2A plate incubated for 6 weeks under aerobic conditions at 22 to 25°C; genomic DNA (gDNA) was extracted using the standard Qiagen Genomic-tip protocol for bacteria. Whole-genome sequencing was completed at the Joint Genome Institute on the PacBio RS platform (4). The reads were assembled using the Hierarchical Genome Assembly Process (HGAP) version 2.3.0 (5), leading to a single contig 3.763 Mb in size with 178× coverage.

Gene prediction and functional annotation were performed using the U.S. Department of Energy Joint Genome Institute annotation pipeline (6). The genome contains 3,273 predicted coding sequences, of which, 73.0% were assigned a function and 9.4% were predicted to contain signal peptides using SignalP (7). The genome contains 47 tRNAs and a single rRNA operon. The genome is available for comparative analysis through the Integrated Microbial Genomes (IMG) data management system (8).

GAS474 is phylogenetically and phenotypically distinct from its closest cultured relatives, sharing less than 88% 16S rRNA identity and less than 73% average nucleotide identity with *Methylacidimicrobium fagopyrum*, *Methylacidimicrobium tartarophylax*, *Methylacidimicrobium cyclopophantes*, *Methylacidiphilum fumariolicum* SolV, and *Methylacidiphilum kamchatkense* Kam1. GAS474 lacks the particulate methane monooxygenase and RuBisCO genes that enable its closest relatives to survive as autotrophic aerobic

Received 20 November 2017 Accepted 27 November 2017 Published 25 January 2018

Citation Pold G, Conlon EM, Huntemann M, Pillay M, Mikhailova N, Stamatis D, Reddy TBK, Daum C, Shapiro N, Kyrpides N, Woyke T, DeAngelis KM. 2018. Genome sequence of *Verrucomicrobium* sp. strain GAS474, a novel bacterium isolated from soil. *Genome Announc* 6:e01451-17. <https://doi.org/10.1128/genomeA.01451-17>.

Copyright © 2018 Pold et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Kristen M. DeAngelis, deangelis@microbio.umass.edu.

methanotrophs (9, 10). Instead, its genome shows enrichment in genes for secondary metabolite processing (Clusters of Orthologous Groups [COG] category Q; 1.5× increase in fraction of protein-coding reads compared to four closest relatives), pilus formation (COG category W; 5.3× greater), signal transduction (COG category T; 2× greater), and motility (COG category N; 6× greater). GAS474 also shows an expanded ability to process diverse saccharides, having more than twice the fraction (8.5 versus 4.1%, determined using dbCAN [11]) and diversity (80 versus 38 distinct carbohydrate-active enzyme families, and 29 versus 12 glycoside hydrolase families) of genes and the potential to process a broader range of substrates than the primarily peptidoglycan- and α -glycosidic limited abilities (GH113 and GH57) of its methanotrophic relatives. We therefore introduce the genome of *Verrucomicrobium* sp. strain GAS474 as a representative of a novel cultivated lineage in the verrucomicrobial phylogenetic tree.

Accession number(s). This whole-genome shotgun project has been deposited in GenBank under the accession no. [LT629781](https://doi.org/10.1126/science.1074153). The version described in this paper is the first version, LT629781.1.

ACKNOWLEDGMENTS

We acknowledge Andrew Billings, Samantha Murphy, and Isa Bushko for assistance in the isolation and extraction of gDNA of bacterial isolates.

Funding for this project came from the University of Massachusetts–Amherst and the U.S. Department of Energy (DOE) Joint Genome Institute, a DOE Office of Science User Facility, which is supported by the Office of Science of the DOE under contract no. DE-AC02-05CH11231. This work was also supported by the DOE Genomic Sciences Program under contract no. DE-SC0016590.

REFERENCES

- Melillo JM, Steudler PA, Aber JD, Newkirk K, Lux H, Bowles FP, Catricala C, Magill A, Ahrens T, Morrisseau S. 2002. Soil warming and carbon-cycle feedbacks to the climate system. *Science* 298:2173–2176. <https://doi.org/10.1126/science.1074153>.
- Bergmann GT, Bates ST, Eilers KG, Lauber CL, Caporaso JG, Walters WA, Knight R, Fierer N. 2011. The under-recognized dominance of *Verrucomicrobia* in soil bacterial communities. *Soil Biol Biochem* 43:1450–1455. <https://doi.org/10.1016/j.soilbio.2011.03.012>.
- Davis KER, Joseph SJ, Janssen PH. 2005. Effects of growth medium, inoculum size, and incubation time on culturability and isolation of soil bacteria. *Appl Environ Microbiol* 71:826–834. <https://doi.org/10.1128/AEM.71.2.826-834.2005>.
- Eid J, Fehr A, Gray J, Luong K, Lyle J, Otto G, Peluso P, Rank D, Baybayan P, Bettman B, Bibillo A, Bjornson K, Chaudhuri B, Christians F, Cicero R, Clark S, Dalal R, Dewinter A, Dixon J, Foquet M, Gaertner A, Hardenbol P, Heiner C, Hester K, Holden D, Kearns G, Kong X, Kuse R, Lacroix Y, Lin S, Lundquist P, Ma C, Marks P, Maxham M, Murphy D, Park I, Pham T, Phillips M, Roy J, Sebra R. 2009. Real-time DNA sequencing from single polymerase molecules. *Science* 323:133–138. <https://doi.org/10.1126/science.1162986>.
- Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <https://doi.org/10.1038/nmeth.2474>.
- Huntemann M, Ivanova NN, Mavromatis K, Tripp HJ, Paez-Espino D, Palaniappan K, Szeto E, Pillay M, Chen IM, Pati A, Nielsen T, Markowitz VM, Kyrpides NC. 2015. The standard operating procedure of the DOE-JGI Microbial Genome Annotation Pipeline (MGAP v.4). *Stand Genomic Sci* 10:86. <https://doi.org/10.1186/s40793-015-0077-y>.
- Petersen TN, Brunak S, von Heijne G, Nielsen H. 2011. SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods* 8:785–786. <https://doi.org/10.1038/nmeth.1701>.
- Chen IM, Markowitz VM, Palaniappan K, Szeto E, Chu K, Huang J, Ratner A, Pillay M, Hadjithomas M, Huntemann M, Mikhailova N, Ovchinnikova G, Ivanova NN, Kyrpides NC. 2016. Supporting community annotation and user collaboration in the integrated microbial genomes (IMG) system. *BMC Genomics* 17:307. <https://doi.org/10.1186/s12864-016-2629-y>.
- van Teeseling MC, Pol A, Harhangi HR, van der Zwart S, Jetten MS, Op den Camp HJ, van Niftrik L. 2014. Expanding the verrucomicrobial methanotrophic world: description of three novel species of *Methylacidimicrobium* gen. nov. *Appl Environ Microbiol* 80:6782–6791. <https://doi.org/10.1128/AEM.01838-14>.
- Mohammadi S, Pol A, van Alen TA, Jetten MSM, Op den Camp HJM. 2017. *Methylacidiphilum fumariolicum* SolV, a thermoacidophilic ‘Knallgas’ methanotroph with both an oxygen-sensitive and -insensitive hydrogenase. *ISME J* 11:945–958. <https://doi.org/10.1038/ismej.2016.171>.
- Yin Y, Mao X, Yang J, Chen X, Mao F, Xu Y. 2012. dbCAN: a Web resource for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res* 40:W445–W451. <https://doi.org/10.1093/nar/gks479>.