

2020

Defoliation severity is positively related to soil solution nitrogen availability and negatively related to soil nitrogen concentrations following a multi-year invasive insect irruption

Emma Conrad-Rooney
Wellesley College

Audrey Barker Plotkin
Harvard University

Valerie J. Pasquarella
Boston University

Joseph Elkinton
University of Massachusetts Amherst

Jennifer L. Chandler
University of Massachusetts Amherst

Follow this and additional works at: https://scholarworks.umass.edu/nrc_faculty_pubs



Part of the [Environmental Monitoring Commons](#), and the [Natural Resources and Conservation Commons](#)
See next page for additional authors

Recommended Citation

Conrad-Rooney, Emma; Plotkin, Audrey Barker; Pasquarella, Valerie J.; Elkinton, Joseph; Chandler, Jennifer L.; and Matthes, Jaclyn Hatala, "Defoliation severity is positively related to soil solution nitrogen availability and negatively related to soil nitrogen concentrations following a multi-year invasive insect irruption" (2020). *Aob Plants*. 427.
<https://doi.org/10.1093/aobpla/plaa059>

This Article is brought to you for free and open access by the Environmental Conservation at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Environmental Conservation Faculty Publication Series by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

Authors

Emma Conrad-Rooney, Audrey Barker Plotkin, Valerie J. Pasquarella, Joseph Elkinton, Jennifer L. Chandler, and Jaclyn Hatala Matthes

STUDIES

Defoliation severity is positively related to soil solution nitrogen availability and negatively related to soil nitrogen concentrations following a multi-year invasive insect irruption

Emma Conrad-Rooney¹, Audrey Barker Plotkin², Valerie J. Pasquarella³, Joseph Elkinton⁴, Jennifer L. Chandler⁴ and Jaclyn Hatala Matthes^{1*}

¹Department of Biological Sciences, Wellesley College, Wellesley, MA 02481, USA, ²Harvard Forest, Harvard University, Petersham, MA 01366, USA, ³Department of Earth and Environment, Boston University, Boston, MA 02215, USA, ⁴Department of Environmental Conservation, University of Massachusetts Amherst, Amherst, MA 01003, USA

*Corresponding author's e-mail address: jmatthes@wellesley.edu

Plants, Ecosystems & Climate. Chief Editor: Mary Heskell

Associate Editor: Jeff Atkins

Abstract

Understanding connections between ecosystem nitrogen (N) cycling and invasive insect defoliation could facilitate the prediction of disturbance impacts across a range of spatial scales. In this study we investigated relationships between ecosystem N cycling and tree defoliation during a recent 2015–18 irruption of invasive gypsy moth caterpillars (*Lymantria dispar*), which can cause tree stress and sometimes mortality following multiple years of defoliation. Nitrogen is a critical nutrient that limits the growth of caterpillars and plants in temperate forests. In this study, we assessed the associations among N concentrations, soil solution N availability and defoliation intensity by *L. dispar* at the scale of individual trees and forest plots. We measured leaf and soil N concentrations and soil solution inorganic N availability among individual red oak trees (*Quercus rubra*) in Amherst, MA and across a network of forest plots in Central Massachusetts. We combined these field data with estimated defoliation severity derived from Landsat imagery to assess relationships between plot-scale defoliation and ecosystem N cycling. We found that trees in soil with lower N concentrations experienced more herbivory than trees in soil with higher N concentrations. Additionally, forest plots with lower N soil were correlated with more severe *L. dispar* defoliation, which matched the tree-level relationship. The amount of inorganic N in soil solution was strongly positively correlated with defoliation intensity and the number of sequential years of defoliation. These results suggested that higher ecosystem N pools might promote the resistance of oak trees to *L. dispar* defoliation and that defoliation severity across multiple years is associated with a linear increase in soil solution inorganic N.

Keywords: Defoliation; gypsy moth; insect disturbance; *Lymantria dispar*; nitrogen; soil solution nitrogen.

Introduction

Forests are an important global carbon sink that buffer the accumulation of atmospheric carbon dioxide (CO₂) from the combustion of fossil fuels causing climate change (IPCC 2014).

The uptake and sequestration of CO₂ during the regrowth of Northeastern US forests following agricultural abandonment over the last two centuries continues to play a significant role in

Received: 13 July 2020; Editorial decision: 28 October 2020; Accepted: 7 November 2020

© The Author(s) 2020. Published by Oxford University Press on behalf of the Annals of Botany Company.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

reducing the global greenhouse gas burden (Goodale et al. 2002; Pugh et al. 2019). The carbon sequestered by forest growth is only possible through important interactions with ecosystem nitrogen (N) cycling (Melillo et al. 2011). However, the ability of trees to grow and sequester carbon can also be impacted by disturbances such as invasive insect outbreaks, which are potentially increasing in frequency and severity with climate change (Seidl et al. 2017).

Gypsy moth (*Lymantria dispar*) is an invasive forest insect that was introduced to New England in the late 1860s, and since then *L. dispar* has periodically exhibited large irruptions during which populations increase by several orders of magnitude (Elkinton and Liebhold 1990; Liebhold et al. 2000). During a population irruption, *L. dispar* caterpillars consume large quantities of leaves, and extensive leaf herbivory often leads to widespread defoliation at the spatial scale of entire forest stands (Elkinton and Liebhold 1990). During the summers of 2015–18, forests of Southern New England experienced sustained irruptions of *L. dispar*, where the defoliated forest area was estimated to have at least doubled each year from 2015 to 2017 (Pasquarella et al. 2018b). Oaks (*Quercus* spp.) are the preferred hosts for *L. dispar* caterpillars, and although trees can typically withstand a single year of defoliation, successive years of *L. dispar* defoliation can lead to oak tree mortality (Barron and Patterson 2008; Dietze and Matthes 2014; Morin and Liebhold 2016).

Nitrogen (N) is a critical nutrient for both caterpillar and tree growth. Nitrogen from soil organic matter becomes available to tree roots through microbial decomposition that converts organic N into soluble inorganic ammonium (NH_4^+) and nitrate (NO_3^-) that trees take up from the soil solution through their roots (Bormann et al. 1977). Inorganic N taken up by tree roots is used to create the enzyme rubisco and pigments in leaves for photosynthesis. Nitrogen is also a critical nutrient for animals, such as *L. dispar* caterpillars, since they have high N requirements for protein synthesis (Mattson 1980). Therefore, the N concentrations of oak leaves could influence the extent and/or intensity of caterpillar herbivory and potentially play a role in the intensity of overall forest defoliation (Lindroth et al. 1997; Foss and Rieseke 2003; Milanović et al. 2014).

In laboratory environments, *L. dispar* caterpillars sometimes prefer leaves with higher leaf N concentrations, but higher leaf N concentrations can also suppress the relative caterpillar leaf consumption rates. Lindroth et al. (1997) found that higher leaf N concentrations suppress relative caterpillar leaf consumption rates (though only for males) and increase caterpillar survival and development rates. Laboratory-feeding preference studies have found that *L. dispar* caterpillars preferentially consume the *Quercus* species with the highest leaf N concentrations for some *Quercus* assemblages (Milanović et al. 2014), but other foliar differences might be more important than N concentrations for caterpillar consumption in other *Quercus* species assemblages (Foss and Rieseke 2003). While these experiments provide some insight into caterpillar herbivory and N dynamics in a laboratory context, behaviours under field conditions may differ.

In addition to the potential direct effects of leaf N concentrations on rates of herbivory, defoliation by *L. dispar* caterpillars also indirectly influences ecosystem N cycling. Herbivory redistributes N from tree leaves to the soil organic N pool through the production of frass (caterpillar faeces) and the accumulation of insect biomass which can subsequently become available to trees through microbial decomposition and mineralization. Insect herbivory can also either increase or decrease rates of N availability within soils through complex direct and indirect feedbacks (Hartley and Jones 2008).

Overall, higher inorganic soil N availability within a particular site could potentially increase *L. dispar* caterpillar herbivory (and hence internal N cycling rates) through higher leaf N concentrations. Alternatively, *L. dispar* caterpillar herbivory might decrease with higher N availability if caterpillars become satiated more quickly with higher N leaves and reduce their overall consumption as in laboratory studies (Lindroth et al. 1997). This study investigated the relationships among *L. dispar* herbivory, defoliation severity, soil and leaf N concentrations, and soil solution inorganic N availability at both individual trees and forest plots in a region defoliated by *L. dispar* in the years 2015–18. We tested two hypotheses assessing the relationships among defoliation and N concentrations and soil solution N at the scale of both individual trees and forest plots:

1. Higher concentrations of leaf and soil N are associated with higher rates of herbivory within trees, and higher concentrations of soil N are associated with higher rates of defoliation at the scale of forest stands.
2. Higher rates of inorganic N availability in soil solution are positively correlated to plot-scale defoliation intensity, since caterpillar herbivory should deliver a pulse of labile N to the forest floor.

To test these hypotheses, we measured the total N concentrations of leaves and soil and soil solution N availability in mixed temperate forests within areas that had been impacted by the *L. dispar* irruption in 2015–18. At individual trees we measured the extent of leaf herbivory, total leaf N concentrations and total soil N concentrations. Within forest plots we measured total soil N concentrations and rates of accumulation of soil solution N, and we assessed plot-level defoliation using a forest condition assessment product based on Landsat imagery (Pasquarella et al. 2017).

Methods

Study design and measurements of individual trees

Data for this study were collected at two scales: (i) individual red oak (*Quercus rubra*) trees and associated soil in the Amherst, MA, USA area and (ii) oak-dominated mixed forest plots in Central Massachusetts, USA. At each of these scales we measured N concentrations, herbivory and/or defoliation that are detailed in the following sections (Table 1). At the individual tree scale, 12 red oak (*Q. rubra*) trees growing alongside suburban roads and fields in the Amherst, MA area were selected as part of a previous study of *L. dispar* populations and their defoliation of oak trees. In early October 2018, we visually classified the percentage of overall tree defoliation on each tree as the percent of the tree canopy that was missing, presumably due to defoliation. The defoliation had been caused almost entirely by *L. dispar* larvae in May and June 2018. At the same time, leaf samples ($n = 30$) were

Table 1. Data were collected at individual trees and within forest plots.

Individual trees	Forest plot
<ul style="list-style-type: none"> • Leaf N concentrations (% N) • Soil N concentrations (% N) • Leaf herbivory (% herbivory) • Tree defoliation (% defoliation) 	<ul style="list-style-type: none"> • Soil N concentrations (% N) • Soil solution inorganic N (mg $\text{kg}^{-1} \text{ day}^{-1}$) • Defoliation intensity (data product from satellite imagery)

collected by hand from reachable branches around each tree to gather a spatially distributed representative sample and the percentage of leaf herbivory for each sampled leaf was visually scored by percent categories for herbivory in 10 % increments (e.g. 0–9 % consumed, 10–19 % consumed, etc.) in the laboratory. The average percentage of herbivory among the leaves for each tree was then multiplied by the number of leaves in each herbivory category by the midpoint percentage of the category (e.g. 4.5 % for the 0–9 % category) to estimate the average percent herbivory for each tree.

Study design and measurements of trees in forest plots

At the forest plot scale, ten 90 m × 90 m plots were established in 2019 in Central Massachusetts near the Quabbin Reservoir and the Harvard Forest to study the effects of sequential years of *L. dispar* defoliation. Each 90 m × 90 m plot was composed of three 20 m by 20 m subplots of oak-mixed forest (Fig. 1). One of the 10 plots was excluded from the analysis because it had only two oak trees in it (2.5 % of the trees in the plot were oaks), and therefore the drivers of *L. dispar* defoliation could have been different for this plot compared to the oak-mixed forest plots. All other forest plots had between 12 and 28 oak trees larger than 5 cm in diameter (mean = 20 cm, SD = 6.20 cm) and oaks comprised 15–33 % of the total trees in these plots.

These plots were selected to capture a range of defoliation intensities and durations, informed by data from a forest condition assessment product derived from Landsat imagery (Pasquarella 2017). We used data from this forest condition assessment product to quantify the cumulative severity of growing season defoliation at each plot across the 2015, 2016, 2017 and 2018 growing seasons (Pasquarella 2018a). Specifically, this satellite-based data product averaged the changes in growing season vegetation greenness relative to a fitted baseline model

for 2000–10 at each Landsat pixel in the broader New England region (Pasquarella 2017). To facilitate interpretation of results in this analysis, we multiplied the values of the data product by –1 so that a higher value represented a larger disturbance (i.e. as opposed to a more negative number representing a larger negative departure from mean greenness). Thus, the interpretation of the numeric values from this data product are analogous to a statistical z-score, where larger values reflect a larger departure from baseline conditions and are associated with more severe defoliation by *L. dispar* (Pasquarella 2017; Pasquarella et al. 2018b). For each plot, we calculated the average growing season defoliation for each year (2015–18) as the average of the data product pixels within a 45-m radius of the main plot centre. To represent the cumulative defoliation intensity within each plot, we summed the average growing season departures from mean greenness for the 2015–18 time period (sensu Foster 2017).

For some analyses, we used a categorical version of the Landsat-based forest condition assessment product, which separated the continuous greenness departure values into four greenness change classes: slight change (deviations 1–2 model root-mean-squared error; RMSE), moderate change (deviations 2–3 times model RMSE), large change (3–4 times model RMSE) and very large change (4–5 times model RMSE) (Pasquarella 2017, 2018a). For each plot, we calculated the average change class for each year (2015–18) as the average of the data product pixels within a 45-m radius of the main plot centre. We then classified plots as being defoliated within each year if the average change category was ‘moderate change’ or higher. Across the 2015–18 growing seasons, we calculated the total number of years defoliated for each plot as the sum of years that were classified as defoliated. This value of the number of years each plot was defoliated was used in conjunction with the cumulative defoliation intensity metric for some analyses.




Figure 1. Conceptual diagram outlining the data collected at the scales of individual trees in the Amherst, MA area ($n = 12$) and forest plots in Central Massachusetts ($n = 9$).

Soil and leaf nitrogen concentrations

To measure soil N concentrations for individual trees and forest plots, we collected samples of the soil organic horizon up to 10 cm depth using a 5-cm-diameter split PVC corer and a mallet. For the Amherst individual trees, three soil cores were taken from around each tree 0.5 m away from the trunk in different directions. If a tree was close to the road (less than about 3 m from road), the side of the tree closest to the road was avoided. The three soil cores associated with each tree were then homogenized into one composite sample. For each of the Quabbin forest subplots (Fig. 1), four soil samples were collected 5 m inward towards the centre of the plot from each of the four plot corners and then homogenized into a composite sample. Soil samples were dried for 24 h at 100 °C, sieved through a 2-mm sieve, and ground and homogenized with a ball mill for 5 min. For each soil sample, a 10-mg subsample was then packed into tins for elemental analysis and measured for the percentage of carbon and nitrogen in each soil sample (vario Micro Cube, Elementar Analysensysteme GmbH).

We also measured the N concentrations of leaves collected in October 2018 that were scored for herbivory from individual Amherst trees. It is important to note that since they were collected at the end of the growing season, they could have started senescing. The 30 leaves from each tree that were scored for herbivory were pooled and ground and homogenized together using a Wiley mill. Next, the ground leaves were more finely ground with a ball mill for 15 min. Then, 2 mg of each leaf sample were packed into elemental analysis tins and analysed for the percent nitrogen content (vario Micro Cube, Elementar Analysensysteme GmbH).

Soil solution N with resin bags

In order to measure soil solution N over time in the forest plots, we used an ion exchange resin bag method to capture mobile nitrate and ammonium using standard methods (Orwig et al. 2008; Templer and McCann 2010). For each resin bag, 10 g of resin were added to a packet created with a mesh stocking. A resin bag was buried 5–10 cm below the soil surface at each of the 10 Quabbin forest plots in the centre of the first subplot in early June (from 11 to 14 June). At each plot, the resin bag was collected 75–78 days (~2 months) after initial burying date to quantify accumulation of soil solution N during a time period that reflects the total soil solution N accumulated during the peak growing season (Sanders-DeMott et al. 2018).

After field collection, nitrate (NO_3^-) and ammonium (NH_4^+) were extracted from the resin using 2 M KCl following Orwig et al. (2008). We measured the resulting KCl extracts for NO_3^- and NH_4^+ concentrations with an Astoria-Pacific Discrete Analyser using colorimetric analysis (Astoria-Pacific, Inc., Clackamas, OR, USA). Each sample was run twice for replication and quality assurance. The resulting concentrations of NO_3^- and NH_4^+ in ppm were converted to mg per kg dry resin, the two replicates were averaged, and that value was divided by the number of days that the resin bag was deployed in order to get the average rates of NO_3^- and NH_4^+ production in mg per kg dry resin per day during the June–August summer growing season period. To calculate the total rate of inorganic N production for each resin bag, we added the mean rates of NO_3^- and NH_4^+ availability.

Statistical analyses

We calculated Pearson's correlation coefficients to test for correlations among N concentrations and soil solution N and

defoliation metrics. We fit one-way ANOVA models with post hoc Tukey honestly significant differences tests to determine whether the number of years a plot experienced defoliation (1, 2 or 3 years, defined as a categorical variable) explained variation in plot-level soil N concentrations or soil solution N. For soil and leaf % N variables we log-transformed the data before statistical testing. All statistical analyses were conducted and figures were produced in R (R Core Team 2019); we used base R (R Core Team 2019) in addition to R packages dplyr and tidyr to organize (Wickham et al. 2019; Wickham and Henry 2020), ggplot2 and cowplot to visualize data relationships (Wickham 2016; Wilke 2019) and agricolae to conduct the Tukey HSD test (de Mendiburu 2019). We defined the alpha value as 0.1 due to our small sample size, and throughout the text we report the actual P-value and r or r^2 value with each statistical test. We also used packages rgdal and raster in R to calculate the cumulative defoliation scores and to quantify the number of years defoliated from the Landsat-derived data products (Pebesma and Bivand 2005; Bivand et al. 2013, 2019; Hijmans 2020).

Results

Leaf and soil total N concentrations and defoliation at individual trees

Within individual *Q. rubra* trees measured in this study, there was no significant relationship between the extent of leaf herbivory and the N concentrations within leaves (Fig. 2A, $r = -0.35$, $P = 0.26$). This result is contrary to the hypothesis that more herbivory occurs on leaves with higher N concentrations. Tree defoliation had a significant negative correlation with soil total N concentrations, where at higher values of soil total N concentrations there was less total tree defoliation (Fig. 2B, $r = -0.57$, $P = 0.052$).

We found no significant relationship between soil total N concentrations and leaf total N concentrations at individual trees (Fig. 2C, $r = 0.23$, $P = 0.47$). Overall, there was larger variation in the soil total N concentrations (mean = 0.624 %, SD = 0.204, CV = 32.8 %) than the leaf total N concentrations (mean = 1.57 %, SD = 0.255, CV = 16.2 %). We found a statistically significant positive correlation between leaf-level herbivory and overall tree-level defoliation that supported a strong scaling relationship from the leaf to canopy scale (Fig. 2D, correlation = 0.86, $P < 0.01$).

Soil N cycling and plot-level defoliation

At the spatial scale of forest plots, there was a significant negative correlation between soil N concentration and the cumulative defoliation intensity from 2015 to 2018 (Fig. 3, $r = -0.65$, $P = 0.060$). This trend paralleled the negative relationship between defoliation and soil N concentrations associated with individual trees (Fig. 2B). There was not a significant relationship between the soil N concentrations and the number of sequential years of plot-level defoliation (Fig. 3, $P = 0.18$, $r^2 = 0.25$).

Plot-level soil solution N availability was significantly negatively correlated with the plot-level soil N concentrations (Fig. 4A, $r = -0.73$, $P = 0.026$), and soil solution N availability was significantly positively correlated with plot-level cumulative defoliation intensity (Fig. 4B, $r = 0.94$, $P < 0.01$). We found a strong positive effect of the number of sequential years of defoliation on soil solution N availability, where plots that were defoliated for three sequential years had amounts of inorganic N in soil solution that were up to 400 % larger than plots that were defoliated for only 1 year (Fig. 4B, $F(2,6) = 26.2$, $P < 0.01$, $r^2 = 0.86$).




Figure 2. Comparisons among *Q. rubra* leaf herbivory, tree defoliation, leaf % N and soil % N at the individual tree level. Each data point represents an individual tree and the soils associated with them ($n = 12$ trees). (A) Percent herbivory for individual leaves was not associated with leaf % N ($P = 0.26$). (B) The percentage of tree defoliation decreased as the amount of soil % N increased ($P = 0.052$). (C) There was no significant relationship between the leaf % N and soil % N ($P = 0.47$). (D) The measurements of individual leaf herbivory are significantly correlated with defoliation of the entire tree ($P < 0.01$).




Figure 3. Cumulative defoliation intensity from 2015 to 2018 at the plot scale ($n = 9$ plots) was negatively correlated with soil % N ($r = -0.65$, $P = 0.060$). This pattern suggested that there was a similar inverse relationship between defoliation and soil N at the plot level that was also found at the tree level (Fig. 2).

Discussion

Nitrogen concentrations and defoliation severity

Overall, there was a consistent negative correlation between soil N concentrations and defoliation severity at the tree and plot scales, where trees and plots with higher soil N concentrations

were associated with less intense defoliation (Figs 2B and 3). This result was in contrast to our hypothesis that higher ecosystem N concentrations would be associated with higher rates of caterpillar herbivory. While results from laboratory studies testing caterpillar herbivory preference with respect to leaf N concentrations have been mixed (Lindroth et al. 1997; Foss and Rieseke 2003; Milanović et al. 2014), our results suggest that trees and plots with lower soil N concentrations experienced more herbivory and overall defoliation. A potential mechanism for the observed pattern is that instead of travelling in search of higher N quality leaves, the caterpillars consume the leaves that they have access to and consume a relatively larger quantity of leaf mass to meet their nutritional requirements. This interpretation is in line with research that indicates that although *L. dispar* caterpillars travel within a tree, there is very little caterpillar dispersal between trees aside from initial wind dispersal, which is not directed dispersal and therefore does not select for high-quality trees (Weseloh 1987).

Though we found evidence of a relationship between soil N concentrations and *L. dispar* defoliation, it is challenging to infer causality since local soil N concentrations can change in response to tree defoliation. A prior field experiment on *Quercus* defoliation effects found that inorganic and soil N concentrations initially declined in the months following defoliation, potentially due to tree-regulated changes in root growth, activity, turnover or rhizodeposition in response to herbivory (Frost and Hunter 2004). However, in the subsequent year following the experimental defoliation, the soil N reservoir at 0–5 cm depth was a large sink for enhanced N deposition from insect frass (Frost and Hunter 2007).




Figure 4. (A) There was a significant negative relationship between plot-level soil N concentrations and the amount of N in soil solution that accumulated between early June and late August ($r = -0.73$, $P = 0.026$, $n = 9$ plots). (B) There was a significant positive relationship between N in soil solution, the defoliation intensity and the number of years defoliated ($r = 0.94$, $P < 0.01$, $n = 9$ plots).

Although the soil N concentrations were negatively associated with defoliation, the leaf total N concentrations were not significantly correlated to herbivory intensity (Fig. 2A). This pattern is potentially due to the lower range of variation in the leaf total N concentrations compared to the soil total N concentrations, and/or factors of leaf location (Niinemets 1997), seasonality (Xu and Griffin 2006; Yang *et al.* 2017), and/or phytochemical responses (Frost and Hunter 2008; War *et al.* 2018) that could have influenced our measurements of leaf N concentrations, or low sample size. Previous studies of forest N cycling using $\delta^{15}\text{N}$ have suggested that the below-ground soil N dynamics are a better indicator for ecosystem N cycling compared to above-ground N dynamics due to integrative lags in soil processes (Templer *et al.* 2007). In addition to the sources of measurement variation addressed in the previous section, this would also potentially help to explain the lack of coherence between soil and leaf N concentrations and the lack of a significant relationship between leaf N concentrations and herbivory in comparison to soil total N concentrations and defoliation (Fig. 2A and B).

Landscape-level studies have found that N levels in leaves are positively correlated with rates of atmospheric N deposition (Pitcairn *et al.* 2001), and that higher leaf N concentrations caused by the addition of N from human activities can lead to an increase in the size of insect populations (Throop and Lerdau 2004). However, results from this study are more in line with leaf-level studies that indicate that feeding rates of caterpillars increase when leaf N quality is lower (Lindroth *et al.* 1997). Recent research has found decreasing rates of atmospheric N deposition in North America (Lloret and Valiela 2016) and lower N availability in Northeastern forests (Durán *et al.* 2016; Elmore *et al.* 2016), which could potentially further increase defoliation intensity.

Plot soil solution nitrogen availability and defoliation severity

Soil solution N reflects the availability of inorganic N to plant roots, whereas the soil total N concentration represents the long-term pools of N in the soil that are mostly composed of soil organic matter. Defoliation transfers N from the canopy to the soil, particularly in the forms of frass (caterpillar faeces), green leaves that fall, and insect biomass. An increase in availability of soil N for trees with defoliation could potentially help trees to recover following stress from herbivory. Within forest plots, cumulative defoliation intensity and soil solution N availability were significantly positively correlated (Fig. 4B). There was also

a strong relationship between the number of years of sequential defoliation and soil solution N availability, where plots that were defoliated 3 years in a row had much higher amounts of soil solution N than plots defoliated 1 or 2 years.

Other studies have found ephemeral enhancements in inorganic N available in soil solution in the months following *L. dispar* defoliation (Lovett *et al.* 2002; Russell *et al.* 2004). In both studies, frass N was quickly taken up from the soil with enhanced plant N uptake in a poplar plantation (Russell *et al.* 2004) and elevated microbial N uptake and immobilization in a mixed hardwood forest (Lovett *et al.* 2002; Christenson *et al.* 2002). Both studies found that the N redistributed to the soil solution inorganic N pool by a single severe defoliation event was not lost by leaching at the ecosystem and watershed scales (Lovett *et al.* 2002; Russell *et al.* 2004). A legacy of elevated inorganic N was also observed in birch forests that experienced 2–3 severe sequential defoliation events and was explained as enhanced decomposer activity (Kaukonen *et al.* 2013). The pattern in our study could potentially be the result of lag effects following defoliation where N mineralization might be enhanced through higher mortality in tree fine roots or more dynamic microbial growth and mortality in microbial population cycles.

There could be alternate explanations for the positive relationship between soil solution N availability and defoliation severity found in this study, such as other site conditions that influence available N. For example, it is possible that particular site conditions such as soil moisture influence soil N cycling (Landesman and Dighton 2010) and that these conditions also co-vary with favourable conditions for *L. dispar* growth and survival at these plots. It is also possible that the severe tree stress caused by 3 years of sequential defoliation led to higher rates of root mortality, which could represent an additional pulse of organic matter to the soil N pool and the higher availability of soil solution N observed in our study. The results from our study highlight the utility of field observations and experiments that capture high and recurrent disturbance severities and monitor changes in ecosystem processes over several years.

Coherence with ecosystem N cycling at trees and plots

In comparing leaf-level to tree-level caterpillar defoliation measurements, we found that individual leaf-level caterpillar herbivory was significantly positively correlated with

assessments of tree-level defoliation (Fig. 2D). This pattern suggested that the impact of caterpillar herbivory of individual leaves scales proportionally to overall defoliation at the tree level, even when leaves were only collected from branches that could be reached from the ground.

We found no correlation between soil and leaf total N concentration at the scale of individual trees (Fig. 2C). Our measurements for the leaf total N concentrations could be impacted by the collection of leaves that could only be reached from the ground, since shade leaves tend to have lower N concentrations than sunlit leaves (Niinemets 1997). Trees that experience herbivory have also been observed to reduce leaf N concentration in response to defoliation (Frost and Hunter 2008) and phytochemical induction of plant defences can also impact leaf N concentrations (War et al. 2018). However, another study found that hemlock foliar N increased following herbivory by hemlock woolly adelgid (Schaeffer et al. 2017). Additionally, leaves in this study were collected at the end of the growing season in October 2018, when leaf N concentrations are typically at the lowest levels of the season, particularly compared to the levels in May and June when *L. dispar* larvae were feeding (Scriber and Slansky 1981; Xu and Griffin 2006; Yang et al. 2017). We found that the values of *Q. rubra* foliar N concentrations (mean = 1.57, SD = 0.255) in this study were about 30 % lower than *Q. rubra* foliar N measurements from a prior Harvard Forest study conducted from 1988 to 1991 (mean = 2.28, SD = 0.23; Martin and Aber 1992). Together, the factors of shaded leaf collection, the seasonal timing of collection during autumn and/or phytochemical responses to defoliation could help to explain the lower-than-expected concentrations of leaf N and lack of a relationship with defoliation.

Across the different forest plots, soil total N concentrations and the amount of N in soil solution were negatively correlated (Fig. 4A). This pattern could be due to the fact that the soil organic matter contains most of the total soil N, and a primary source of inorganic N in soil solution is the decomposition of this soil organic matter (McClaugherty et al. 1985; Aber et al. 1990). Therefore, it is consistent that forest plots with higher amounts of soil solution N have lower soil N concentrations, while soils with larger N concentrations have lower amounts of N in soil solution. The measurements of ammonium and nitrate in soil solution in this study were within the same range of variation as other studies that used this same method in mixed Northeastern US forests (Sanders-DeMott et al. 2018).

Ecosystem level implications

Lymantria dispar defoliation events have both short- and long-term impacts on forest dynamics. In the short term, defoliation events can alter canopy light transmittance, temperature and moisture content of the soil, and forest nutrient cycling (Gandhi and Herms 2010). Over time, *L. dispar* defoliation events in the Northeast can lead to large-scale oak mortality and changes in forest stand composition and dynamics including a decrease in oak dominance (Barron and Patterson 2008; Morin and Liebhold 2016). Changes in forest composition due to oak mortality from *L. dispar* outbreaks are predicted to negatively impact eastern North American wildlife, which rely heavily on the consumption of acorns from oak masts (McShea et al. 2007). Increased climate variability including drought events may impact the intensity and frequency of *L. dispar* outbreaks because one of the main controls on *L. dispar* populations, the fungal pathogen *Entomophaga maimaiga*, proliferates with higher levels of precipitation (Hajek and Humber 1997; Reilly et al. 2014).

Variation in defoliation rates across the broader region of Northeastern US forests could drive variation in relationships

between defoliation and ecosystem N cycling (Davidson et al. 2001). Patterns within our study could also have been influenced by trends in regional patterns of declining N deposition in Northeastern US forests (Lloret and Valiela 2016) where impacts of climate change, including increasing temperatures, lower snow accumulation and changing seasonality, may be contributing to a decline in N availability (Durán et al. 2016; Elmore et al. 2016). This pattern has implications for our study because of the potential relationship between higher rates of defoliation at lower soil N concentrations, which could predict a larger extent of defoliation with future declines in N pollution.

Conclusions

Overall, we found that at the scales of individual trees and forest stands, there was a consistent negative relationship between soil N concentrations and defoliation intensity. We found no significant relationship between leaf N concentrations and the amount of herbivory. This study provides important in situ characterization of nitrogen cycling at multiple scales that could help to improve predictions of nutrient cycling within ecosystem models in response to recurrent and severe landscape-level defoliation events. Declines in dominant forest canopy species like oaks can have widespread impacts on forest ecosystem processes such as decomposition, carbon sequestration and nutrient cycling, as well as on processes in nearby aquatic ecosystems (Ellison et al. 2005). Forest insects and pathogens also impact ecosystem services such as water quality and carbon sequestration (Dale et al. 2001), which are highly influenced by ecosystem N cycling. Additionally, changes in climate can directly impact the *E. maimaiga* fungal control to *L. dispar* populations, potentially increasing the likelihood of future *L. dispar* outbreaks (Hajek and Humber 1997; Reilly et al. 2014). The strong relationships we found between ecosystem N cycling and defoliation following a severe, multi-year invasive insect irruption highlight the importance of investigating the long-term effects of severe and recurring ecosystem disturbances. Particularly considering that these types of disturbances are likely to increase in frequency and intensity, long-term monitoring efforts are critical to understanding ecosystem feedbacks and potential thresholds.

Data Availability

Data associated with this manuscript are available through the Environmental Data Initiative (Conrad-Rooney et al. 2020a) and in the Harvard Forest Data Archive (Conrad-Rooney et al. 2020b).

Sources of Funding

This research was supported by the Wellesley Science Center Summer Research Program and the Frost Endowed Environmental Science Studies Fund, the US National Science Foundation (NSF) MacroSystems Biology grant NSF-163840 and the Harvard Forest NSF Research Experience for Undergraduates (REU) program grant DBI-1459519.

Contributions by the Authors

All authors collaboratively designed the overall study and collected data. A.B.P. and V.J.P. led the establishment of forest

plots used in this study and J.E. and J.L.C. designed the study of individual trees. E.C.-R. led the data analysis, figure development, and writing for the manuscript, with support from J.H.M. A.B.P., V.J.P., J.E., and J.L.C. provided feedback for data analysis and edited the manuscript.

Conflict of Interest

None declared.

Acknowledgements

Thank you to Elizabeth Bouchard, Jonathan Gewirtzman, Alden Griffith, Kristina Jones, Danelle Laflower, David Orwig, Manisha Patel, Pamela Templer and Greta VanScoy for their guidance, support and assistance with this project. Thank you to Savanna Brown, Sofia Kruszka, Brianna Martinez, Samantha Matson, Elise Miller and Nathan Oalican for their assistance with field and laboratory work.

Literature Cited

- Aber JD, Wessman CA, Peterson DL, Melillo JM, Fownes JH. 1990. Remote sensing of litter and soil organic matter decomposition in forest ecosystems. In: Hobbs RJ, Mooney HA, eds. *Remote sensing of biosphere functioning*. New York: Springer, 87–103.
- Barron ES, Patterson WA. 2008. Monitoring the effects of gypsy moth defoliation on forest stand dynamics on Cape Cod, Massachusetts: sampling intervals and appropriate interpretations. *Forest Ecology and Management* 256:2092–2100.
- Bivand R, Keitt T, Rowlingson B. 2019. *rgdal: bindings for the “geospatial” data abstraction library*. R package version 1.4-8.
- Bivand RS, Pebesma EJ, Gomez-Rubio V. 2013. *Applied spatial data analysis with R*, 2nd edn. New York: Springer, 83–125.
- Bormann FH, Likens GE, Melillo JM. 1977. Nitrogen budget for an aggrading northern hardwood forest ecosystem. *Science* 196:981–983.
- Christenson LM, Lovett GM, Mitchell MJ, Groffman PM. 2002. The fate of nitrogen in gypsy moth frass deposited to an oak forest floor. *Oecologia* 131:444–452.
- Conrad-Rooney E, Barker Plotkin A, Matthes JH. 2020a. Leaf and soil nitrogen following gypsy moth defoliation in Central Massachusetts 2018–2019 ver 1. *Environmental Data Initiative*. doi:10.6073/pasta/b9361a509c3371cd58bdcd1da7a8b71f.
- Conrad-Rooney E, Barker Plotkin A, Matthes JH. 2020b. Leaf and soil nitrogen following Gypsy Moth Defoliation in Central Massachusetts 2018–2019. *Harvard Forest Data Archive*: HF344.
- Dale VH, Joyce LA, McNulty S, Neilson RP, Ayres MP, Flannigan MD, Hanson PJ, Irland LC, Lugo AE, Peterson CJ, Simberloff D, Swanson FJ, Stocks BJ, Wotton BM. 2001. Climate change and forest disturbances: climate change can affect forests by altering the frequency, intensity, duration, and timing of fire, drought, introduced species, insect and pathogen outbreaks, hurricanes, windstorms, ice storms, or landslides. *BioScience* 51:723–734.
- Davidson CB, Johnson JE, Gottschalk KW, Amateis RL. 2001. Prediction of stand susceptibility and gypsy moth defoliation in coastal plain mixed pine-hardwoods. *Canadian Journal of Forest Research* 31:1914–1921.
- Dietze MC, Matthes JH. 2014. A general ecophysiological framework for modelling the impact of pests and pathogens on forest ecosystems. *Ecology Letters* 17:1418–1426.
- Durán J, Morse JL, Groffman PM, Campbell JL, Christenson LM, Driscoll CT, Fahey TJ, Fisk MC, Likens GE, Melillo JM, Mitchell MJ, Templer PH, Vadeboncoeur MA. 2016. Climate change decreases nitrogen pools and mineralization rates in northern hardwood forests. *Ecosphere* 7:e01251.
- Elkinton JS, Liebholt AM. 1990. Population dynamics of gypsy moth in North America. *Annual Review of Entomology* 35:571–596.
- Ellison AM, Bank MS, Clinton BD, Colburn EA, Elliott K, Ford CR, Foster DR, Kloeppel BD, Knoepp JD, Lovett GM, Mohan J, Orwig DA, Rodenhouse NL, Sobczak WV, Stinson KA, Stone JK, Swan CM, Thompson J, Holle BV, Webster JR. 2005. Loss of foundation species: consequences for the structure and dynamics of forested ecosystems. *Frontiers in Ecology and the Environment* 3:479–486.
- Elmore AJ, Nelson DM, Craine JM. 2016. Earlier springs are causing reduced nitrogen availability in North American eastern deciduous forests. *Nature Plants* 2:16133.
- Foss LK, Riese LK. 2003. Species-specific differences in oak foliage affect preference and performance of gypsy moth caterpillars. *Entomologia Experimentalis et Applicata* 108:87–93.
- Foster JR. 2017. Xylem traits, leaf longevity and growth phenology predict growth and mortality response to defoliation in northern temperate forests. *Tree Physiology* 37:1151–1165.
- Frost CJ, Hunter MD. 2004. Insect canopy herbivory and frass deposition affect soil nutrient dynamics and export in oak mesocosms. *Ecology* 85:3335–3347.
- Frost CJ, Hunter MD. 2007. Recycling of nitrogen in herbivore feces: plant recovery, herbivore assimilation, soil retention, and leaching losses. *Oecologia* 151:42–53.
- Frost CJ, Hunter MD. 2008. Insect herbivores and their frass affect *Quercus rubra* leaf quality and initial stages of subsequent litter decomposition. *Oikos* 117:13–22.
- Gandhi KJK, Herms DA. 2010. Direct and indirect effects of alien insect herbivores on ecological processes and interactions in forests of eastern North America. *Biological Invasions* 12:389–405.
- Goodale CL, Apps MJ, Birdsey RA, Field CB, Heath LS, Houghton RA, Jenkins JC, Kohlmaier GH, Kurz W, Liu S, Nabuurs G-J, Nilsson S, Shvidenko AZ. 2002. Forest carbon sinks in the northern hemisphere. *Ecological Applications* 12:891–899.
- Hajek AE, Humber RA. 1997. Formation and germination of *Entomophaga maimaiga* azygospores. *Canadian Journal of Botany* 75:1739–1747.
- Hartley SE, Jones TH. 2008. Insect herbivores, nutrient cycling, and plant productivity. In: Weisser WW, Siemann E, eds. *Insects and ecosystem function*. Berlin: Springer, 27–52.
- Hijmans RJ. 2020. *raster: geographic data analysis and modeling*. R package version 3.0-12.
- IPCC. 2014. Climate change 2014: synthesis report. In: Core Writing Team, Pachauri RK, Meyer LA, eds. *Contribution of working groups I, II and III to the fifth assessment report of the Intergovernmental Panel on Climate Change*. Geneva: IPCC.
- Kaukonen M, Ruotsalainen AL, Wäli PR, Männistö MK, Setälä H, Saravesi K, Huusko K, Markkola A. 2013. Moth herbivory enhances resource turnover in subarctic mountain birch forests? *Ecology* 94:267–272.
- Landesman WJ, Dighton J. 2010. Response of soil microbial communities and the production of plant-available nitrogen to a two-year rainfall manipulation in the New Jersey Pinelands. *Soil Biology and Biochemistry* 42:1751–1758.
- Liebholt A, Elkinton J, Williams D, Muzika R-M. 2000. What causes outbreaks of the gypsy moth in North America? *Population Ecology* 42:257–266.
- Lindroth RL, Klein KA, Hemming JDC, Feuker AM. 1997. Variation in temperature and dietary nitrogen affect performance of the gypsy moth (*Lymantria dispar* L.). *Physiological Entomology* 22:55–64.
- Lloret J, Valiela I. 2016. Unprecedented decrease in deposition of nitrogen oxides over North America: the relative effects of emission controls and prevailing air-mass trajectories. *Biogeochemistry* 129:165–180.
- Lovett GM, Christenson LM, Groffman PM, Jones CG, Hart JE, Mitchell MJ. 2002. Insect defoliation and nitrogen cycling in forests. *BioScience* 52:335–341.
- Martin M, Aber J. 1992. Canopy chemistry study at Harvard Forest 1992. *Harvard Forest*.
- Mattson WJ. 1980. Herbivory in relation to plant nitrogen content. *Annual Review of Ecology and Systematics* 11:119–161.
- McClougherty CA, Pastor J, Aber JD, Melillo JM. 1985. Forest litter decomposition in relation to soil nitrogen dynamics and litter quality. *Ecology* 66:266–275.
- McShea WJ, Healy WM, Devers P, Fearer T, Koch FH, Stauffer D, Waldon J. 2007. Forestry matters: decline of oaks will impact wildlife in hardwood forests. *Journal of Wildlife Management* 71:1717–1728.
- Melillo JM, Butler S, Johnson J, Mohan J, Steudler P, Lux H, Burrows E, Bowles F, Smith R, Scott L, Vario C, Hill T, Burton A, Zhou Y-M, Tang J.

2011. Soil warming, carbon–nitrogen interactions, and forest carbon budgets. *Proceedings of the National Academy of Sciences of the United States of America* **108**:9508–9512.
- de Mendiburu F. 2019. *agricolae: statistical procedures for agricultural research*. R package version 1.3-1.
- Milanović S, Lazarević J, Popović Z, Miletic Z, Kostić M, Radulović Z, Karadžić D, Vuleta A. 2014. Preference and performance of the larvae of *Lymantria dispar* (Lepidoptera: Lymantriidae) on three species of European oaks. *European Journal of Entomology* **111**:371–378.
- Morin RS, Liebhold AM. 2016. Invasive forest defoliator contributes to the impending downward trend of oak dominance in eastern North America. *Forestry* **89**:284–289.
- Niinemets Ü. 1997. Role of foliar nitrogen in light harvesting and shade tolerance of four temperate deciduous woody species. *Functional Ecology* **11**:518–531.
- Orwig DA, Cobb RC, D'Amato AW, Kizlinski ML, Foster DR. 2008. Multi-year ecosystem response to hemlock woolly adelgid infestation in Southern New England forests. *Canadian Journal of Forest Research* **38**:834–843.
- Pasquarella VJ. 2018a. *Landsat-based gypsy moth defoliation assessment (Southern New England) (version 3.0)*. Zenodo: doi:10.5281/zenodo.1163679.
- Pasquarella VJ, Bradley BA, Woodcock CE. 2017. Near-real-time monitoring of insect defoliation using Landsat time series. *Forests* **8**:275.
- Pasquarella VJ, Elkinton JS, Bradley BA. 2018b. Extensive gypsy moth defoliation in Southern New England characterized using Landsat satellite observations. *Biological Invasions* **20**:3047–3053.
- Pebesma EJ, Bivand RS. 2005. Classes and methods for spatial data in R. *R News* **5**.
- Pitcairn CER, Leith ID, Fowler D, Hargreaves KJ, Moghaddam M, Kennedy VH, Granat L. 2001. Foliar nitrogen as an indicator of nitrogen deposition and critical loads exceedance on a European scale. *Water, Air, and Soil Pollution* **130**:1037–1042.
- Pugh TAM, Lindeskog M, Smith B, Poulter B, Arneth A, Haverd V, Calle L. 2019. Role of forest regrowth in global carbon sink dynamics. *Proceedings of the National Academy of Sciences of the United States of America* **116**:4382–4387.
- R Core Team. 2019. *R: a language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Reilly JR, Hajek AE, Liebhold AM, Plymale R. 2014. Impact of *Entomophaga maimaiga* (Entomophthorales: Entomophthoraceae) on outbreak gypsy moth populations (Lepidoptera: Erebidae): the role of weather. *Environmental Entomology* **43**:632–641.
- Russell CA, Kosola KR, Paul EA, Robertson GP. 2004. Nitrogen cycling in poplar stands defoliated by insects. *Biogeochemistry* **68**:365–381.
- Sanders-DeMott R, Sorensen PO, Reinmann AB, Templer PH. 2018. Growing season warming and winter freeze–thaw cycles reduce root nitrogen uptake capacity and increase soil solution nitrogen in a northern forest ecosystem. *Biogeochemistry* **137**:337–349.
- Schaeffer RN, Soltis NE, Martin JL, Brown AL, Gómez S, Preisser EL, Orians CM. 2017. Seasonal variation in effects of herbivory on foliar nitrogen of a threatened conifer. *AoB PLANTS* **9**:plx007; doi:10.1093/aobpla/plx007.
- Scriber JM, Slansky F Jr. 1981. The nutritional ecology of immature insects. *Annual Review of Entomology* **26**:183–211.
- Seidl R, Thom D, Kautz M, Martin-Benito D, Peltoniemi M, Vacchiano G, Wild J, Ascoli D, Petr M, Honkaniemi J, Lexer MJ, Trotsiuk V, Mairota P, Svoboda M, Fabrika M, Nagel TA, Reyer CPO. 2017. Forest disturbances under climate change. *Nature Climate Change* **7**:395–402.
- Templer PH, Arthur MA, Lovett GM, Weathers KC. 2007. Plant and soil natural abundance delta (15)N: indicators of relative rates of nitrogen cycling in temperate forest ecosystems. *Oecologia* **153**:399–406.
- Templer PH, McCann TM. 2010. Effects of the hemlock woolly adelgid on nitrogen losses from urban and rural northern forest ecosystems. *Ecosystems* **13**:1215–1226.
- Throop HL, Lerdau MT. 2004. Effects of nitrogen deposition on insect herbivory: implications for community and ecosystem processes. *Ecosystems* **7**:109–133.
- War AB, Taggar GK, Hussain B, Taggar MS, Nair RM, Sharma HC. 2018. Plant defence against herbivory and insect adaptations. *AoB PLANTS* **10**:ply037; doi:10.1093/aobpla/ply037.
- Weseloh RM. 1987. Dispersal and survival of gypsy moth larvae. *Canadian Journal of Zoology* **65**:1720–1723.
- Wickham H. 2016. *ggplot2: elegant graphics for data analysis*. New York: Springer-Verlag.
- Wickham H, Henry L. 2020. *tidyr: tidy messy data*. R package version 1.0.2.
- Wickham H, François R, Henry L, Müller K. 2019. *dplyr: a grammar of data manipulation*. R package version 0.8.3.
- Wilke CO. 2019. *cowplot: streamlined plot theme and plot annotations for “ggplot2”*. R package version 1.0.0.
- Xu C-Y, Griffin KL. 2006. Seasonal variation in the temperature response of leaf respiration in *Quercus rubra*: foliage respiration and leaf properties. *Functional Ecology* **20**:778–789.
- Yang H, Yang X, Heskell M, Sun S, Tang J. 2017. Seasonal variations of leaf and canopy properties tracked by ground-based NDVI imagery in a temperate forest. *Scientific Reports* **7**:1267.