

University of Massachusetts Amherst

**ScholarWorks@UMass Amherst**

---

Environmental Conservation Faculty Publication  
Series

Environmental Conservation

---

2020

## **Evaluating the Likelihood of Tree Failure in Naples, Florida (United States) Following Hurricane Irma**

Ryan W. Klein

Andrew K. Koeser

Brian Kane

Shawn M. Laundry

Heather Shields

*See next page for additional authors*

Follow this and additional works at: [https://scholarworks.umass.edu/nrc\\_faculty\\_pubs](https://scholarworks.umass.edu/nrc_faculty_pubs)



Part of the [Environmental Monitoring Commons](#), and the [Natural Resources and Conservation Commons](#)

---

---

**Authors**

Ryan W. klein, Andrew K. Koeser, Brian Kane, Shawn M. Laundry, Heather Shields, Stephen Lloyd, and Gail Hansen

---

## Article

# Evaluating the Likelihood of Tree Failure in Naples, Florida (United States) Following Hurricane Irma

Ryan W. Klein <sup>1,\*</sup>, Andrew K. Koeser <sup>2</sup>, Brian Kane <sup>3</sup> , Shawn M. Landry <sup>4</sup>, Heather Shields <sup>5</sup>, Stephen Lloyd <sup>6</sup> and Gail Hansen <sup>7</sup>

<sup>1</sup> Department of Environmental Horticulture, CLCE, IFAS, University of Florida, 100A Mehrhof Hall, Gainesville, FL 32611, USA

<sup>2</sup> Department of Environmental Horticulture, CLCE, IFAS, University of Florida–Gulf Coast Research and Education Center, 14625 County Road 672, Wimauma, FL 33598, USA; akoeser@ufl.edu

<sup>3</sup> Department of Environmental Conservation, University of Massachusetts, 160 Holdsworth Way, Amherst, MA 01003, USA; bkane@eco.umass.edu

<sup>4</sup> School of Geosciences, University of South Florida, 4202 East Fowler Avenue, Tampa, FL 33620, USA; landry@usf.edu

<sup>5</sup> City Arborist–Project Manager, 280 Riverside Circle, Naples, FL 34102, USA; hshields@naplesgov.com

<sup>6</sup> Florida Forest Service–U&CF Partnership Coordinator, 4316 International Speedway Blvd., Daytona Beach, FL 02124, USA; Stephen.Lloyd@freshfromflorida.com

<sup>7</sup> Department of Environmental Horticulture, CLCE, IFAS, University of Florida, Building 550 Room 101, Gainesville, FL 32611, USA; ghansen@ufl.edu

\* Correspondence: ryanwklein@ufl.edu

Received: 14 April 2020; Accepted: 23 April 2020; Published: 25 April 2020



**Abstract:** Trees in residential landscapes provide many benefits, but can injure persons and damage property when they fail. In hurricane-prone regions like Florida, USA, the regular occurrence of hurricanes has provided an opportunity to assess factors that influence the likelihood of wind-induced tree failure and develop species failure profiles. We assessed open-grown trees in Naples, Florida, following the passage of Hurricane Irma in September 2017 to determine the effect of relevant factors on the degree of damage sustained by individual trees. Of 4034 assessed individuals ( $n = 15$  species), 74% sustained no damage, 4% sustained only minor damage (i.e., minimal corrective pruning needed), 6% sustained significant damage (i.e., major corrective pruning needed), and 15% were whole-tree failures (i.e., overturned trees or trees requiring removal). The proportion of individuals in each damage category varied among species, stem diameter at 1.4 m above ground, and the presence of utility lines, which was a proxy for maintenance. We compared our results with the findings of seven previous hurricanes in the region to explore species' resilience in hurricanes.

**Keywords:** hurricane; tree risk assessment; urban forest strike team; species failure profile; likelihood of failure

## 1. Introduction

Wind loads often induce tree failure, and many studies have quantified the effects of wind events on forests. In an extensive review, Everham and Brokaw [1] synthesized the findings of 42 studies of tree failure during catastrophic wind events to compile a database of wind resistance ratings for different species in temperate and tropical climates. However, their synthesis only included studies on forest- or plantation-grown trees, which resulted in two important limitations. First, the form of open-grown trees differs from that of forest- or plantation-grown trees [2]; secondly, ornamental species common in residential landscapes are typically absent from natural forest stands or plantations. Studies have examined the wind-induced failure of open-grown trees in temperate [3] and tropical and

subtropical climates [4–7], but predicting failure is problematic because empirical data on relevant parameters are sparse.

Hurricanes annually impact communities along the Atlantic and Gulf Coasts in the United States, sometimes causing widespread and costly damage. Hurricanes often induce tree failure, and studies have (i) quantified the cost of tree damage and debris [8], (ii) evaluated the likelihood of tree failure during catastrophic wind events [1,4–7,9], (iii) assessed damaged trees [10] and property [11,12], (iv) considered their effect on urban tree populations [12], and (v) gauged perceptions of the urban forest following a hurricane [13]. However, predicting the likelihood of tree failure during storms remains a challenge.

Many factors influence the likelihood of tree failure. Factors can be categorized as intrinsic or extrinsic to the tree. Intrinsic factors include size, wood density, and the presence or absence of structural defects (e.g., decay, weakly attached branches) and foliage. Studies have consistently reported a greater likelihood of failure for taller trees with larger trunk diameter for trees in both cooler [3,9,14,15] and warmer climates [4,11]. Larger trees experience greater drag and drag-induced bending moments because of a greater crown area, a higher center of pressure, and greater exposure to the wind. The likelihood of tree failure tends to be lower with denser wood [7,11,15–18]. Leafless trees experience considerably less drag than in-leaf trees [19], which presumably explains the lower likelihood of failure in catastrophic wind events of dormant trees [9] or trees that readily shed leaves [4,20]. An integral part of visual tree assessment is identifying the presence and severity of structural defects [21–25], but only severe defects, rather than simply the presence of defects, have been linked to increased likelihood of tree failure [9].

Extrinsic factors, such as site conditions, can influence and even mask species-specific intrinsic factors because they influence growth form. For several species, the form of open-grown individuals differed substantially from that of forest- or plantation-grown trees [2]. However, within a particular site, many studies have demonstrated that the likelihood of failure varies among species [1,3,4,6,7,9,11,20]. Studies have also shown instances where native species were more resilient to wind than nonnative species [5,12,26] especially in hurricane-prone regions, but the converse has also been documented [11,27].

In developed landscapes, likelihood of failure was lower when trees had a greater volume of soil for roots [11,12], and among trees growing in clusters [12]. In developed landscapes, management practices may also influence the likelihood of failure in catastrophic wind events, but a consensus is lacking among the few studies that have investigated this effect. Some studies have found that pruned trees were less likely to fail than trees that had not been pruned [5,11,12], but one [9] reported a minimal effect of pruning on the likelihood of failure. It is also important to remember that the effects of management may be negligible when winds are catastrophic and sufficient to cause even structurally sound trees to fail.

Given the complexity of modeling tree failure in catastrophic wind events, species failure profiles can be useful. Failure profiles commonly involve an evaluation of structural and environmental factors that lead to tree failure. For example, Edberg and Berry [28] developed a failure profile for coast live oak in California (*Quercus agrifolia* Née) using data from the California Tree Failure Report Program (CTFRP). Structural factors included the location of the failure (trunk; branch; root; above, at, or below ground level); environmental factors included precipitation, wind speed, soil, and topography. They also considered the size of the tree, stand characteristics, common defects, cultural practices, and the presence of decay [28]. The most common type of failure found in coast live oak (83%) was related to decay found in the lower trunk and roots, resulting from saturated soil conditions [28]. Edberg et al. [29] also used data from the CTFRP to develop a failure profile for Monterey pine (*Pinus radiata* D. Don) and found that 75% of failures were of the heavy lateral limbs; of these, 74% failed away from the branch attachment, suggesting that wood strength or load distribution, not a weak branch attachment, was the cause.

In the United States, hurricanes regularly occur in Florida, providing ample opportunities to develop species failure profiles [5,10–12]. Since climate change models predict an increase in hurricane frequency and severity, the objectives of this study were to determine (i) factors that influenced the likelihood of failure, and (ii) what type of failure was most common.

## 2. Materials and Methods

Hurricane Irma reached coastal Florida as a category 3 hurricane on 10 September 2017. As it moved northward to Naples, Florida, USA (26.1420° N, 81.7948° W), the storm diminished in strength, with the southern and eastern portions of the city experiencing category 2 hurricane winds and the northern portion of the city experiencing category 1 winds [30]. According to the Saffir–Simpson scale, category 2 hurricanes will cause “many shallowly rooted trees [to] be snapped or uprooted and block numerous roads” and category 1 hurricanes will cause “large branches of trees [to] snap and shallowly rooted trees may be toppled” [31]. Prior to Hurricane Irma, Naples had a complete inventory of 19,638 public trees, which included 79 species. The current study focused on a subset of the inventory that included city-owned street trees in the public right-of-way, but excluded trees in parks and medians (i.e., the strip of unpaved land that separates lanes of opposing traffic on a divided highway).

Following the hurricane and initial cleanup, from 30 October 2017, through 3 November 2017, the city of Naples conducted an assessment of trees in the public right-of-way. The assessment was carried out by a Florida Forest Service Urban Forest Strike Team (UFST). The UFST consisted of four arboricultural professionals who were divided into two crews. Each crew had one assessor with the International Society of Arboriculture’s (ISA) Tree Risk Assessment Qualification (TRAQ). All four assessors were ISA-Certified Arborists with previous experience or training in post-disaster tree risk assessment and incident command systems [32]. The crews located trees using smartphones and the city’s inventory, which included species, trunk diameter 1.4 m above ground (DBH), height, and GPS coordinates in a GIS-based data collection app (Collector, Esri, Redlands, CA, USA).

The crews conducted a level 1 limited visual assessment [25] of each tree, using a one-year timeframe as the inspection period. The crews performed a walk-by visual inspection (partial or 360-degree) of trees from ground level to identify obvious defects without using any inspection tools. For each tree, crews assessed the three components of a risk rating: likelihood of failure, likelihood of impact, and severity of consequences [24,25]. From the components, the crews assigned each tree a damage category corresponding to mitigation options. Damage categories included the following: “n/a”—missing tree, “none”—no damage present, “minor”—minimal corrective pruning needed, “significant”—major corrective pruning needed, and “whole tree”—severely damaged crown or whole-tree failure, requiring tree removal.

From the population of assessed trees ( $n = 5094$  individuals;  $n = 38$  species), we excluded any species with fewer than 30 individuals, paring the sample to  $n = 4194$  individuals of 15 species. We used descriptive statistics to describe the proportion of individuals in each species and the range of individual DBHs. We used ordinal logistic regression to investigate whether measured parameters (species, DBH, height, presence of utility lines) influenced the proportion of individuals in each damage category. We considered the presence of utility lines a surrogate for pruning, assuming that individuals near utility lines would, on average, be pruned more frequently than individuals away from utility lines. We computed survivorship as the proportion of each species that remained standing following the hurricane. We used SPSS (v 26.0, IBM Corporation, Armonk, NY, USA) for ordinal regression models and Excel (Microsoft Corporation, Redmond, Washington, DC, USA) to compute odds ratios and descriptive statistics. We made statistical inferences at a 0.05 level of significance.

## 3. Results

Table 1 lists 15 species with at least 30 individuals assessed after Hurricane Irma in Naples, including the number of individuals assessed and each species’ proportion of the total sample. Nearly half of all individuals were *Swietenia mahogany* (L.) Jacq., and more than 70% of the sample

was composed of three species: *S. mahogani*, *Ficus retusa* L., and *Quercus virginiana* Mill. In contrast, individuals of the eight least-common species made up less than 10% of the sample (Table 1).


**Table 1.** Count of individuals within each species assessed prior to the passage of Hurricane Irma in Naples, Florida, USA. Only species with at least 30 individuals are included; species are arranged in descending order of their proportion of the total sample ( $n = 4194$ ).

Species	Individuals Assessed	Proportion of Total Sample	Cumulative Proportion
<i>Swietenia mahogani</i> (L.) Jacq.	1800	42.92%	42.92%
<i>Ficus retusa</i> L.	650	15.50%	58.42%
<i>Quercus virginiana</i> Mill.	536	12.78%	71.20%
<i>Bucida buceras</i> L.	390	9.30%	80.50%
<i>Ilex</i> $\times$ <i>attenuata</i> Ashe	184	4.39%	84.89%
<i>Delonix regia</i> (Bojer ex Hook.) Raf.	133	3.17%	88.06%
<i>Tabebuia heterophylla</i> (DC.) Britton	92	2.19%	90.25%
<i>Ligustrum japonicum</i> Thunb.	74	1.76%	92.01%
<i>Peltophorum pterocarpum</i> (DC.) Backer ex K.Heyne	60	1.43%	93.44%
<i>Taxodium distichum</i> L.	57	1.36%	94.80%
<i>Tabebuia impetiginosa</i> (Mart. ex DC.) Standl.	51	1.22%	96.02%
<i>Bauhinia</i> $\times$ <i>blakeana</i> Dunn	47	1.12%	97.14%
<i>Melaleuca viminalis</i> (Sol. ex Gaertn.) Byrnes.	43	1.03%	98.17%
<i>Cassia fistula</i> L.	43	1.03%	99.20%
<i>Bulnesia arborea</i> (Jacq.) Engl.	34	0.80%	100.00%

Figure 1 shows the distribution of DBH for each species, except *F. retusa*, in Table 1. We did not include *F. retusa* in the latter analyses because of its habit of growing aerial roots, which created uniquely large DBH values. The median DBH for species ranged from 15.2 cm for *Bulnesia arborea* (Jacq.) Engl., *Ilex*  $\times$  *attenuata* Ashe, and *Melaleuca viminalis* (Sol. ex Gaertn.) Byrnes. to 66.0 cm for *S. mahogani*. The interquartile range of DBH was greatest for *S. mahogani* (61.0 cm), nearly twice the range of the species with the next largest value (34.3 cm), *Delonix regia* (Bojer ex Hook.) Raf. Species, with the smallest interquartile range (2.5 cm) found for the species *M. viminalis*, *I. x attenuata*, and *Tabebuia heterophylla* (DC.) Britton.

Of 4194 individuals assessed before Hurricane Irma, 162 were not found during the post-hurricane assessments, and we excluded them from further analysis. Of the remaining 4034 individuals, 74% sustained no damage, 4% sustained only minor damage, 6% sustained significant damage and 15% were whole-tree failures or individuals with sufficient crown damage to justify whole tree removal. Survivorship and the proportion of individuals in each damage category varied among species (Figure 2). More than 90% of individuals of eight species (*Bucida buceras* L., *Cassia fistula* L., *D. regia*, *I. x attenuata*, *Ligustrum japonicum* Thunb., *M. viminalis*, *Q. virginiana*, *Taxodium distichum* L.) survived the hurricane. In contrast, less than 60% of individuals of three species (*B. arborea*, *Bauhinia*  $\times$  *blakeana* Dunn, *T. heterophylla*) survived the hurricane.

Species differences also existed with respect to the proportion of individuals in each damage category (Figure 2). For five species (*D. regia*, *I. x attenuata*, *L. japonicum*, *M. viminalis*, *T. distichum*), more than 90% of individuals sustained no damage. For an additional four species (*B. buceras*, *C. fistula*, *F. retusa*, *Q. virginiana*), more than 80% of individuals sustained no or minor damage. The remaining six species fared worse. For *B. arborea* and *T. heterophylla*, more than 50% of individuals sustained significant damage or whole-tree failure; for the remaining four species (*B. blakeana*, *Peltophorum pterocarpum* (DC.) Backer ex K.Heyne, *S. mahogany*, *Tabebuia impetiginosa* (Mart. ex DC.) Standl.), at least 30% of individuals sustained significant damage or whole-tree failure.



**Figure 1.** Box and whisker plot for stem diameter 1.4 m above ground (DBH) of species listed in Table 1, except *Ficus retusa* L., which was not included because of the presence of aerial roots. Complete binomials are in Table 1. Means represented as square symbols and outliers represented as rhombus symbols.



**Figure 2.** Proportion of each damage type (described in the text) for each species. Numbers at the left edge of each bar indicate the sample size excluding individuals not found in post-hurricane assessments; numbers at the right edge of the chart indicate survivorship, computed as the proportion of trees of each species rated as having no damage, minor damage, or significant damage after the hurricane. Complete binomials are in Table 1.



Table 2 includes the output from the ordinal logistic regression. We excluded *F. retusa* from the model because of its uniquely large DBH values and used *Q. virginiana* as the comparison species because previous studies have shown it to be a storm-resistant species [5,11,12]. Species, DBH, and the presence of utility lines influenced the proportion of individuals in each damage category. For each cm increase in DBH, the odds of an individual moving to a lesser damage category increased by a factor of 1.01. Similarly, individuals near utility lines were 9.09 times more likely to be in a lower damage category than individuals not near a utility line. Compared to *Q. virginiana*, individuals of six species (*B. × blakeana*, *B. arborea*, *P. pterocarpum*, *S. mahogany*, *T. heterophylla*, and *T. impetiginosa*) were significantly more likely to be in a higher damage category, and individuals of two species (*I. × attenuata*, *L. japonicum*) were significantly more likely to be in a lower damage category. There was weaker evidence ( $p = 0.07$ ) that *M. viminalis* and *T. distichum* were more likely to be in a lower damage category than individuals of *Q. virginiana*.

**Table 2.** Ordinal logistic regression output including factors that influenced tree damage in Naples, Florida, USA following Hurricane Irma. Positive coefficients indicate an increase in damage category (e.g., from no damage to minor damage); negative coefficients indicate a decrease in damage category (e.g., from significant damage to minor damage). *Ficus retusa* L. was not included in the model because of uniquely large DBH values; sample sizes for the remaining species are in Figure 2. Complete binomials are in Table 1.

Factor	Coefficient	Standard Error	<i>p</i> -Value <sup>z</sup>	Odds-Ratio <sup>y</sup>	95% CI Lower	95% CI Upper
Species— <i>B. × blakeana</i>	1.86	0.33	<0.00	6.43	3.35	12.26
Species— <i>B. buceras</i>	−0.09	0.20	0.64	1.10	1.63	1.34
Species— <i>B. arborea</i>	2.21	0.43	<0.00	9.10	3.97	21.42
Species— <i>C. fistula</i>	0.27	0.43	0.54	1.31	0.52	2.88
Species— <i>D. regia</i>	−0.56	0.35	0.11	1.75	3.70	1.09
Species— <i>I. × attenuata</i>	−0.63	0.29	0.03	1.89	3.45	1.08
Species— <i>L. japonicum</i>	−2.51	1.01	0.01	12.50	219.72	2.70
Species— <i>M. viminalis</i>	−1.34	0.74	0.07	3.85	25.00	1.14
Species— <i>P. pterocarpum</i>	1.27	0.30	<0.00	3.54	1.93	6.38
Species— <i>S. mahogani</i>	1.37	0.14	<0.00	3.94	3.00	5.22
Species— <i>T. heterophylla</i>	1.19	0.32	<0.00	3.30	1.72	6.12
Species— <i>T. impetiginosa</i>	2.07	0.25	<0.00	7.95	4.91	12.95
Species— <i>T. distichum</i>	−1.10	0.61	0.07	3.03	12.5	1.08
DBH	−0.01	<0.00	<0.00	1.01	1.02	1.01
Utility Line	−2.19	0.40	<0.00	9.09	20.00	4.16

<sup>z</sup> *p*-Values for species indicate whether the proportion of individuals in each damage category for a species differed from the baseline species (*Quercus virginiana* Mill.). <sup>y</sup> The odds-ratio indicates the magnitude of the change in likelihood of being in a higher (for positive coefficients) or lower (for negative coefficients) damage category (i) between a species and the baseline species, (ii) for a 1 cm increase in DBH, or (iii) if utility lines were present.

#### 4. Discussion

Hurricanes and other catastrophic wind events will continue to threaten urban forests. Because most climate models predict the increasing frequency and severity of such storms—especially in coastal regions like the Atlantic and Gulf Coasts in the United States—the economic and human costs associated with storm-induced tree failures will likely increase in the future. At the same time, however, communities and homeowners continue to plant trees to capitalize on the many benefits they provide to residents. Understanding the factors that influence the likelihood of failure and the degree of damage sustained by amenity trees is integral to sustainable communities.


Our work considered two intrinsic factors that influence the likelihood of tree failure (species and size), as well as the presence of utility lines, which we assumed reflected the effect of pruning. However, we did not consider the effect of structural defects, adding uncertainty to our findings, even though presence/absence was not a factor in [9]; nor did we investigate the effect of extrinsic factors such as wind speed. A careful spatial analysis would help to correlate local wind speed data with the likelihood of failure or damage, but a more comprehensive sample will be necessary to include other relevant factors like species, size, previous maintenance, and presence/severity of structural defects.



We expected species to influence the proportion of individuals in each damage category because many previous studies have shown differences in survivorship between species in forests and plantations [1,16,20,33,34], as well as urban forests [3,4,6,7,10,12]. Our data include open-grown individuals of eight species (*B. arborea*, *B. × blakeana*, *C. fistula*, *F. retusa*, *I. × attenuata*, *L. japonicum*, *P. pterocarpum*, *T. impetiginosa*) not previously studied, and seven species (*B. buceras*, *D. regia*, *M. viminalis*, *Q. virginiana*, *S. mahogani*, *T. distichum*, *T. heterophylla*) included in previous studies [5,11,12]. Consistent with previous regional studies of open-grown trees [5,7,11,12], our data revealed species differences, although we cannot rule out the possibility that such differences were the result of differences in unmeasured variables, such as the presence of defects.

Categorizing individuals in terms of damage—in addition to survivorship—was important, because it may help communities prepare not only for the proportion of trees that may be removed following a storm, but also the volume of debris that needs to be managed. Measuring survivorship alone will likely underestimate the volume of debris that accumulates from individuals that sustained significant damage but remained standing and would not be removed following a storm.

Despite this limitation, we computed survivorship to more readily compare our data with previous studies that reported survivorship of open-grown species from seven previous hurricanes in the region [5,11,12]. Figure 3 shows survivorship by species for all eight hurricanes, ranging in sustained wind speed from 122 km/h to 265 km/h. Generally, there is a decrease in survivorship in the reported species as wind speed increases. For example, the survivorship of *B. buceras* was noticeably higher following Hurricane Irma (122 km/h) than Hurricane Charley (233 km/h). A similar overall trend can be noted among the seven data sets featuring *Pinus elliotii* Engelm. In light of this, variability among species within each hurricane and within a species among hurricanes is evident. For example, the survivorship of *T. heterophylla* was noticeably lower following Hurricane Irma (122 km/h) than hurricanes Andrew (265 km/h) or Georges (188 km/h). The survivorship of *Q. virginiana*, with a reputation among arborists and urban foresters as a particularly resilient species, was high throughout the range of wind speeds. Similarly, *T. distichum* survivorship was nearly identical for Hurricanes Irma and Charley, despite their differing intensities. Such disparities underscore the inherent variability of survivorship, which depends on many intrinsic and extrinsic factors.



**Figure 3.** Scatter plot of survivorship of open-grown individuals vs. wind speed for 16 species and eight hurricanes in Florida, USA. Data from the current study (15 species) align with 122 km/h (average sustained wind across six measurement points [30]); the remaining data come from [5,11,12]. Note that the axes do not intersect at zero. Complete binomials are in Table 1, except for *Pinus elliotii* Engelm.

Inherent variability was also evident with the effect of DBH on the proportion of individuals in each damage category. Previous studies from the region [5,7,11] and of other sub-tropical [6] and temperate climates [1,3] have shown that the likelihood of failure or the severity of damage is greater for larger trees. Following Hurricane Irma, however, larger trees in Naples were less likely to have sustained greater damage. It was unclear why this occurred, although it was possible that there was a confounding interaction between species and DBH. It is also possible that the effect was an artifact of the damage rating system: the relative severity of a given limb falling decreased with increased crown size and complexity. Although the effect of DBH was statistically significant, the odds ratio was comparatively small. Applying the odds ratio in practice, a tree would have to be 75 cm smaller in DBH to have twice the likelihood of being in the next highest damage category.

Consistent with a previous study in Florida which indicated that pruned trees were less likely to sustain damage [5], trees near utility lines in Naples were less likely to be in a more severe damage category. We assumed that trees near utility lines were more likely to have been pruned than trees not in proximity to utility lines, which would remove defective branches from parts of the crown near the lines. Repeated pruning may also have reduced the crown area which, in turn, would reduce the wind-induced bending moment [19].

## 5. Conclusions

Our findings contribute to a growing body of knowledge that describes the likelihood of failure and the severity of damage to open-grown trees in subtropical climates. However, even in a hurricane-prone region such as Florida, which has provided many opportunities to study wind-induced tree failure, a more comprehensive meta-analysis would be necessary to investigate the interactions between intrinsic (size, species, presence of defects, tree maintenance history, and maintenance of infrastructure like sidewalks and roads that often damages roots and induces windthrow) and extrinsic (wind speed, sheltering, soil texture and volume) factors that influence wind-induced tree failures and damage. Such work is challenging, however. Unlike forest or plantation settings, wind-induced damage in residential settings must be cleaned up relatively quickly, often precluding the collection of detailed measurements of individual trees.

Species selection is an important aspect of urban forest management; in communities that experience regular and severe storms, arborists and urban foresters should consider which species have a greater likelihood of failure or damage. However, restricting selection to species that have consistently demonstrated storm resilience would unnecessarily limit species diversity. Maintaining a resilient and diverse urban forest will require additional knowledge of species performance in storms. It will also require careful planning to provide shelter for less-resilient species and sound arboricultural practice to reduce wind loads and minimize the likelihood of defects that compromise structural integrity.

**Author Contributions:** Conceptualization, B.K., R.W.K., A.K.K., S.L., S.M.L., and H.S.; methodology, S.L. and H.S.; validation, B.K. and A.K.K.; formal analysis, R.W.K. and A.K.K.; resources, S.L. and H.S.; data curation, R.W.K., A.K.K., S.L., and H.S.; writing—original draft preparation, B.K., R.W.K., and A.K.K.; writing—review and editing, G.H., S.L., S.M.L. and H.S.; visualization, B.K. and R.W.K.; supervision, G.H. and A.K.K.; project administration, A.K.K., S.L., and H.S.; funding acquisition, A.K.K., S.L., and S.M.L. All authors have read and agreed to the published version of the manuscript.

**Funding:** The data collection for this research was funded by the Southern Group of State Foresters. This paper is part of a larger effort to assemble and analyze urban forestry storm data funded by the USDA Forest Service National Urban and Community Forestry Advisory Council (NUCFAC). APC was funded by the University of Florida Library System.

**Acknowledgments:** We would like to thank Gregory Dunn, Arthur Clothier, and Clark Ryals for their efforts in performing the post-storm inspections. All three serve as County Foresters for the Florida Forest Service and are International Society of Arboriculture-Certified Arborists.

**Conflicts of Interest:** The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

## References

1. Everham, E.M.; Brokaw, N.V.L. Forest damage and recovery from catastrophic wind. *Bot. Rev.* **1996**, *62*, 113–185. [\[CrossRef\]](#)
2. MacFarlane, D.W.; Kane, B. Neighbour effects on tree architecture: Functional trade-offs balancing crown competitiveness with wind resistance. *Funct. Ecol.* **2017**, *31*, 1624–1636. [\[CrossRef\]](#)
3. Gibbs, J.N.; Greig, B.J.W. Survey of parkland trees after the great storm of October 16, 1987. *Arboric. J.* **1990**, *14*, 321–347. [\[CrossRef\]](#)
4. Francis, J.K.; Gillespie, A.J.R. Relating gust speed to tree damage in Hurricane Hugo, 1989. *J. Arboric.* **1993**, *19*, 368–373.
5. Duryea, M.L.; Blakeslee, G.M.; Hubbard, W.G.; Vasquez, R.A. Wind and trees: A survey of homeowners after Hurricane Andrew. *J. Arboric.* **1996**, *22*, 44–50.
6. Jim, C.Y.; Liu, H.T. Storm damage on urban trees in Guangzhou, China. *Landsc. Urban Plan.* **1997**, *38*, 45–59. [\[CrossRef\]](#)
7. Francis, J.K. Comparison of hurricane damage to several species of urban trees in San Juan, Puerto Rico. *J. Arboric.* **2000**, *26*, 189–197.
8. Escobedo, F.J.; Luley, C.J.; Bond, J.; Staudhammer, C.; Bartel, C. Hurricane debris and damage assessment for Florida urban forests. *J. Arboric.* **2009**, *35*, 100–106.
9. Kane, B. Tree failure following a windstorm in Brewster, Massachusetts, USA. *Urban For. Urban Green.* **2008**, *7*, 15–23. [\[CrossRef\]](#)
10. Mayer, H.; Trump, T.; Escobedo, F. Tree damage and restoration following Hurricanes Katrina and Wilma at Miami Metrozoo. *Proc. Fla. State Hort. Soc.* **2007**, *120*, 320–322.
11. Duryea, M.L.; Kampf, E.; Littell, R.C. Hurricanes and the urban forest: I. Effects on Southeastern United States coastal plain tree species. *Arboric. Urban For.* **2007**, *33*, 83–97.
12. Duryea, M.L.; Kampf, E.; Littell, R.C.; Rodriguez-Pedraza, C.D. Hurricanes and the urban forest: II. Effects on tropical and subtropical tree species. *Arboric. Urban For.* **2007**, *33*, 98–112.
13. Hull, R.B. How the public values urban forests. *J. Arboric.* **1992**, *18*, 98–101.
14. Foster, D.R. Disturbance history, community organization and vegetation dynamics of the old-growth Pisgah Forest, South-Western New Hampshire, U.S.A. *J. Ecol.* **1988**, *76*, 105–134. [\[CrossRef\]](#)
15. Webb, S.L. Contrasting windstorm consequences in two forests, Itasca State Park, Minnesota. *Ecology* **1989**, *70*, 1167–1180. [\[CrossRef\]](#)
16. Walker, L.R. Tree damage and recovery from Hurricane Hugo in Luquillo Experimental Forest, Puerto Rico. *Biotropica* **1991**, *23*, 379–385. [\[CrossRef\]](#)
17. Putz, F.E.; Coley, P.D.; Lu, K.; Montalvo, A.; Aiello, A. Uprooting and snapping of trees: Structural determinants and ecological consequences. *Can. J. For. Res.* **1983**, *13*, 1011–1020. [\[CrossRef\]](#)
18. Zimmerman, J.K.; Everham, E.M.; Waide, R.B.; Lodge, D.J.; Taylor, C.M.; Brokaw, N.V.L. Responses of tree species to hurricane winds in subtropical wet forest in Puerto Rico: Implications for tropical tree life histories. *J. Ecol.* **1994**, *82*, 911–922. [\[CrossRef\]](#)
19. Smiley, E.T.; Kane, B. The effects of pruning type on wind loading of *Acer rubrum*. *Arboric. Urban For.* **2006**, *32*, 33–39.
20. Putz, F.E.; Sharitz, R.R. Hurricane damage to old-growth forest in Congaree Swamp National Monument, South Carolina, U.S.A. *Can. J. For. Res.* **1991**, *21*, 1765–1770. [\[CrossRef\]](#)
21. Kennard, D.K.; Putz, F.E.; Niederhofer, M. The predictability of tree decay based on visual assessments. *J. Arboric.* **1996**, *22*, 249–254.
22. Pokorny, J.D. *Urban Tree Risk Management, A Community Guide to Program Design and Implementation*; USDA Forest Service Northeastern Area State and Private Forestry: St. Paul, MN, USA, 2003; 19p.
23. Fink, S. Hazard tree identification by visual tree assessment (VTA): Scientifically solid and practically approved. *Arboric. J.* **2009**, *32*, 139–155. [\[CrossRef\]](#)
24. Dunster, J.A.; Smiley, E.T.; Matheny, N.; Lilly, S. *Tree Risk Assessment Manual*, 2nd ed.; International Society of Arboriculture: Champaign, IL, USA, 2017; 194p.
25. Smiley, E.T.; Matheny, N.; Lilly, S. *Best Management Practices: Tree Risk Assessment*, 2nd ed.; International Society of Arboriculture: Champaign, IL, USA, 2017; 86p.

26. Ostertag, R.; Silver, W.L.; Lugo, A.E. Factors affecting mortality and resistance to damage following hurricanes in a rehabilitated subtropical moist forest. *Biotropica* **2005**, *37*, 16–24. [[CrossRef](#)]
27. Wiersma, Y.F.; Davis, T.L.; Eberendu, E.C.; Gidge, I.; Jewison, M.; Martin, H.C.; Parsons, K.C.; Patterson, H.; Quirke, A. Hurricane Igor impacts at northern latitudes: Factors influencing tree fall in an urban setting. *Arboric. Urban For.* **2012**, *38*, 92–98.
28. Edberg, R.; Berry, A. Patterns of structural failures in urban trees: Coast live oak (*Quercus agrifolia*). *J. Arboric.* **1999**, *25*, 48–55.
29. Edberg, R.J.; Berry, A.M.; Costello, L.R. Patterns of structural failure in Monterey pine. *J. Arboric.* **1994**, *20*, 297–304.
30. Hurricane Irma Local Report/Summary. Available online: <https://www.weather.gov/mfl/hurricaneirma> (accessed on 21 April 2020).
31. Saffir-Simpson Hurricane Wind Scale. Available online: <https://www.nhc.noaa.gov/aboutsshws.php> (accessed on 31 October 2019).
32. Urban Forest Strike Team (UFST). *Urban Forest Strike Team Post-Disaster Rapid Tree Risk Assessment Specification*; Version 7.0; 2018; Unpublished.
33. Frangi, J.L.; Lugo, A.E. Hurricane damage to a flood plain forest in the Luquillo Mountains of Puerto Rico. *Biotropica* **1991**, *24*, 324–335. [[CrossRef](#)]
34. Gresham, C.A.; William, T.M.; Lipscombe, D.J. Hurricane Hugo wind damage to Southeastern U.S. coastal forest tree species. *Biotropica* **1991**, *23*, 420–426. [[CrossRef](#)]



© 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).