1-1-1977

Shifts in perception of size after adaptation gratings.

Francine Sara Frome
University of Massachusetts Amherst

Follow this and additional works at: https://scholarworks.umass.edu/dissertations_1

Recommended Citation

Frome, Francine Sara, "Shifts in perception of size after adaptation gratings." (1977). *Doctoral Dissertations 1896 - February 2014*. 2770.
<https://doi.org/10.7275/t7r5-rh05> https://scholarworks.umass.edu/dissertations_1/2770

This Open Access Dissertation is brought to you for free and open access by ScholarWorks@UMass Amherst. It has been accepted for inclusion in Doctoral Dissertations 1896 - February 2014 by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

UMASS/AMHERST

312066013585426

SHIFTS IN PERCEPTION OF SIZE
AFTER ADAPTATION GRATINGS. II.

A Dissertation Presented

By

FRANCINE SARA FROME

Submitted to the Graduate School of the
University of Massachusetts in partial fulfillment
of the requirements for the degree of

DOCTOR OF PHILOSOPHY

May 1977

Psychology Department

(c) Francine Sara Frome 1977

All Rights Reserved

SHIFTS IN PERCEPTION OF SIZE
AFTER ADAPTATION GRATINGS. II.

A Dissertation Presented

By

FRANCINE SARA FROME

Approved as to style and content by:

William Eichelman, Chairperson

Arnold Well, Member

Edward Riseman, Member

Bonnie Strickland, Chairperson
Psychology Department

ABSTRACT

Shifts in Perception of Size
after Adaptation Gratings. II.

June 1977

Francine Sara Frome, B.A., University of Maryland
M.S., University of Massachusetts
Ph.D., University of Massachusetts

Directed by: Professor William Eichelman

Frome et al. (1974) have shown that after adaptation to vertical sinusoidal gratings, a rectangle looks wider, regardless of whether an adapting grating bar was smaller than, larger than, or equal to the size of the test rectangle. The direction of these size after-effects is very different from those found by Blakemore et al. (1907) when they used sinusoidal gratings as both their test and adapting stimuli and found shifts in perceived frequency away from the adapting frequency. Four experiments tested possible reasons for the differences in the direction of these after-effects. The first experiment examined the possibility that the differences could be due to the judgments being made.

Experiment 1 compared judgments of size with judgments of spatial frequency. A short grating of multiple filled rectangles was used as the test stimulus for size or spatial frequency judgments, both pre- and post-adaptation to a grating. When observers judged spatial frequency after adaptation, they all obtained the perceived spatial frequency

shift with the short rectangle grating as test stimulus, just as Blakemore and Sutton did in their demonstration (1969) using tall gratings. However, when observers made size judgments by adjusting the height of the bars in the short grating until all the bars appeared square before and after adaptation to a vertical sinusoidal grating, the multiple test rectangles appeared wider after adaptation, regardless of whether the bars in the adapting grating were smaller than, larger than, or equal to the width of the squares in the short grating. This is the same resulting size after-effect obtained by Frome (1974) when she used a single rectangle as test stimulus. These results showed striking dissociations between judgments of perceived spatial frequency and judgments of perceived size: adapting to a slightly lower frequency grating, increased the perceived frequency shift, while an increase in the perceived width of the multiple squares was still observed. These results led to the conclusion that for supra-threshold judgments like these, "width" and "spatial frequency" are not equivalent. The possibility that subjects attend only to a single rectangle in the short grating during their size judgment was discussed.

Experiment 2 explored the contribution of apparent duty cycle to the perception of size or spatial frequency of a multiple square grating. The effect of adaptation on the perceived duty cycle of a multiple square grating could have

accounted for both the size and spatial frequency results of Experiment 1 if there were differential adaptation effects on duty cycle across the combinations of adapting and test grating frequencies that were used. However, these results were not obtained. For nearly all period ratios, after adaptation to a vertical sinusoid, the rectangles in the multiple perceived squares grating became perceptually narrower than the spaces. These after-effects were often small and not significant. Therefore, the effects of adaptation on apparent duty cycle cannot account for the dissociation of size and spatial frequency found in Experiment 1.

Experiment 3 explored the possibility that the size after-effects with square test stimuli were different than those obtained with sinusoidal stimuli because the effective stimulus width of the bars of the sinusoidal adapting grating may have been psychologically smaller than they were physically since contours of sinusoidal bars are indistinct. Subjects adjusted the height of a rectangle until it appeared square before and after adaptation to either a sine wave or a square wave grating. After adaptation to both sine wave and square wave gratings with vertically oriented bars, the rectangle appeared wider for all width ratios. Sine wave and square wave effects were nearly identical. That square wave adaptation produced identical effects to sine wave adaptation showed that the sine wave after-effect of only apparent widening of the single rectangle could not

have been due to the bars having a narrower psychological than physical width because of their blurry edges.

Experiment 4 introduced an outline rectangle as a test target. Subjects adjusted the height of the rectangle before and after adaptation to a vertical sinusoidal grating, as with the other test targets; yet this stimulus has a very different frequency spectrum than the others. After adaptation to a vertical sinusoidal grating, the outline rectangle always appeared wider, regardless of whether the bars in the grating were smaller than, larger than, or equal to the rectangle width. These results were essentially the same as those with a filled rectangle as test stimulus and a simple neural size channels hypothesis predicts neither result. Frome et al. (1974) have shown that the size after-effects obtained with a single filled rectangle can be explained by the operation of spatial frequency specific channels. Further psychophysical measurements or ad hoc assumptions must be made if the data from the outline rectangle is to be explained in terms of the distribution of activity in the spatial frequency specific channels. In addition, a new sort of size model was called for, since similar size ratios of adapting grating to test stimulus width yield similar results, regardless of the spectral composition of the test target.

TABLE OF CONTENTS

Introduction	.1
Experiment 1	22
Experiment 2	57
Experiment 3	73
Experiment 4	93
References	.115
Appendix A: Terms describing a grating.	.123

LIST OF TABLES

1. Adapting grating, and test rectangle sizes used to obtain P_A/P_T conditions. 41
2. Multiple filled squares: t tests for the mean of the ratios ($n=11$) of the $\bar{v}$ vertical extent needed for squareness in the adaptation condition to the mean vertical extent needed for squareness in the control condition 43
3. Multiple filled squares - naive University of Maryland subjects: t tests for the mean of the ratios ($n=11$) of the $\bar{v}$ vertical extent needed for squareness in the adaptation condition to the mean vertical extent needed for squareness in the control condition 49
4. Perceived multiple squares duty cycle: t tests for the mean of the ratios ($n=11$) of the $\bar{h}$ horizontal extent of the dim spaces needed for perceived 50% duty cycle in the adaptation condition to the mean horizontal extent of the dim spaces needed for perceived 50% duty cycle in the control condition. 69
5. Sine wave versus square wave adaptation: t tests for the mean of the ratios ($n=11$) of the vertical extent needed for squareness in the adaptation condition to the mean vertical extent needed for squareness in the control condition 89
6. Outline Square: t tests for the mean of the ratios ($n=11$) of the vertical extent needed for squareness in the adaptation condition to the mean vertical extent needed for squareness in the control condition 107

LIST OF FIGURES

1. Two parallel descriptions of a sine wave grating. The left part of the figure specifies the grating as amplitude versus position. The right part of the figure specifies the grating as amplitude versus frequency 3
2. Figure 2 illustrates many terms used in the description of a grating 5
3. The modulation transfer function (MTF) for a typical observer. The MTF is a measure of contrast sensitivity of the human visual system to sinusoidal grating as a function of their spatial frequency 8
4. Pattern for demonstration of the perceived spatial shift. (After Blakemore and Sutton, 1969. Copyright 1969 by the American Association for the Advancement of Science.) 16
5. Pattern for demonstration of the perceived spatial frequency shift with a short grating as test stimulus 23
6. An individual subject's data from Experiment 1, which had multiple filled squares, that is a short grating, as test object. The abscissa is the ratio of adapting bar width (P_A) to test rectangle widths (P_T). Each ordinate is the mean of the ratios of the vertical extent needed for perceived squareness in the experimental condition (E), after grating adaptation, to the mean vertical extent needed for squareness in the control condition before adaptation ($\bar{C}$). Standard errors of the mean are plotted for each datum. The left side of the figure illustrates results obtained when adapting with horizontally oriented grating bars, and the right side illustrates results when vertically oriented adapting grating bars were used. These same coordinate axes are used for all other data in Experiment 1, Figures 7-11 29

7.	An individual subject's data from Experiment 1 . . .	31
8.	An individual subject's data from Experiment 1 . . .	33
9.	An individual subject's data from Experiment 1 . . .	35
10.	An individual subject's data from Experiment 1 . . .	37
11.	An individual subject's data from Experiment 1 . . .	39
12.	Seven naive subjects' data for a P_A/P_T ratio of 1.5 in Experiment 1	47
13.	Quantitative results of Blakemore <u>et al.</u> (1970) obtained with vertical sine wave gratings as both adapting and test stimuli which have been replotted on the same coordinate axes as the data in the present experiments.	51
14.	The one-dimensional spectral distributions of a single filled square and of multiple filled squares	55
15.	An individual subject's data from Experiment 2. Subjects set a perceived multiple square grat- ing to 50% duty cycle (equal bar and space width) both before and after adaptation to a vertical sine wave grating. The abscissa is the ratio of adapting grating period (P_A) to short test grating period (P_T). Each ordinate is the mean of the ratios of the horizontal extent of the dim spaces needed for perceived 50% duty cycle of the perceived multiple square grating in the experimental condition (E), after grating adaptation, to the mean horizon- tal extent of the dim spaces needed for perceived 50% duty cycle in the control condition before adaptation ($\bar{C}$). Standard errors of the mean are plotted for each datum. These same coordinate axes are used for all other data in Experiment 2, Figures 16 and 17.	60
16.	An individual subject's data from Experiment 2 . . .	62
17.	An individual subject's data from Experiment 2 . . .	64

18. An illustration and explanation of subjects' setting and perception of a 50% duty cycle grating before and after adaptation to a vertical grating 67
19. The one-dimensional spectral distribution of a 7.5 min arc filled rectangle. The frequency spectrum may be described by the function $\sin x/x$. The open arrows indicate the fundamental frequencies of the adapting gratings that were used in Experiment 3. The filled arrows indicate the frequencies of the third harmonics of the square waves that were used 75
20. An individual subject's data from Experiment 3, which compares results from vertical square wave and sine wave adapting gratings. Standard errors of the mean are plotted for each datum. The coordinate axes used for all data in Experiment 3, Figures 20-24, are the same as those used in Experiment 1. 79
21. An individual subject's data from Experiment 3 . . . 81
22. An individual subject's data from Experiment 3 . . . 83
23. An individual subject's data from Experiment 3 . . . 85
24. The average of all subjects' data from Experiment 3, with the standard error of the mean between subjects indicated for each datum. 87
25. The one-dimensional spectral distributions of 5' and 10' wide test stimuli: a single filled square, multiple filled squares, and an outline square. As the legend shows, arrows indicate different ratios of adapting grating bar width to square width. 94
26. An individual subject's data from Experiment 4, which uses an outline square as test object before and after adaptation to a vertical sine wave grating. Standard errors of the mean are plotted for each datum. The coordinate axes used for all data in Experiment 4, Figures 26-30, are the same as those used in Experiments 1 and 3. 97

27. An individual subject's data from Experiment 4. . . 99
28. An individual subject's data from Experiment 4. . . 101
29. An individual subject's data from Experiment 4. . . 103
30. The average of all subjects' data in Experiment 4, with the standard error of the mean between subjects indicated for each datum.105
31. The average of all subjects' data when a single filled square was used as test object before and after adaptation to a vertical sinusoidal grating (After Frome et al., 1974). The standard error of the mean between subjects is indicated for each datum. The data have been plotted on the same coordinate axes as the data in the present experiments.110

INTRODUCTION

Until recently the discussion of psychological factors involved in human perception of spatial dimensions has had rather limited scope and has lacked general theoretical structure (Graham, 1966). In order to be comprehensive such a discussion should include: 1) a quantifiable description of all possible stimuli with a set of physically measurable variables; 2) a demonstrated underlying neurophysiological substrate; and 3) a proposed operation of physiological mechanisms which has predictive value for human perception (Cornsweet, 1970).

A recent approach to the specification of a visual pattern has been to apply a specific kind of linear systems analysis, Fourier analysis, to the stimulus. Fourier's theorem states that any waveform (in this case, the spatial light intensity distribution) can be completely described as the sum of sinusoidal waves of appropriate frequency, amplitude and phase. This theorem states that there are two parallel domains such that specification of a function in one domain (one domain specified as amplitude versus position) completely specifies the corresponding function in the parallel domain (the parallel domain specified as amplitude versus frequency). A sine wave grating (a repetitive pattern of blurry light and dark bars which has a luminance

distribution that varies sinusoidally) may be considered a primitive in the Fourier analysis of pattern, since it contains only one frequency, and in only one dimension. The two parallel descriptions of a sine wave grating may be seen in Figure 1.

Fourier synthesis is the procedure for finding a waveform that is produced when a set of sine waves are added together. Fourier analysis is the procedure for finding the particular set of sine waves that must be added in order to obtain some given waveform. Those resultant sine waves are called the Fourier components of the given waveform. The definitions that are necessary to completely describe a grating may be found in the appendix. Figure 2 illustrates many of these terms.

In order to use gratings to describe the transmission of spatial information through the visual system, Campbell and Robson (1968) presented a particular spatial frequency grating at a fixed average luminance to an observer, and adjusted the grating's contrast until the observer reported just seeing the grating. This is the observer's contrast threshold for that grating. The reciprocal of the contrast threshold is the sensitivity of the observer to the grating. The measure of contrast sensitivity of the human visual system to sinusoidal gratings as a function of their spatial

Figure 1. Two parallel descriptions of a sine wave grating. The left part of the figure specifies the grating as amplitude versus position. The right part of the figure specifies the grating as amplitude versus frequency.

Figure 2. Figure 2 illustrates many terms used in the description of a grating.

frequencies defines the modulation transfer function (MTF) for an observer. A typical MTF is plotted in Figure 3. A typical observer might be most sensitive to a grating of about 4 c/d with less sensitivity to both higher and lower spatial frequencies. The limit of resolution for the visual system is typically a grating with bars of 1 minute of arc width. Thus, a frequency analysis approach has met the first criterion of a comprehensive theory of perception of spatial dimensions--it has described the physical stimulus by showing how effectively any light intensity distribution passes through the human visual system. But do these physical parameters have a basis in neurophysiology?

Neurophysiological studies in animals have found individual neurons in their visual systems which are selectively sensitive to the orientation and/or dimensions of retinal images. Lettvin et al. (1959) recorded from the frog's optic tectum and found neurons which responded rapidly when small "bug" shapes are moved across its retina. Hubel and Wiesel (1962, 1965, 1968) identified neurons in the visual cortex of the cat which are sensitive to the orientation of targets like bright slits of light, or black-white edges. Campbell, Cooper, and Entroth-Cugell (1969) also recorded from single cells in the visual cortex of the cat, but they used moving grating patterns of variable spatial frequency

Figure 3. The modulation transfer function (MTF) for a typical observer. The MTF is a measure of contrast sensitivity of a human visual system to sinusoidal grating as a function of their spatial frequency.

as stimuli. It is clear from their work that a cat's cortex has cells that respond to the orientation and spatial frequency of grating stimuli, as well as to a single bar or line. Campbell, Cooper, Robson, and Sachs (1969) have also shown such neural units in the monkey visual cortex.

The evidence that the human visual system has cells like these is inferred from psychophysical experiments. Campbell and Robson (1968) originally proposed the visual frequency channels hypothesis that the human visual system contains independent channels which each respond selectively to a particular limited range of spatial frequencies. This hypothesis suggests that the pattern of excited frequency selective neurons constitutes a frequency code of the spatial content of the visual world. If this hypothesis is correct, then human observations should be differentially sensitive to the frequency components in a visual stimulus. Campbell and Robson (1968) showed that the threshold for detection of a square wave grating was predictable from the threshold of its fundamental Fourier component (which has an amplitude 1.27 times that of a sine wave grating of the same fundamental frequency and contrast). They also found that the threshold for discriminating a square wave from a sine wave grating was predicted from the independent threshold of the third harmonic of the square wave. A system which re-

sponds to bars on the basis of their size cannot account for the effects upon the third harmonic observed in Campbell and Robson's (1968) experiments, whereas the effects are easily explained if the frequency components are detected independently.

A basic psychophysical technique in studies testing the frequency and orientation specific channels hypothesis employs a sensory adaptation phenomenon. Blakemore and Campbell (1969a) found that prolonged inspection of a grating increases the contrast threshold for gratings of similar orientation. Blakemore and Campbell (1969b) also found that this long grating inspection will also cause a decrease in the amplitude of the occipital potential evoked by an oscillating grating of nearly the same orientation and spatial frequency as the adapting grating. These and other findings suggest that, as in other animals, there are human frequency and orientation specific neurons and that constant exposure to a grating decreases the sensitivity of those neural channels which would normally respond to that stimulus. This phenomenon of sensory adaptation to gratings is supported neurophysiologically by the work of Maffei, Fiorentini and Bisti (1973). They found when recording from simple cells of the cat striate cortex that exposure to drifting gratings of high contrast consequently reduced the response of the

cells to low contrast gratings.

Adaptation to a grating will not produce any decrease of human contrast sensitivity in frequencies more than an octave away (Pantle & Sekular, 1968; Blakemore & Campbell, 1969). This result is consistent with the notion that there are multiple independent channels tuned to different bands of spatial frequency. The trichromatic theory of color vision is also a separate channels hypothesis. Each of three cone pigments is maximally sensitive to a different wavelength of light and the interactions of the signals from these channels produce the perception of all color. Graham and Nachmias (1971) tested the multiple channels hypothesis as it applies to spatial frequency channels. According to this hypothesis if two spatial frequencies in a stimulus are very different, they should be simultaneously processed by different frequency channels. Therefore, detection threshold should occur when any of the components reaches its own independent threshold, regardless of the phase relation of the components. A single channel neural processing model predicts that the phase relationship of the frequency components that yields the largest amplitude should be the most detectable. The data show that this is not the case, upholding the multiple channels model prediction that detectability was not affected by the phase relations of the

components. Regardless of the phase relation of the component parts, detection threshold is predicted by the component independent frequency threshold. However, a study by Tolhurst (1972) indicates that above threshold, the frequency specific channels are not acting independently. If the fundamental frequency, f , and third harmonic, $3f$, of a supra-threshold square wave adapting grating were treated independently by the visual system, then the threshold elevation of sine wave frequency $3f$ (that of the third harmonic in the square wave adapting grating) should be the same as the threshold elevation caused by adaptation to a single sine wave of frequency $3f$, assuming the amplitudes of the single and component $3f$ frequency were equated. Tolhurst (1972) found this not to be the case, in that the $3f$ frequency caused less threshold elevation as a component in a complex grating than the $3f$ frequency presented alone. Thus, although the work of Graham and Nachmias (1971) indicates independent multiple frequency channels at threshold; above threshold, frequency channels seem to interact.

The studies reported above use detection of pattern as the criterion response, but what a stimulus looks like is a different criterion. Although Graham and Nachmias (1971) found that detection threshold is independent of the phase relation of component parts, Nachmias and Weber (1975) have

shown that well above threshold (4 times), discrimination is almost unlimited for stimuli with the exact same frequency components and that differ only in component phase relations.

One may use a different model, a multiple channels hypothesis which assumes size-specific, rather than frequency specific neurons to make predictions about both detection and appearance of spatial stimuli. A size specific model is supported by evidence like that of Bagrash (1973) who found that adaptation to a luminous disc of a given diameter reduces sensitivity to disks of similar diameter much more than to larger or smaller disks. In contrast to this finding, Sullivan, Georgeson and Oatley (1972) found that adaptation to bars raised the threshold for bars of all widths and for all frequency sine wave gratings. These authors also found that adaptation to 5.5 c/d raised the thresholds of only those sine waves near that frequency, but raised the threshold for all bar widths. This result can be explained by the spatial frequency hypothesis. Since all bars have a continuous multiple frequency spectrum, with some energy in the most sensitive, but in this case, most adapted part of the visual system, the detection threshold for all width bars is raised. The differential effects of adapting to bars and to sinusoidal gratings led these authors to maintain that for contrast sensitivity functions the stimulus property "width"

is not equivalent to the stimulus property "spatial frequency."

One must also ask if the stimulus properties of width and spatial frequency are equivalent with regards to human perception of size. Blakemore and Sutton (1969) and Blakemore, Nachmias, and Sutton (1970) have proposed that frequency selective neural channels are the basic units used to encode the spatial dimensions of suprathreshold stimuli.

These authors have shown that under certain conditions, adaptation to a grating pattern leads to small shifts in the frequencies of other gratings. Figure 4 is the Blakemore and Sutton (1969) demonstration. After adapting for at least one minute by scanning back and forth between the patterns on the right, when one fixates between the two parallel equal-frequency gratings on the left, they no longer appear equal in frequency. Adapting to the low frequency grating, on the lower right, makes the grating to its left appear to rise in frequency, and conversely, for the high frequency adaptation grating.

Blakemore, Nachmias, and Sutton (1970) quantified these perceived frequency shifts. In their experiments, the subject's task was to adjust the frequency of a grating occupying a retinal area previously exposed to an adapting grating. Both the demonstration and experiments were clear in their

Figure 4. Pattern for demonstration of the perceived spatial shift. (After Blakemore and Sutton, 1969. Copyright 1969 by the American Association for the Advancement of Science.).

result--adaptation tends to make gratings appear more different from the adapting frequency than they really are. Their explanation is as follows (from Blakemore, Nachmias, & Sutton, 1970, p. 729):

When a grating of suprathreshold contrast is viewed, it activates frequency-selective neurones in proportion to their sensitivity to the grating's frequency. Imagine that the brain distinguishes, say, the repetition period of a grating by the identity of the most active frequency-selective neurone in the visual cortex, or by some other measure of central tendency derived from the distribution of activity among the whole population of neurones. Assume that prolonged stimulation reduces the overall sensitivity of a neurone without altering the characteristic frequency--the frequency to which it is most sensitive. Consequently, adaptation at one particular spatial frequency selectively depresses the sensitivity of neurones whose characteristic frequencies are in the vicinity of that frequency. Subsequent observation of a test grating of somewhat different spatial frequency will produce a distribution of excitation different from normal. The central tendency of the distribution will be shifted away from the adapting frequency, and hence so should the perceived frequency of the test grating.

Blakemore et al. (1970) stated that the perceived spatial frequency shifts obtained after adaptation were evidence

-
1. This explanation of an adaptation after-effect by a shift in the relative output distribution of neural channels is not specific to spatial frequency channels. In color channels, an analogous after-effect might occur: after staring at a red light, a yellow light will appear slightly greener than previously. The red channel has had its sensitivity reduced due to adaptation, thereby making its contribution to the output signal of the color channels relatively less than previously when a yellow light is observed.

for frequency selective neurons in the human brain which were "actually involved in the encoding and perception of the size of simple patterns" (p. 727). However, these authors used sine wave gratings as adapting and test stimuli. Since frequency and period are simply inverses of each other when sine wave gratings are used as stimuli, frequency and bar width have a perfect correlation. Therefore, it is possible to describe their results in terms of width or size, as well as frequency. When adapting bars are narrower than the test bars, the test bars appear even wider; when the adapting bars are wider than the test bars, the test bars appear even narrower; when adapting and test bars are equal² there is no size shift.

The Blakemore et al. (1970) after-effect is equally consistent with size and frequency domain models. This has been supported by the computer simulation work of MacLeod and Rosenfeld (1974). If one uses a multiple size channels hypothesis assuming size-specific rather than frequency-specific

-
2. Classical figural after-effects (Kohler & Wallach, 1944) demonstrated that contours of a fixated test pattern are perceptually changed in the direction away from the features of the previously fixated inspection pattern. These data are the same as those obtained by Blakemore and Sutton (1969), however, different mechanisms for the after-effects were proposed. The proposed physiological correlate of figural aftereffects was displacement of the path of cortical current flow. Unlike proposed frequency selective neurons, this mechanism has since been discounted (Sperry & Miner, 1955).

neurons, adaptation should make similar size stimuli appear more different from the adapting size than they really are, regardless of the spatial frequency composition of the stimuli. Because of the single frequency, one dimensional nature of the sine wave adapting and test stimuli, the Blakemore et al. (1970) after-effect cannot discriminate between size and spatial frequency models of perception of spatial dimensions. What is needed, then, is to find the effects of adapting to a sine wave stimulus on a test stimulus where the correlation between size and spatial frequency is more complex.

Frome et al. (1974) asked observers to judge the size of an object with a continuous, multiple frequency spectrum. The observers adjusted the height of a rectangle so as to make the object look square, both before and after adaptation to a sine wave grating. Their results were dramatically different from those of Blakemore et al. adaptation to a sine wave grating produced an apparent enlargement of the size of the rectangles on the dimension orthogonal to the bars of the grating, whether the grating bar was smaller than, larger than or equal to the size of the rectangle width. In other words, after adaptation to a vertical (horizontal) grating, a square looked squat (skinny). These results are not consistent with a neural size channels

hypothesis or with classical figural after-effect data, as were Blakemore et al.'s (1970). The following experiments were designed to test several reasons why we obtained differences: the first experiment examined the possibility that the differences could be due to the judgments being made.

EXPERIMENT 1

The purpose of Experiment 1 is to compare frequency judgments with size judgments. The way to compare these judgments is to find one test stimulus that an observer can use to make a size or a spatial frequency judgment, both pre- and post-adaptation to a grating. The chosen test stimulus is a periodic pattern of multiple filled rectangles, that is, a very short grating.

Figure 5 is similar to the Blakemore and Sutton (1969) demonstration, with the short grating as test stimulus. As the reader may demonstrate for herself, frequency shifts like those obtained by Blakemore and Sutton (1969) result. That is, after adaptation, perceived frequency shifts in the short test grating are obtained to both lower and higher frequencies. Figure 5 demonstrates that a mere change in the length of the bars in the test grating will not change the direction of the perceived frequency shift. It is therefore unlikely that the differences in the results of Blakemore et al. (1970) and Frome et al. (1974) are due to Frome's test object being shorter.

In Experiment 1 a short grating is used as a test stimulus pre- and post-adaptation to a sine wave grating. In this experiment, as opposed to the demonstration in Figure 5, size judgments, not spatial frequency judgments are being

Figure 5. Pattern for demonstration of the perceived spatial frequency shift with short grating as test stimulus.

made. The direction of perceived frequency shifts is apparent; the direction of perceived size shifts with the same stimuli is being tested.

Methods

Subjects

There were four undergraduate subjects who made observations at the University of Massachusetts. Of these four subjects, two (D.R. and J.S.) were naive with respect to the purpose and expected direction of results of the experiment. These subjects were paid for their observations. Subjects D.W. and J.B. were undergraduate research assistants who were receiving academic credit for participating in the research described in Experiments 1 and 2.

Two subjects made observations in a similar experiment conducted at the University of Maryland. Subject M.W. was a paid naive subject, while J.L. was a highly practiced psychophysical observer who was a colleague in this experiment.

Seven unpaid naive subjects at the University of Maryland each made observations on only 2 conditions.

When necessary, observers wore glasses to bring them to 20:20 vision by the Snellen test.

Apparatus and Procedure

Adapting stimuli were three high contrast (40%) sinusoidal gratings of different spatial frequencies: 3, 4, and

6 c/deg. These were chosen because they are in the region of maximal sensitivity of the visual system and have shown large magnitude adaptation effects. These gratings were used in both horizontal and vertical orientations. Every adapting grating was used with each test stimulus.

Test stimuli were short square wave gratings of variable height and fixed spatial frequency--3, 4, and 6 c/deg. The observer's task was to adjust the height of the short grating to make the elements look square ($n = 11$).

Each experimental session consisted of the control condition with a homogeneous adapting field, a 10 minute rest period, and the experimental grating adaptation condition. The homogeneous field in the control condition was of the same average luminance as the grating field (135 nits). The initial adaptation in both the control and grating adaptation was 5 minutes long. During adaptation, the observer scanned the field in the direction orthogonal to the bars of the adapting grating being used. This scanning was an attempt to avoid the formation of an afterimage. After the initial adaptation there was a judgment of the vertical extent needed for perceived squareness of the short test grating. After each judgment ($n = 11$) there were 15 seconds of re-adaptation. These adaptation times were chosen to maintain adaptation at a maximum level in accordance with the

findings of Blakemore and Campbell (1969). A practice session for which no results were analyzed was conducted first in order to acquaint the observers with the task. Each observer completed only one experimental session per day.

The short square wave test grating was electronically generated on a cathode ray tube equipped with a P-31 phosphor. The equipment was masked so that the screen of the CRT was seen through a circular aperture cut in a large surround. There were always a minimum of 9 cycles of the grating on the screen. The observer viewed the test and adapting gratings binocularly while positioned in a chinrest. The adapting gratings were 35mm slides, projected on a screen at right angles to the CRT. The adapting field was a six degree circular field.

None of the naive subjects were shown the modified Blakemore and Sutton demonstration (Figure 5) until after the experiment was complete. No subjects had any difficulty obtaining the perceived frequency shift with the short grating as test object. No quantitative measurements of the perceived frequency shift with this test object were made, as the direction of the shift in perceived frequency is apparent.

Results

Figures 6-11 are graphs of individual subject data, when subjects were asked to make the bars in a very short grating square. Figures 6-9 are results obtained at the University of Massachusetts. Figures 10 and 11 were results obtained at the University of Maryland. The abscissae in these figures are the ratio of adapting grating bar width (P_A) to test rectangle widths (P_T). Abscissa values less than 1 indicate that a bar in the adapting grating was smaller than the width of a bar in the test grating, and an abscissa value greater than 1 indicates that the adapting grating bar was wider than the test grating bar. Table 1 indicates the adapting frequencies and rectangle sizes used to obtain these ratios. Each ordinate point is the mean of the ratios of the vertical extent needed for perceived squareness in the experimental grating adaptation (E) to the mean vertical extent needed for squareness in the control condition ($\bar{C}$), before grating adaptation. The height judgments are taken as a measure of perceived width. Ordinate values greater than 1, above the horizontal axis, indicate apparent widening after adaptation; in other words, after adaptation a greater vertical extent was required for perceived squareness. Ordinate values less than 1, below the axis, indicate apparent narrow-

Figure 6. An individual subject's data from Experiment 1, which had multiple filled squares, that is a short grating, as test object. The abscissa is the ratio of adapting bar width (P_A) to test rectangle widths (P_T). Each ordinate is the mean of the ratios of the vertical extent needed for perceived squareness in the experimental condition (E), after grating adaptation, to the mean vertical extent needed for squareness in the control condition before adaptation ($\bar{C}$). Standard errors of the mean are plotted for each datum. The left side of the figure illustrates results obtained when adapting with horizontally oriented grating bars, and the right side illustrates results when vertically oriented adapting gratings bars were used. These same coordinate axes are used for all other data in Experiment 1, Figures 7-11.

Figure 7. An individual subject's data from Experiment 1.

Figure 8. An individual subject's data from Experiment 1.

Figure 9. An individual subject's data from Experiment 1.

Figure 10. An individual subject's data from Experiment 1.

Figure 11. An individual subject's data from Experiment 1.

TABLE 1

Adapting grating, and test rectangle sizes used to obtain P_A/P_T conditions.

Bar Width Rectangle Width	$\frac{P_A}{P_T}$	Adapting Grating (P_A) (Bar Width)	Rectangle Width (P_T)
.5		6 c/deg. (5')	10'
. $\bar{6}$		6 c/deg. (5')	7.5'
.75		4 c/deg. (7.5')	10'
1.0		3 c/deg. (10')	10'
		4 c/deg. (7.5')	7.5'
		6 c/deg. (5')	5'
1. $\bar{3}$		3 c/deg. (10')	7.5'
1.5		4 c/deg. (7.5')	5'
2.0		3 c/deg. (10')	5'

ing after adaptation; that is after adaptation the rectangle appeared taller since less vertical extent was needed for the figure to be perceived as square. The right sides of these figures illustrate the results obtained when adapting with vertically oriented grating bars, and the left sides illustrate results obtained when horizontally oriented adapting grating bars were used. The bars on each datum indicate one standard error of the mean.

Except for one obvious point in Figure 6 and one in Figure 7, these results indicate that for nearly all width ratios less than 2.0, adaptation to a vertical (horizontal) sine wave grating produced significant widening (narrowing) of a short grating of multiple-filled squares. Significance test results are shown in Table 2.

Figure 12 is a nominal graph for six totally naive observers at the University of Maryland. Each made unpaid judgments at bar to rectangle ratio 1.5. There were few significant results with these subjects. Significance test results are shown in Table 3. For those subjects with a significant result with one grating there was no significant result in the opposite direction with the orthogonal grating.

TABLE 2

Multiple filled squares: t tests for the mean of the ratios ($n=11$) of the vertical extent needed for squareness in the adaptation condition to the mean vertical extent needed for squareness in the control condition.

$\frac{P_A}{P_T}$	Subject	Orientation (Horizontal or Vertical)	$\frac{E}{C}$	S.E.	t	Signif- icance
.5	D.W.	H	.930	.0191	-3.681	$p < .05$
		V	1.031	.0159	1.893	N.S.
	J.B.	H	.980	.0061	-3.262	$p < .05$
		V	1.036	.0100	3.600	$p < .05$
	D.R.	H	.902	.0103	-9.476	$p < .05$
		H	.946	.0122	-4.418	$p < .05$
		V	.983	.0138	-1.246	N.S.
	M.W.	V	1.015	.0146	.993	N.S.
	D.W.	H	.993	.0158	-.424	N.S.
		V	1.058	.0108	5.333	$p < .05$
.6	J.B.	H	.967	.0081	-4.025	$p < .05$
		V	1.031	.0101	3.030	$p < .05$
	D.R.	V	.991	.0127	-.724	N.S.
	M.W.	V	1.033	.0068	4.882	$p < .05$
	J.L.	H	.994	.008	-.739	N.S.
	D.W.	H	.958	.009	-4.644	$p < .05$
		H	.968	.020	-1.606	N.S.
		H	.940	.0155	-3.897	$p < .05$
	D.W.	H	.958	.009	-4.644	$p < .05$
		H	.968	.020	-1.606	N.S.
.75	M.W.	V	.996	.0112	-.384	N.S.
1.0	D.W.	H	.958	.009	-4.644	$p < .05$
		H	.968	.020	-1.606	N.S.
		H	.940	.0155	-3.897	$p < .05$

TABLE 2 (continued)

$\frac{P_A}{P_T}$	Subject	Orientation (Horizontal or Vertical)	$\frac{E}{\bar{C}}$	S.E.	$\underline{t}$	Signif- icance
1.0	D.W.	H	1.082	.0138	6.167	$p < .05$
		H	.989	.0150	- .720	N.S.
		H	.992	.0098	- .816	N.S.
		V	1.059	.0120	4.900	$p < .05$
		V	1.067	.0140	4.757	$p < .05$
	J.B.	V	1.060	.0124	4.855	$p < .05$
		H	.970	.0043	-6.884	$p < .05$
		H	.906	.0080	-11.738	$p < .05$
		H	1.075	.0181	4.149	$p < .05$
		H	.974	.0172	-1.506	N.S.
		H	.963	.0077	-4.844	$p < .05$
		V	.990	.0134	- .784	N.S.
		V	1.065	.0131	4.931	$p < .05$
	D.R.	H	.949	.0098	-5.245	$p < .05$
		H	.942	.0144	-4.063	$p < .05$
		H	.945	.0093	-5.970	$p < .05$
		V	1.065	.0150	4.300	$p < .05$
		V	1.004	.0142	.268	N.S.
		V	1.043	.0165	2.576	$p < .05$
	J.S.	H	.968	.0256	-1.258	N.S.
		V	1.023	.0338	.669	N.S.
	M.W.	H	.984	.0199	-1.047	N.S.
		H	.963	.0153	-2.405	$p < .05$
		V	1.132	.0149	8.873	$p < .05$
		V	1.023	.0133	1.729	N.S.
		V	1.009	.0146	.644	N.S.
	J.L.	H	.989	.0041	-2.488	$p < .05$
1.3	D.W.	H	.978	.0105	-2.105	N.S.
		V	1.019	.0128	1.492	N.S.
		H	.997	.0095	- .242	N.S.
		V	1.014	.0092	1.544	N.S.

TABLE 2 (continued)

$\frac{P_A}{P_T}$	Subject	Orientation (Horizontal or Vertical)	$\frac{E}{C}$	S.E.	t	Signif- icance
1.3	D.R.	H	.912	.0155	-5.658	$p < .05$
		V	.998	.0133	-.165	N.S.
	M.W.	H	.973	.0110	-2.482	$p < .05$
		V	1.003	.0119	.210	N.S.
	J.L.	V	1.007	.0080	.813	N.S.
	D.W.	H	.947	.0135	-3.948	$p < .05$
		H	.902	.0133	-7.368	$p < .05$
		H	.961	.0179	-2.220	$p < .05$
		V	1.019	.0138	1.406	N.S.
		V	1.110	.0148	7.446	$p < .05$
		V	1.131	.0141	9.319	$p < .05$
1.5	J.B.	H	.949	.0160	-3.168	$p < .05$
		H	.967	.0100	-3.300	$p < .05$
		V	1.060	.0210	2.870	$p < .05$
		V	1.073	.0093	7.871	$p < .05$
	D.R.	H	.921	.0292	-2.712	$p < .05$
		H	.942	.0166	-3.494	$p < .05$
		H	.928	.0199	-3.623	$p < .05$
		V	1.096	.0319	2.994	$p < .05$
		V	1.039	.0211	1.825	N.S.
		V	1.062	.0218	2.821	$p < .05$
	J.S.	H	.973	.0238	-1.143	N.S.
		V	1.417	.0337	12.365	$p < .05$
	M.W.	H	.972	.0103	-2.718	$p < .05$
		H	.987	.0126	-1.008	$p < .05$
		H	.989	.0077	-1.389	N.S.
		V	1.120	.0133	9.030	$p < .05$
		V	1.012	.0107	1.084	N.S.
		V	1.015	.0131	1.130	N.S.
		V	1.032	.0156	2.019	N.S.
		V	1.014	.0196	.694	N.S.

TABLE 2 (continued)

$\frac{P_A}{P_T}$	Subject	Orientation (Horizontal or Vertical)	$\frac{E}{\bar{C}}$	S.E.	$\underline{t}$	Signif- icance
1.5	J.L.	H	.980	.0050	-3.920	$p < .05$
		H	.977	.0116	-1.966	N.S.
		V	1.021	.0143	1.469	N.S.
		V	1.013	.0116	1.112	N.S.
		V	.998	.0092	-.272	N.S.
2.0	D.W.	H	1.016	.0157	1.000	N.S.
		H	.982	.0180	-1.050	N.S.
	J.B.	H	.977	.0130	-1.777	N.S.
		V	.991	.0092	-.967	N.S.
	D.R.	H	.966	.0227	-1.502	N.S.
		V	.997	.0235	-.145	N.S.
	M.W.	V	.969	.0201	-1.562	N.S.

Figure 12. Seven naive subjects' data for a P_A/P_T ratio of 1.5 in Experiment 1.

TABLE 3

Multiple filled Squares, naive University of Maryland Subjects: t tests for the mean of the ratios ($n=11$) of the vertical extent needed for squareness in the adaptation condition to the mean vertical extent needed for squareness in the control condition.

$\frac{P_A}{P_T}$	Subject	Orientation (Horizontal or Vertical)	$\frac{E}{\bar{C}}$	S.E.	$\underline{t}$	Signif- icance
1.5	P.M.	H	.994	.0095	- .674	N.S.
		V	.999	.0165	- .061	N.S.
	T.M.	H	.989	.0054	-2.130	N.S.
		V	1.000	.0082	- .061	N.S.
	P.H.	H	.986	.0151	- .947	N.S.
		V	.964	.0150	-2.373	$p < .05$
		V	.973	.0110	-2.491	$p < .05$
	S.P.	H	1.011	.0120	.908	N.S.
		V	.985	.0143	-1.049	N.S.
		V	1.008	.0068	1.471	N.S.
	D.L.	H	.899	.0139	-7.245	$p < .05$
		V	1.013	.0146	.897	N.S.
	M.H.	H	1.007	.0095	.695	N.S.
		V	1.033	.0407	.813	N.S.
		V	1.076	.0191	3.963	$p < .05$
	D.W.	V	.988	.0126	- .909	N.S.

Discussion

In general the multiple test rectangles appeared wider (narrower) after adaptation, regardless of whether the bars in the vertical (horizontal) adapting grating are smaller than, equal to, or larger than the bars in the short test grating. The size after-effect results obtained using a short multiple square grating as test stimulus are different from those obtained by Blakemore et al. (1970) using "tall" gratings. Figure 13 is their results plotted on the same axes as the results of the size after-effect with the short grating test stimulus. Since they only used adaptation to vertical gratings to obtain these results, this curve is only comparable to the right half of Figures 6-11. Note the dip in the right part of Figure 13. This portion of the curve indicates that after adapting to a grating with wider bars than the test grating bars, the test grating is judged to be higher in spatial frequency than previously. The corresponding size ratios of 1.0-2.0 in Figures 6-11 illustrate size judgments of a grating, show no such dip. I must therefore conclude, as Sullivan, Georgeson, and Oatley (1972) did for threshold judgments, that "width" is not equivalent to "spatial frequency."

In this experiment, the same test stimuli produced

Figure 13. Quantitative results of Blakemore et al. (1970) obtained with vertical sine wave gratings as both adapting and test stimuli which have been replotted on the same coordinate axes as the data in the present experiments.

different functions for different tasks. Carpenter and Ganz (1972) had similar findings when the same test stimuli produced different masking functions for a detection and a vernier acuity task. They proposed an attentional mechanism such that the output of spatial frequency analyzers are weighted differentially, depending on the task:

This model could also provide a parsimonious explanation of the data in this experiment. Both a single filled rectangle (Frome, 1975) and a short grating (multiple filled rectangles) appeared wider whether a bar in the sinusoidal adapting grating was smaller than, equal to, or larger than the filled rectangle(s) width. It is possible that in the multiple rectangle condition, the visual system attends only to a single rectangle, and thus the size after-effect is the same as that for the single rectangle condition. If Carpenter and Ganz's (1972) model of weighted spatial frequency analyzers as the mechanism for attention were correct, then for the visual system to attend to only one of the bars in the multiple rectangle condition, the frequency spectrum of the multiple rectangle stimulus must be weighted in a very specific way. According to their model, attention to a single bar the multiple bar grating would mean weighting the output of the frequency channels in such a way that they would produce an output more like the distribution of

activity produced by an unweighted single bar. Figure 14 shows the frequency spectra of these stimuli. As can be seen, if a periodic stimulus is not extended infinitely (the spectrum of the grating assumes infinite repetition), the spectral lines are smeared, as in the case of a single rectangle. In order to produce an output which resembles the activity produced by a single rectangle by weighting the activity normally produced by the multiple rectangle grating, those frequency selective channels which are not maximally stimulated by the spectral lines of the grating must be weighted relatively more in the final size judgment. This would produce a weighted output function resembling the $\sin x/x$ spectral distribution of a single bar.

In other words, if a subject is attending to a single bar in the grating because that is where the salient visual dimensions for judging size lie, a weighting function model like that proposed by Carpenter and Ganz (1972) provides a plausible mechanism for this process. However, there is little direct evidence that the attentional mechanisms just described actually occur in the visual system.

Figure 14. The one-dimensional spectral distributions of a single filled square and of multiple filled squares.

EXPERIMENT 2

Experiment 2 explores the contribution of a factor which might have influenced the perception of size or spatial frequency of a multiple square grating. Adaptation to a sine wave grating may cause the perceived duty cycle (light to dark bar width ratio) of the test grating to change. For example, it may be that after adaptation to a grating, the light rectangles tend to fill in the dark spaces. This might account for the paradoxical "judgment" of wider size, but higher spatial frequency; however, at the same time, not also account for perceptions of both wider sizes and lower spatial frequencies. In other words, if the effect of adaptation on the perceived duty cycle of a multiple square grating were to account for both the size and spatial frequency judgments, there must be differential adaptation effects on duty cycle across the combinations of adapting grating and test grating frequencies. Specifically, there would have to be little or constant effect of adaptation on perceived duty cycle where adaptation produces judgments of wider size and lower spatial frequency (that is, at adapting period to test period ratios less than 1). In addition, adaptation must cause the light rectangles to fill in the dark spaces at adapting period to test period ratios from a little less than 1 to almost a ratio of 2,

that is at those conditions where the directions of the Blakemore et al. (1970) and Frome et al. (1974) effects diverge.

This experiment, therefore, tests the possibility that adaptation to a sine wave grating may cause the perceived duty cycle of the test grating to change. At the same time, this experiment tests the possibility that adaptation effects on the perceived duty cycle of a multiple square grating would account for both the judgments of size and spatial frequency.

Methods

Subjects

There were three undergraduate subjects who made observations at the University of Massachusetts. One subject, G. K., was naive with respect to the purpose and expected direction of results of the experiment. He was paid for his observations. Subjects D. W. and J. B. were undergraduate research assistants who were receiving academic credit for participating in the research described in Experiments 1 and 2.

Apparatus and Procedure

The apparatus and general procedures described in Experiment 1 were also used in this experiment. The specific procedures necessary for this experiment are as

follows.

Observers were asked to set the elements of the short 50% duty cycle grating so that they look square, using the same procedure as in Experiment 1. The height of the rectangles was then fixed to the computed mean of these judgments. Observers were then asked to set the squares to an apparent 50% duty cycle ($n=11$). Following these judgments, observers adapted to a vertical sinusoidal grating and then set the perceived multiple square pattern to 50% duty cycle again ($n=11$). Through this procedure, any effect in the perceived size change of the multiple square pattern after adaptation that is due to an apparent change in duty cycle may be obtained.

Results

Figures 15-17 are individual subject data when subjects were asked to set a perceived multiple square grating to 50% duty cycle both before and after adaptation to a vertical sine wave grating. The abscissae in these figures are the ratio of adapting grating period to short test grating period. Abscissa values less than 1 indicate that the period of the adapting grating was smaller than the period of the test grating. Abscissa values greater than 1 indicate that the period of the adapting grating was

Figure 15. An individual subject's data from Experiment 2. Subjects set a perceived multiple square grating to 50% duty cycle (equal bar and space width) both before and after adaptation to a vertical sine wave grating. The abscissa is the ratio of adapting grating period (P_A) to short test grating period (P_T). Each ordinate is the mean of the ratios of the horizontal extent of the dim spaces needed for perceived 50% duty cycle of the perceived multiple square grating in the experimental condition (E), after grating adaptation, to the mean horizontal extent of the dim spaces needed for perceived 50% duty cycle in the control condition before adaptation ($\bar{C}$). Standard errors of the mean are plotted for each datum. These same coordinate axes are used for all other data in Experiment 2, Figures 16 and 17.

Figure 16. An individual subject's data from Experiment 2.

Figure 17. An individual subject's data from Experiment 2.

larger than that of the short test grating. Each ordinate point is the mean of the ratios of the horizontal extent of the dim spaces needed for perceived 50% duty cycle of the perceived multiple square test grating after adaptation to the mean horizontal extent of the dim spaces needed for the same criterion (equal bar and space width) before sine wave adaptation. The bars on each datum indicate one standard error of the mean. The use of these coordinate ratio measures allows easy comparison to the results from Experiment 1, in that ordinate values greater than 1 indicate perceived widening of the bright squares after adaptation, while ordinate values less than 1 indicate perceived narrowing of the bright squares after adaptation. This point is illustrated and explained in Figure 18.

As can be seen in Figures 15-17, for nearly all period ratios, after adaptation to a vertical sinusoid, the bright bars in a short grating (multiple perceived squares) become perceptually narrower than the dim spaces. The significance test results are shown in Table 4. As both the figures and t-tests show, the aftereffects from this experiment were often small and not significant. However, in no case was the effect of adaptation to cause the bright squares to fill in some of the dim space.

Figure 18. An illustration and explanation of subject's setting and perception of a 50% duty cycle grating before and after adaptation to a vertical grating.

Physically 50% cycle grating:

BEFORE ADAPTATION:

Ss setting:

The physically smaller horizontal extent of the bright bars is perceived as equal to the extent of the wider spaces. Therefore, a physically measured 50% duty cycle grating would appear to have wider bright bars than dark spaces.

Ss psychological perception of a physically 50% duty cycle grating:

AFTER ADAPTATION:

If adaptation causes perceived 'narrowing' of the bright bars:

S's setting:

The physically larger horizontal extent of the bright bars is perceived as equal to the extent of the narrower spaces.

Therefore, the bright bars of a 50% duty cycle grating would then be perceptually narrower than the dim spaces.

Ss psychological perception of a physically 50% duty cycle grating would then be:

Outline rectangles represent bright bars. The physical dimensions are exaggerated for purposes of explanation.

TABLE 4

Perceived multiple squares duty cycle: t tests for the mean of the ratios ($n=11$) of the horizontal extent of the dim spaces needed for perceived 50% duty cycle in the adaptation condition to the mean horizontal extent of the dim spaces needed for perceived 50% duty cycle in the control condition.*

$\frac{P_A}{P_T}$	Subject	$\frac{E}{C}$	S.E.	t	Significance
.5	D.W.	.988	.0059	-2.102	N.S.
	J.B.	.983	.0062	-2.742	$p < .05$
	G.K.	1.001	.0034	.294	N.S.
.6	D.W.	.952	.0083	-5.819	$p < .05$
	J.B.	.982	.0069	-2.580	$p < .05$
	G.K.	.988	.0041	-2.829	$p < .05$
.75	J.B.	.993	.0037	-1.946	N.S.
1.0	D.W.	.976	.0045	-5.289	$p < .05$
	J.B.	.995	.0057	-.947	N.S.
	G.K.	.984	.0030	-5.000	$p < .05$
1.3	J.B.	1.002	.0068	.2790	N.S.
	G.K.	1.005	.0025	1.880	N.S.

TABLE 4 (continued)

$\frac{P_A}{P_T}$	Subject	$\frac{E}{\bar{C}}$	S.E.	$\underline{t}$	Significance
1.5	D.W.	.996	.0054	- .667	N.S.
	J.B.	.990	.0062	-1.661	N.S.
	G.K.	.979	.0050	-4.140	$p < .05$
2.0	J.B.	.997	.0040	- .700	N.S.
	G.K.	.992	.0039	-2.077	N.S.

* All adapting grating orientations were vertical.

Discussion

Prior to this experiment, it was thought that effects of adaptation on the apparent duty cycle of a multiple square grating might provide a sensory level explanation of the paradoxical effect found in Experiment 1, that adaptation to a lower frequency leads observers to judge a test grating to be higher in spatial frequency but wider in size. This would require differential effects from adaptation on apparent duty cycle across the combinations of adapting and test frequencies such that adaptation to a frequency that is equal to or lower than the frequency of the test grating would decrease the apparent amount of space between squares. The findings of this experiment do not meet these requirements and cannot provide such a sensory level explanation since adaptation to a lower frequency grating seems to make some small increases in the apparent amount of space between squares.

The findings of this experiment merely strengthen the paradox established in Experiment 1: adaptation which decreases the apparent width of bright bars per cycle, also yields judgments of higher spatial frequency, and can also give rise to perceptions of wider size. One could adjust the findings of Experiment 1 to eliminate the adaptation

effects on perceived duty cycle. Since duty cycle adaptation effects are in the direction of narrowing, adjustment for this would slightly increase the apparent widening effects found in Experiment 1.

EXPERIMENT 3

Experiment 3 controls for the possibility that the effective psychological width of a bar in the sine wave adapting grating was so much smaller than its physical ($1/2$ period) width, that in fact, the obtained results of no perceived narrowing may be due to a displacement of size of the filled test rectangles away from the smaller psychological width of the sine wave bar size. One reason that the effective stimulus width of the bars of the adapting grating may have been psychologically smaller than they were physically is that the contours in the adapting stimulus were not distinct, since they had a sinusoidal luminance distribution. In Experiment 3, the comparison of square wave gratings to sine wave gratings as adapting stimuli control for this possibility, since a square wave bar can have the same physical width as a sine wave bar but will have distinct contours.

A square wave has a very different frequency spectrum than a sine wave. However, precautions have been taken to minimize these differences so that the difference in contour between the sine and square wave is the major variable allowed to effect the results. In general, a square wave of the same fundamental frequency, has a

fundamental of greater amplitude and in addition has all the odd harmonics, with amplitudes reciprocal to the number of their harmonic. In this experiment the amplitudes of the fundamentals of the same frequency sine and square wave have been matched as nearly as possible. Figure 19 shows the frequency spectrum of a 7.5 min filled rectangle, described by the function $\sin x/x$. The open arrows indicate the fundamental frequencies of the adapting gratings that were used. The filled arrows indicate the frequencies of the third harmonics of the square waves that were used. The combination of the increasing decrement of amplitude in the sine function and the smaller amplitude of the third harmonics at higher frequencies will work together to reduce the effectiveness of the component third harmonic adaptation in the square waves.

Adaptation with the fundamental sinusoidal gratings shown in Figure 19 should replicate Frome et al.'s (1974) original finding that although a single bar in these gratings are larger than, equal to, or smaller than the rectangle width, apparent widening should always result. If this result is due to the fact that sine wave bars have a narrower psychological than physical width because of their contour, then results from adaptation to square wave gratings should be very different. If results from adaptation

Figure 19. The one-dimensional spectral distribution of a 7.5 min arc filled rectangle. The frequency spectrum may be described by the function $\sin x/x$. The open arrows indicate the fundamental frequencies of the adapting gratings that were used in Experiment 3. The filled arrows indicate the frequencies of the third harmonics of the square waves that were used.

to a square wave were consistent with a simple neural size channels hypothesis which predicts shifts in apparent size away from the adapting size, then the results from adapting to a square wave with vertical bars should look like those obtained by Blakemore et al. (1970), illustrated in Figure 13.

Methods

Subjects

There were four undergraduate subjects who made observations at the University of Massachusetts. All subjects were paid and were naive with respect to the purpose and expected direction of results.

Apparatus and Procedure

The apparatus and general procedures described in Experiment 1 were also used in this experiment. The specific difference was that only one bar in the generated square wave test object was visible to the subject. The observer's task was to adjust the vertical dimension of that rectangle so as to make the figure look square ($n=11$), before and after adaptation to a vertical grating. Two subjects were adapted to a sine wave at a particular size ratio two days prior to their next adaptation, which was to a square wave, until all conditions were complete. The other two subjects were run in the square wave adaptation

condition first, followed by sine wave exposure on different days.

Results

Figures 20-23 are individual subject data obtained with adaptation to both sine and square wave gratings as described above. The abscissae and ordinates in these figures are the same as in Experiment 1. The bars on each datum indicate one standard error of the mean. The subjects in Figures 20 and 21 were adapted to a sine wave at a particular size ratio two days prior to their next adaptation, which was to a square wave. Figures 22 and 23 are data similarly obtained, except they were run in the square wave adaptation condition first, followed by sine wave exposure on different days. There are no obvious differences due to this running order. Figure 24 shows the average for all observers' data, with the standard error of the mean between subjects indicated for each data point.

As Figures 20-24 show, after adaptation to both sine wave and square wave gratings with vertically oriented bars, the rectangle appeared wider for all width ratios. The amplitude of the after-effect seems nearly equal at all width ratios. Significance test results are shown in Table 5. Sine wave and square wave adaptation effects are

Figure 20. An individual subject's data from Experiment 3, which compares results from vertical square wave and sine wave adapting gratings. Standard errors of the mean are plotted for each datum. The coordinate axes used for all data in Experiment 3, Figures 20-24, are the same as those used in Experiment 1.

Figure 21. An individual subject's data from Experiment 3.

Figure 22. An individual subject's data from Experiment 3.

Figure 23. An individual subject's data from Experiment 3.

Figure 24. The average of all subjects' data from Experiment 3, with the standard error of the mean between subjects indicated for each datum.

TABLE 5

Sine wave versus square wave adaptation: t tests for the mean of the ratios ($n=11$) of the vertical extent needed for squareness in the adaptation condition to the mean vertical extent needed for squareness in the control condition.*

$\frac{P_A}{P_T}$	Subject	Square or Sine Wave	$\frac{E}{C}$	S.E.	$\underline{t}$	Signif- icance
.6	D.A.	Square	1.033	.0129	2.527	$p < .05$
		Sine	1.017	.0127	1.339	N.S.
	S.K.	Square	1.013	.0134	.970	N.S.
		Sine	1.033	.0107	3.037	$p < .05$
	N.S.	Square	1.075	.0205	3.663	$p < .05$
		Sine	1.092	.0192	4.797	$p < .05$
	T.B.	Square	1.082	.0115	7.087	$p < .05$
		Sine	1.076	.0060	12.683	$p < .05$
	Mean	Square	1.076	.0160	3.188	$p < .05$
		Sine	1.054	.0177	3.073	$p < .05$
1.0	D.A.	Square	1.020	.0215	.916	N.S.
		Sine	1.047	.0126	3.722	$p < .05$
	S.K.	Square	1.031	.0155	2.026	N.S.
		Sine	1.031	.0245	1.249	N.S.
	N.S.	Square	1.121	.0190	6.369	$p < .05$
		Sine	1.049	.0137	3.540	$p < .05$
	T.B.	Square	1.037	.0240	1.550	N.S.
		Sine	1.111	.0224	4.964	$p < .05$
	Mean	Square	1.052	.0228	2.281	$p < .05$
		Sine	1.059	.0177	3.333	$p < .05$
1.3	S.K.	Square	1.017	.0169	1.174	N.S.
		Sine	1.115	.0198	5.788	$p < .05$

TABLE 5 (continued)

$\frac{P_A}{P_T}$	Subject	Square or Sine Wave	$\frac{E}{C}$	S.E.	$\underline{t}$	Signif- icance
1.3	N.S.	Square	1.086	.0161	5.366	$p < .05$
		Sine	1.032	.0157	2.026	N.S.
	T.B.	Square	1.052	.0217	2.406	$p < .05$
		Sine	1.034	.0307	1.098	N.S.
	Mean	Square	1.066	.0108	6.148	$p < .05$
		Sine	1.045	.0089	5.056	$p < .05$
1.5	D.A.	Square	1.025	.0178	1.405	N.S.
		Sine	1.077	.0318	2.421	$p < .05$

* All adapting grating orientations were vertical.

nearly identical.

Discussion

Neither adaptation to sine waves or square waves produces perceived narrowing in a single filled rectangle for larger period adapting stimuli, as a simple neural size channels hypothesis would predict. That square wave adaptation produces identical effects to sine wave adaptation implies that the sine wave after-effect of only apparent widening could not be due to a narrower psychological than physical width of the bars because of their blurry edges.

Since size repulsion after-effects do not seem to be occurring, it would be possible to use a square wave adapting stimulus to examine the consistency of these findings with a frequency analysis model. In Experiment 3, the effective amplitudes of the third harmonics in the square waves were small. One could try and maximize the effect of the third harmonic of the square wave so as to match the frequency and contrast of a sine wave grating to which results are compared. If adaptation is truly frequency specific and it is this frequency specificity which produces the size after-effects, one would expect adaptation to a square wave grating to produce perceived size shifts to its large amplitude harmonics as well. Blakemore et al.

(1970) have shown frequency shifts of target gratings at the third harmonic frequencies of adapting square wave gratings.³ It would be of interest then to see if the same size shifts would occur at the ratio of the third harmonic, $1/2$ period to rectangle width, as occurred for a single sine wave $1/2$ period to rectangle width.

-
3. Contrary to Blakemore and Campbell (1969), Nachmias, Sansbury, Vassilev, and Weber (1973) failed to find consistent evidence that the third harmonic in a square wave grating does any adapting at all to detection thresholds of sinusoidal test gratings whose frequency is in the vicinity of the third harmonic in the square wave. However, their failure to replicate the third harmonic effect may be due to procedural differences. These procedural differences are discussed in their paper.

EXPERIMENT 4

Experiment 4 introduces a new test target, an outline square. Previous test targets, a single filled square, and a short grating (a periodic pattern of multiple filled squares) have different frequency spectra, yet produce the same size after-effects. Although the spectra differ, they are similar in important ways: for a given width rectangle in these patterns, the spectral maxima and minima coincide, only the spread of the spectral lines differ (see Figure 14). As discussed above, an attentional mechanism would be a parsimonious explanation of the similarities of size after-effects for these different test targets. By using an outline square as a test target, subjects can use the same task as with the previous test targets; yet this stimulus has a very different frequency spectrum than the other test targets, as can be seen in Figure 25. By using the resultant size after-effects after adaptation, one can begin to weave hypotheses about assumptions which must be made for either size and/or frequency specific models.

Methods

Subjects

There were four subjects who made observations at the University of Maryland. Three subjects, M.W., T.M., and

Figure 25. The one-dimensional spectral distributions of 5' and 10' wide test stimuli: a single filled square, multiple filled squares, and an outline square. As the legend shows, arrows indicate different ratios of adapting grating bar width to square width.

LEGEND $\square$ ARROWS INDICATE (BAR WIDTH / SQUARE WIDTH) = $2.0 \nabla = 1.0 \nabla = 5 \nabla$

P.M., were paid and were naive with respect to the purposes and expected direction of results. Subject J.Z.L. was a highly practiced psychophysical observer who was a colleague in this experiment.

Apparatus and Procedure

The apparatus and general procedures described in Experiment 1 were also used in this experiment. The specific difference was that a single outline rectangle was electronically generated on the cathode ray tube as the test stimulus. The observer's task was to adjust the vertical dimension of the outline rectangle before and after adaptation to a vertical sinusoidal grating ($n=11$).

Results

Figures 26-29 show individual subject data when observers were asked to adjust the height of an outline rectangle so it appears square, both before and after adaptation to a vertical sine wave grating. The abscissae and ordinates in these figures are the same as those used for Experiments 1 and 3. Figure 30 is the average of the subjects' results with between subjects standard error indicated.

These data indicate that after adaptation to a vertical grating, both a filled square and an outline square look squat. These data resemble the filled square data,

Figure 26. An individual subject's data from Experiment 4, which uses an outline square as test object before and after adaptation to a vertical sine wave grating. Standard errors of the mean are plotted for each datum. The coordinate axes used for all data in Experiment 4, Figures 26-30, are the same as those used in Experiments 1 and 3.

Figure 27. An individual subject's data from Experiment 4.

Figure 28. An individual subject's data from Experiment 4.

Figure 29. An individual subject's data from Experiment 4.

Figure 30. The average of all subject's data in Experiment 4, with the standard error of the mean between subjects indicated for each datum.

TABLE 6

Outline Square: t tests for the mean of the ratios ($n=11$) of the vertical extent needed for squareness in the adaptation condition to the mean vertical extent needed for squareness in the control condition.*

$\frac{P_A}{P_T}$	Subject	$\frac{E}{\bar{C}}$	S.E.	t	Significance
.5	M.W.	1.057	.0112	5.080	$p < .05$
	T.M.	1.015	.0066	2.303	$p < .05$
	P.M.	1.029	.0080	3.575	$p < .05$
	J.Z.L.	1.019	.0085	2.282	$p < .05$
.6	M.W.	1.028	.0094	2.840	$p < .05$
	T.M.	1.042	.0086	4.895	$p < .05$
	J.Z.L.	1.010	.0042	2.357	$p < .05$
1.0	M.W.	1.029	.0058	4.966	$p < .05$
		1.036	.0111	3.198	$p < .05$
		1.041	.0114	3.605	$p < .05$
	T.M.	1.045	.0055	8.164	$p < .05$
		1.032	.0091	3.473	$p < .05$
		1.052	.0133	3.910	$p < .05$
	P.M.	1.050	.0121	4.149	$p < .05$
	J.Z.L.	1.028	.0067	4.209	$p < .05$
		1.025	.0084	3.000	$p < .05$
1.3	M.W.	1.001	.0141	.078	N.S.

TABLE 6 (continued)

$\frac{P_A}{P_T}$	Subject	$\frac{E}{\bar{C}}$	S.E.	$\underline{t}$	Significance
1.3	T.M.	1.039	.0062	6.258	$p < .05$
		.987	.0068	-1.912	N.S.
	J.Z.L.	1.017	.0084	2.056	N.S.
		1.033	.0094	3.553	$p < .05$
1.5	M.W.	1.027	.0148	1.845	N.S.
	T.M.	1.040	.0085	4.694	$p < .05$
	J.Z.L.	1.042	.0080	5.300	$p < .05$
2.0	M.W.	.995	.0086	- .535	N.S.
	T.M.	.998	.0127	- .142	N.S.
	P.M.	1.019	.0142	1.331	N.S.
	J.Z.L.	1.008	.0091	.912	N.S.
		.995	.0100	- .490	N.S.
2.67	M.W.	1.011	.0080	1.338	N.S.
	T.M.	1.009	.0117	.735	N.S.
	P.M.	1.061	.0154	3.961	$p < .05$
	J.Z.L.	.989	.0146	- .788	N.S.
		.997	.0132	.182	N.S.

* All adapting grating orientations were vertical.

shown in Figure 31, rather closely, with the exception that it was easier to obtain data for smaller test rectangles (adapting with a 3 c/d grating and testing with a 3.75 min arc rectangle).

Discussion

The perceived size shifts that were obtained after adaptation to a grating with the three test stimuli shown in Figure 25 are strikingly similar. Although the frequency spectra of these stimuli are very different, all test rectangle widths still gave perceived widening whether a bar of a vertical grating was smaller than, equal to, or larger than its width.

The direction of size after-effect results for the sinusoidal grating that Blakemore et al. (1970) used and the test stimuli used in these experiments, is very different. By extension of the size encoding model proposed by those authors, it may be possible to understand the results for solid and outline squares in terms of the frequency spectra of these patterns. Blakemore et al. (1970) propose that after adaptation, stimulus size is coded by the mean (or some other measure of central tendency) of the distribution of activity in frequency selective neural channels. The shift in central tendency of the neural output distri-

Figure 31. The average of all subjects' data when a single filled square was used as test object before and after adaptation to a vertical sinusoidal grating (After Frome et al., 1974). The standard error of the mean between subjects is indicated for each datum. The data have been plotted on the same coordinate axes as the data in the present experiments.

bution due to adaptation presents a more complicated situation in the interpretation of this data than in Blakemore et al.'s because of the multiple frequency distribution in our stimuli. At some ratios of grating bar to rectangle width there was no apparent widening after adaptation. Note especially the abscissa ratio 2.0 in Figures 30 and 31. Assuming Blakemore et al.'s size encoding model there can be two reasons for this:

- 1) the grating has not adapted the visual system, or
- 2) this is the psychological mean of the frequency distribution, since the reduction in sensitivity of the channel responding to the adapting frequency leaves the perceived size of the square unaffected.

Only the second reason can be the case if the data is examined. For both a filled and outline rectangle, there is no size shift at ratio 2.0, which is obtained after adapting to a 3 c/deg grating and then looking at a 5 min arc square. This trend away from increased perceived size cannot be due to failure of adaptation at 3 c/deg because the same adapting grating frequency when used with a 10 min arc square is a size ratio of 1.0 which yields large shifts in the increasing direction. Therefore, given Blakemore et al.'s model, the size ratios for 2.0 are equivalent

psychologically, although as can be seen in Figure 25 the adapting frequency will be effective on different frequency-amplitude distributions. The differences in the frequency distributions of an outline and filled square may be described as follows: The outline square has relatively less low-frequency energy. Much of the higher frequency energy that would be affected by the adaptations in these experiments, appears in phase opposition to the low frequencies. Also an outline rectangle has discrete top and bottom bars, which probably play a role in the perception of the rectangle's width. All these factors have not yet been assessed psychophysically, although they may be with the proper experimental manipulations. If one is to extend Blakemore et al.'s (1970) model of size encoding to other than grating patterns, the data from these experiments must be accounted for. Yet if one is to model such data without further psychophysical assessments, one must introduce ad hoc assumptions, such as an assumption about the effect of adaptation on the negative lobe of the outline square's cosine frequency spectrum.

An alternative hypothesis one could weave about these results is in terms of a size model. Since similar size ratios of adapting grating to test stimulus width yield similar results, regardless of the spectral composition of

the test target, it seems that an explanation of these data in terms of size as a primitive rather than spatial frequency would be more parsimonious. However, a simple neural size channels hypothesis, which would predict size shifts away from the adapting size would not explain the data of the present experiments. It would be of interest to investigate the size after-effects on the test targets in these experiments after adapting to a single bar. Although Sullivan, Georgeson, and Oatley (1972) found no width specific adaptation from bars at detection threshold, size judgments may yield different results. If adaptation to single bars of the same size ratios as those contained in the gratings used in the present experiment yielded curves similar to those in Experiments 1, 3, and 4, then one might interpret the present results as being obtained from adaptation to the size of individual bar width in the grating. If this is not the case, the weaving of size and spatial frequency interaction models resumes.

REFERENCES

- Anstis, S. M. Size adaptation to visual texture and print: Evidence for spatial frequency analysis. Paper presented at annual meeting of Assoc. for Res. in Vis. and Ophthalm., Sarasota, Fla., 1973.
- Bagrash, F. M., Thomas, J. P., & Shimamura, K. K. Size-tuned mechanisms: Correlation of data on detection and apparent size. Vision Res., 1974, 14, 937-942.
- Bagrash, F. M. Size-selective adaptation psychophysical evidence for tuning and the effects of stimulus contour and adapting flux. Vision Res., 1973, 13, 575-598.
- Blakemore, C. & Campbell, F. W. Adaptation to spatial stimuli. J. Physiol., Lond., 1969, 200, 11-13P.
- Blakemore, C. & Campbell, F. W. On the existence of neurones in the human visual system selectively sensitive to the orientation and size of retinal images. J. Physiol., 1969, 203, 237-260.
- Blakemore, C., Nachmias, J., & Sutton, P. Perceived spatial frequency shift: Evidence of frequency-selective neurones in the human brain. J. Physiol., Lond., 1970, 210, 727-750.
- Blakemore, C. & Sutton, P. Size adaptation: A new after-effect. Science, 1969, 166, 245-247.

- Campbell, F. W., Carpenter, R. H. S., & Levinson, J. Z.
Visibility of aperiodic patterns compared with that
of sinusoidal gratings. J. Physiol., 1969, 204, 283-
298.
- Campbell, F. W., Cooper, G. F., & Enroth-Cugell, C. The
spatial selectivity of the visual cells of the cat.
J. Physiol., 1969, 203, 223-235.
- Campbell, F. W., Cooper, G. F., Robson, J. G., & Sachs, M.
The spatial selectivity of visual cells of the cat and
the squirrel monkey. J. Physiol., Lond., 1969, 204,
120P-123P.
- Campbell, F. W. & Maffei, L. Electrophysiological evidence
for the existence of orientation and size detectors in
the human visual system. J. Physiol., 1970, 207, 635-
652.
- Campbell, F. W., Nachmias, J., & Jukes, J. Spatial frequen-
cy discrimination in human vision. J. Opt. Soc. Amer.,
1970, 60, 555-559.
- Campbell, F. W. & Robson, J. G. Application of Fourier ana-
lysis to the modulation response of the eye. J. Opt.
Soc. Amer., 1964, 54, 581A.
- Campbell, F. W. & Robson, J. G. Application of Fourier ana-
lysis to the visibility of gratings. J. Physiol., 1968,
197, 551-566.

- Carpenter, P. A. & Ganz, L. An attentional mechanism in the analysis of spatial frequency. Perception & Psychophysics, 1972, 12, 57-60.
- Cornsweet, T. N. Visual perception. New York: Academic Press, 1970.
- DeValois, K. Black and white: Equal but separate. Paper presented at annual meeting of Assoc. for Res. in Vis. and Ophthalm., Sarasota, Fla., 1973.
- Enroth-Cugell, C. & Robson, J. G. The contrast sensitivity of retinal ganglion cells of the cat. J. Physiol., Lond., 1966, 187, 517-552.
- Fiorentini, A. & Maffei, L. Suprathreshold contrast perception and its electrophysiological correlate. Optica Acta, 1972, 19, 363-364.
- Frome, F., Danielson, J. T., & Levinson, J. Z. Shifts in perception of size after adaptation to gratings. Paper presented at annual meeting of Assoc. for Res. in Vis. and Ophthalm., Sarasota, Fla., 1974.
- Frome, F. & Levinson, J. Z. Shifts in perception of size after adaptation to gratings II. Paper presented at annual meeting of Assoc. for Res. in Vis. and Ophthalm., Sarasota, Fla., 1975.

- Ganz, L. Mechanism of the figural aftereffects. Psychol. Rev., 1966, 73, 128-150.
- Gouras, P. Visual neurophysiology: Feature detecting channels. Investigative Ophthalmology, 1973, 12, 1, 2-3.
- Graham, C. H. (Ed). Vision and visual perception. New York: John Wiley & Sons, 1965.
- Graham, N. & Nachmias, J. Detection of grating patterns containing two spatial frequencies: Comparison of single channel and multiple channel models. Vision Res., 1971, 11, 251-259.
- Gross, C. G., Rocha Miranda, C. E., & Bender, D. B. Visual properties of neurons in inferotemporal cortex of the macaque. J. Neurophysiology, 1972, 35, 96-111.
- Hays, W. Statistics. New York: Holt, Rinehart, & Winston, 1963.
- Hoekstra, J., VanDerGoot, D. P. S., Vandenbrink, G., & Bilsen, F. A. The influence of the number of cycles upon the visual contrast threshold for spatial sine wave patterns. Vision Res., 1974, 14, 365-368.
- Hubel, D. H. & Wiesel, T. N. Receptive fields, binocular interaction and functional architecture in the cat's visual cortex. J. Physiol., Lond., 1962, 160, 106-154.

- Hubel, D. H. & Wiesel, T. N. Shape and arrangement of columns in cat's striate cortex. J. Physiol., Lond., 1963, 165, 559-568.
- Hubel, D. H. & Wiesel, T. N. Receptive fields and functional architecture in two nonstriate visual areas (18 and 19) of the cat. J. Neurophysiology, 1965, 28, 229-289.
- Hubel, D. H. & Wiesel, T. N. Receptive fields and functional architecture of monkey striate cortex. J. Physiol., 1968, 195, 215-243.
- Kaufman, L. Sight and mind, an introduction to visual perception. New York: Oxford University Press, 1974.
- Kelly, D. H. How many bars make a grating? Vision Res., 1975, 15, 625-626.
- Kitterle, F. L. The effect of adapting and test field color on the perception of size. Vision Res., 1975, 15, 7, 883-885.
- Klein, S., Stromeyer, C. F., & Ganz, L. The simultaneous spatial frequency shift: A dissociation between the detection and perception of gratings. Vision Res., 1974, 14, 1421-1432.
- Koehler, W. & Wallach, H. Figural after-effects. Proc. Amer. Phil. Soc., 1944, 88, 269-357.
- Lange, V., Sigel, C., & Stecher, S. Adapted and unadapted spatial frequency channels in human vision. Vision Res.

1973, 13, 11, 2139-2144.

Lettvin, J. Y., Maturana, H. R., McCulloch, W. S., & Pitts, W. H. What the frog's eye tells the frog's brain.

Proc. Inst. Radio Engr., 1959, 47, 1940-1951.

Macleod, I. D. G. & Rosenfeld, A. The visibility of gratings: Spatial frequency channels or bar-detecting units? Vision Res., 1974, 14, 909-915.

Maffei, L. & Fiorentini, A. Processes of synthesis in visual perception. Nature, 1972, 240, 479-481.

Maffei, L. & Fiorentini, A. The visual cortex as a spatial frequency analyser. Vision Res., 1973, 13, 1255-1268.

Maffei, L., Fiorentini, A., & Bisti, S. Neural correlate of perceptual adaptation to gratings. Science, 1973, 182, 1036-1038.

Nachmias, J. & Sansbury, R. V. Grating contrast: discrimination may be better than detection. Vision Res., 1974, 14, 1039-1042.

Nachmias, J., Sansbury, R., Vassilev, A., & Weber, A. Adaptation to square wave gratings: in search of the elusive third harmonic. Vision Res., 1973, 13, 1335-1342.

Nachmias, J. & Weber, A. Discrimination of simple and complex gratings. Vision Res., 1975, 15, 217-223.

- Neisser, U. Cognitive psychology. New York: Appleton-Century-Crofts (Meredith Corporation), 1967.
- Pantle, A. & Sekuler, R. Size detecting mechanisms in human vision. Science, 1968, 162, 1146-1148.
- Schade, O. H. Electro-optical characteristics of TV systems. I. Characteristics of vision and visual systems. RCA Review, 1948, 9, 5-37.
- Sekuler, R. Spatial vision. Ann. Review of Psych., 1974, 195-232.
- Sperry, R. W. & Miner, N. Pattern perception following insertion of mica plates into visual cortex. Journal of Comparative and Physiological Psychology, 1955, 48, 463-469.
- Sullivan, G. D., Georgeson, M. A., & Oatley, K. Channels for spatial frequency selection and the detection of single bars by the human visual system. Vision Res., 1972, 12, 383-394.
- Tolhurst, D. J. Adaptation to square-wave gratings: inhibition between spatial frequency channels in the human visual system. J. Physiol., Lond., 1972, 226, 231-249.
- Veniar, F. A. Difference thresholds for shape distortion of geometrical squares. J. Psychol., 1948, 26, 461-476.
- Virsu, V. Dark adaptation shifts apparent spatial frequency. Vision Res., 1974, 14, 433-435.

Weisstein, N. & Bisaha, J. Gratings mask bars and bars mask gratings: Visual frequency response to aperiodic stimuli. Science, 176, 1047-1049.

APPENDIX A: Terms describing a grating

APPENDIX A

The following definitions are necessary to completely describe a grating:

The spatial frequency of a grating can be expressed in terms of the number of cycles of change in luminance per unit distance. In the case of the human optical system, the unit of distance is taken to be degree of visual angle. Thus, a high frequency grating would contain many cycles per degree (c/d) and looks like many narrow alternating bright and dark bars. A low frequency grating might contain just a few wide bright and dark bars.

Spatial period = $1/\text{spatial frequency}$. Therefore, the spatial period is the visual angle subtended by a single cycle of change in luminance of the grating. Likewise, $1/2$ period would be the visual angle subtended by a single bar, bright or dark.

I max = luminance of the brightest part of the grating

I min = luminance of the dimmest part of the grating

$$\text{Average Luminance} = \frac{I_{\text{max}} + I_{\text{min}}}{2}$$

$$\text{Contrast} = \frac{I_{\text{max}} - I_{\text{min}}}{I_{\text{max}} + I_{\text{min}}}$$

Harmonics = multiples of the fundamental frequency component in stimuli other than a sine wave

Octave = One octave is a change in frequency by a factor

of two.

Duty Cycle = the percent of the distance subtended by luminance above the average luminance, within a single cycle of change in the luminance distribution. If the bright and dark bars in a grating are of equal widths, the grating has a 50% duty cycle. If the bright bars are wider than the dark bars, the duty cycle is greater than 50%.