

2008

2008 Chart Book: Insect Management At-A-Glance

Anne L. Averill

University of Massachusetts Amherst Cranberry Station, averill@eco.umass.edu

Martha M. Sylvia

University of Massachusetts Amherst Cranberry Station, martys@umass.edu

Follow this and additional works at: <https://scholarworks.umass.edu/cranchart>



Part of the [Agriculture Commons](#), and the [Plant Sciences Commons](#)

Averill, Anne L. and Sylvia, Martha M., "2008 Chart Book: Insect Management At-A-Glance" (2008). *Cranberry Chart Book - Management Guide*. 85.

Retrieved from <https://scholarworks.umass.edu/cranchart/85>

This Public Service and Outreach is brought to you for free and open access by the Cranberry Station Outreach and Public Service Activities at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Cranberry Chart Book - Management Guide by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

INSECT MANAGEMENT 2008 AT A GLANCE

Prepared by Anne L. Averill and Martha M. Sylvia

**University of Massachusetts, Cranberry Station
PO Box 569, East Wareham, MA 02538**

ALWAYS READ THE LABEL and REFER TO 2008 CRANBERRY CHART BOOK

PRACTICE	RATE/A	INSECT TARGETED	NOTES
Actara*	2-4 oz	cranberry weevil flea beetle suppression	Do not exceed 12 oz per season, low rate works well, beware bee toxicity. No Zone II in spring.
Admire Pro Admire 2F, Alias 2F	7-14 oz 16-32 oz	striped colaspis, oriental beetle	Best against just hatched larvae. Use drench application.
Avault	6 oz	cranberry weevil blackheaded fireworm	Only for spring weevil populations. No flow through bogs, hold water 1 day.
<i>Bacillus thuringiensis</i> based products		gypsy moth, armyworms cutworms, spanworms	Best in low gallonage, against early stage larvae, in well-timed systems.
Delegate WG SpinTor 2SC	3-6 oz 6-10 oz	blackheaded fireworm <i>Sparganothis</i> fruitworm cutworms, spanworms flea beetle suppression**	Delegate is more active and has greater residual than SpinTor. Better efficacy with shorter rinse times. Delegate only.
Diazinon 50 W* Diazinon AG500* Diazinon AG600* & 5W	4-6 lb 4-6 pt 51-76.5 oz	fireworms, cranberry fruitworm cutworms**, flea beetle** striped colaspis**	<i>For all diazinon formulations:</i> <i>Highly toxic to birds.</i> Hold water at least 3 days, 7 day PHI, 5 day REI, 3 apps/season
Diazinon 14G*	21 lb	cranberry girdler	SLN 2008 only, limited availability.
Flood, fall		cranberry girdler black vine & strawberry root weevil cranberry fruitworm	Sept. 20-30, hold 6 days no later than Sept. 25. 3 days may suffice if water is warm. 10-14 days after harvest; best if water is warm. Sept. 20-30, may impact vines.
Flood, late water		CB fruitworm, southern red mite gypsy moth and cutworms	April 15-20, hold for 30 days.
Flood, summer		cranberry root & white grub	<i>Loss of crop</i> , May 12-July 20.
Imidan 70W	1.33-4 lbs	fireworms, CB fruitworm	Little or varied efficacy data available in MA.
Intrepid 2F* Confirm 2F	10-16 oz 16 oz	blackheaded fireworm, cutworms ** <i>Sparganothis</i> fruitworm gypsy moth, spanworms Cranberry fruitworm**	Insect growth regulators. Intrepid is more active and has greater residual than Confirm. However, Intrepid is Zone II restricted. Very low gallonage only for fruitworm.
Lorsban 4E*, Nufos 4E* & Hatchet*, Chlorpyrifos 4E AG* Lorsban 75 WG*	3 pt 2 lbs	cranberry weevil, cutworms fireworms, spanworms, <i>Sparganothis</i> & CB fruitworms	<i>Extremely toxic to fish.</i> Hold water at least 5 days, use only 2 times/season, 60 day PHI, 24 hr REI. Most populations of weevil and <i>Sparganothis</i> are resistant to Lorsban.
Nematodes		black vine & strawberry root weevil cranberry girdler	Apply in early evening in May and/or Sept. Apply 2 weeks after end of moth flight.
Nexter	3.5-7.0 oz	southern red mite	Low-end rates usually gives excellent control,
Orthene 97, Acephate 97UP Orthene 75S, Acephate 75 WSP 1 1/3 lb	1 lb	gypsy moth, fireworms, spanworms, cutworms	Only 1 app per season, 90 day PHI, beware bee toxicity. 24 hr REI
Pyronyl or pyrenone	up to 12 oz	spanworms	
Sevin XLR Plus Sevin 4F & Carbaryl 4L Sevin 80 WSP & 80S	1.5-2 qt 1.5-2 qt 1.88-2.5 lb	fireworms, cutworms, gypsy moth, CB fruitworm CB flea beetle, striped colaspis	Avoid application within 10 days of start of bloom, 7 day PHI, no more than 5 app/season, 7 day application interval.

* Restricted Use ** FIFRA 2ee recommendation