

Spring 3-2016

2016 Chart Book: Resistance Management

Martha Sylvia

University of Massachusetts Amherst Cranberry Station, martys@umass.edu

Follow this and additional works at: <https://scholarworks.umass.edu/cranchart>



Part of the [Agriculture Commons](#), and the [Plant Sciences Commons](#)

Sylvia, Martha, "2016 Chart Book: Resistance Management" (2016). *Cranberry Chart Book - Management Guide*. 217.
Retrieved from <https://scholarworks.umass.edu/cranchart/217>

This Public Service and Outreach is brought to you for free and open access by the Cranberry Station Outreach and Public Service Activities at ScholarWorks@UMass Amherst. It has been accepted for inclusion in Cranberry Chart Book - Management Guide by an authorized administrator of ScholarWorks@UMass Amherst. For more information, please contact scholarworks@library.umass.edu.

4 RESISTANCE MANAGEMENT

RESISTANCE MANAGEMENT 2016

Prepared by Martha M. Sylvia

In an effort to manage resistance with our pesticides, many labels now come with a “group” number assigned to them. The group ID is specific among insecticides, herbicides and fungicides. Many of our cranberry pesticides are in their own group. The largest groups with the same ID are the organophosphates and neonicotinoids. The following 3 pages show the groupings for our cranberry pesticides. The goal in resistance management is to not repeatedly use compounds that fall within the same group. Resistance management may include alternating products with different modes of action or limiting the total number of applications per season.

Fungicide Resistance Action Committee (FRAC) (<http://www.frac.info/home>)

The group that advises for fungicide resistance is the Fungicide Resistance Action Committee (FRAC). Their goal is to prolong the effectiveness of fungicides that are likely to encounter resistance problems and to limit crop losses should resistance appear. For cranberry, Ridomil and Abound are fungicides that are at high risk for resistance development, while Indar and Proline are at medium risk. They should not be used repeatedly and should be carefully alternated with other fungicides from other groupings. See grouping of cranberry fungicides on page 6. A box like this would appear on the front of the label:

GROUP **11** FUNGICIDE

Herbicide Resistance Action Committee (HRAC) (<http://www.hracglobal.com/pages/Home.aspx>)

The Herbicide Resistance Action Committee developed a classification of herbicides according to their mode of action. A similar system to FRAC has been developed by the Weed Science Society of America (WSSA) using numbers instead of letters to designate the categories. This classification is found on a few herbicide labels, for example Callisto labels have this marking:

GROUP **27** HERBICIDE

Herbicide resistance is a world wide phenomenon with more than 215 documented cases. Selection of herbicide-resistant weed populations is often the result of the continuous use of the same herbicide or herbicides with the same mode of action. A key step in resistance management is to minimize the continuous use of herbicides with the same mode of action through rotations and combinations of products. One of the purposes of these classification systems is to make it easier for farmers and farm advisors to understand which herbicides share the same mode of action without having to actually know the biochemical basis.

In cranberry, our biggest concern is our new reliance on Callisto. Be sure to rotate other compounds into your herbicide schedule. Do not treat the same bog with Callisto year after year. See table of cranberry herbicides by grouping on page 7.

Insecticide Resistance Action Committee (IRAC) (<http://www.iraconline.org/>)

An Insecticide Resistance Action Committee (IRAC) has been formed to assemble the information for insecticides. Their goal is to manage resistance to keep agriculture sustainable. For cranberry, organophosphates and neonicotinoids have the most compounds within their group. We are reliant on several compounds in these groupings. As long as growers remember to alternate between groupings and not repeat same mode-of-action compounds over and over, we should be able to keep newer compounds viable for decades. See Cranberry Insecticides by grouping on the next page. Insecticides are grouped clearly by chemical makeup and most insecticide labels now included markings such as this:

GROUP **5** INSECTICIDE

Insecticide Resistance Action Committee (IRAC) Grouping for cranberry insecticides

IRAC GROUP	TRADE NAME	ACTIVE INGREDIENT	MODE OF ACTION	CHEMICAL FAMILY
1	Diazinon	diazinon	Acetylcholine esterase inhibitor	Organophosphates and carbamates
	Imidan	phosmet		
	Lorsban	chlorpyrifos		
	Orthene	acephate		
	Sevin	carbaryl		
3	Pyreth-It Pyganic	pyrethrin	Sodium channel modulators	Pyrethrins
4, 4A	Actara	thiamethoxam	Nicotinic Acetylcholine receptor agonists	Neonicotinoids
	Admire	imidacloprid		
	Assail	acetamiprid		
	Belay	clothianidin		
	Scorpion	dinotefuran		
5	Delegate	spinetoram	Nicotinic Acetylcholine receptor allosteric activators	Spinosyns
	Entrust	spinosad		
11	Dipel, Xentari Biobit	<i>Bacillus thuringiensis</i>	Microbial disruptors of insect midgut membranes	<i>Bacillus thuringiensis</i>
15	Rimon	novaluron	Inhibitors of chitin biosynthesis	Benzoylureas
18	Confirm	tebufenozide	Ecdysone agonists / molting disruptors	Diacylhydrazines
	Intrepid	methoxyfenozide		
21	Nexter	pyridaben	Mitochondrial complex / electron transport inhibitor	Meti acaracides
22	Avaunt	indoxacarb	Voltage-dependent sodium channel blockers	Oxadiazines
23	Oberon	spiromesifen	Inhibitors of acetyl CoA carboxylase	Tetramic acid derivatives
28	Altacor	chlorantraniliprole	Ryanodine receptor modulators	Diamides

6 RESISTANCE MANAGEMENT

Fungicide Resistance Action Committee (FRAC) Grouping for cranberry fungicides

FRAC GROUP	TRADE NAME	COMMON NAME	MODE OF ACTION	GROUP NAME	CHEMICAL GROUP	Resistance Development Risk
4	Metastar	mefenoxam	A1: RNA polymerase I	PA – fungicides (PhenylAmides)	acylalanines	High Risk
	Ridomil	metalaxyl				
	Ultra					
	Flourish					
11	Abound	azoxystrobin	C3: cytochrome bc1 at Qo site	QoI-fungicides	methoxy-acrylates	High Risk
	Aftershock	fluoxastrobin		Strobilurins	dihydro-dioxazines	
	Evito					
3	Indar	fenbuconazole	G1: c14-demethylase in sterol biosynthesis	DMI-fungicides (DeMethylation Inhibitors)	triazoles	Medium Risk
	Proline	prothioconazole				
19	OSO	polyoxin	H4: chitin synthase	polyoxins	peptidyl pyrimidine nucleoside	Medium Risk
	Ph-D					
33	Aliette	fosetyl-Al	Unknown	phosphonates	ethyl phosphonates	Low Risk
	Legion	aluminum-tris				
	Fosphite	phosphorous acids and salts				
	Fungi-Phite					
K-Phite, Phostrol, ProPhyt, Rampart						
M1	Champ	copper (salts)	M1: Multi-site contact activity	inorganic	inorganic	Low Risk
	Kocide					
M3	Ferbam	ferbam	M3: Multi-site contact activity	dithiocarbamates	dithiocarbamates	Low Risk
	Manzate	mancozebs		EBDC's (Ethylene bis dithio carbamate)		
	Dithane					
	Penncozeb					
M5	Bravo	chlorothalonil	M5: Multi-site contact activity	chloronitriles	chloronitriles	Low Risk
	Chloronil					
	Echo, Equus					
	Initiate					

Herbicide Resistance Action Committee (HRAC) Grouping for cranberry herbicides
Group numbering from Weed Science Society of America (WSSA) at right

HRAC GROUP	TRADE NAME	ACTIVE INGREDIENT	MODE OF ACTION	CHEMICAL FAMILY	WSSA GROUP
A	Fusilade	fluazifop-P-butyl	Inhibition of acetyl CoA carboxylase (ACCase)	Aryloxyphenoxy-propionate 'FOPs'	1
	Select	clethodim		Cyclohexanedione	
	Poast	sethoxydim		'DIMs'	
F1	Evital	norflurazon	Bleaching: Inhibition of carotenoid biosynthesis at the phytoene desaturase step (PDS)	Pyridazinone	12
F2	Callisto	mesotrione	Bleaching: Inhibition of 4-hydroxyphenyl-pyruvate-dioxygenase (4-HPPD)	Triketone	27
G	Roundup	glyphosate	Inhibition of EPSP synthase	Glycine	9
K3	Devrinol	napropramide	Inhibition of VLCFAs (Inhibition of cell division)	Acetamide	15
L	Casoron	dichlobenil	Inhibition of cell wall (cellulose) synthesis	Nitrile	20
	Quinstar	quinclorac		Quinoline carboxylic acid	26
O	Quinstar	quinclorac	Action like indole acetic acid (synthetic auxins)	Quinoline carboxylic acid	4
	2,4-D Weedar 64	2,4-D		Phenoxy-carboxylic acid	
	Stinger	clopyralid		Pyridine carboxylic acid	