



University of
Massachusetts
Amherst

2008 Chart Book: Weed Management At-A-Glance

Item Type	public;article
Authors	Sandler, Hilary A.
Download date	2025-02-07 10:45:11
Link to Item	https://hdl.handle.net/20.500.14394/9497

WEED MANAGEMENT 2008 AT A GLANCE

Prepared by Hilary A. Sandler

University of Massachusetts Cranberry Station
PO Box 569, East Wareham, MA 02538

ALWAYS READ THE LABEL and REFER TO 2008 CHART BOOK

WEEDS CONTROLLED	TIMING	HERBICIDE
	FEBRUARY AND MARCH	
Broadleaf perennials (DO NOT USE ON BOG)		Crossbow/ Weedone CB
Green scum (algae)		Copper sulfate/algaecide products
	MARCH TO MID-APRIL	
Preeemergence herbicides should be applied prior to vines breaking dormancy. Timing may vary depending on weather conditions.		
Aster, blue joint, bracken fern, corn grass, cottongrass, crabgrass, cutgrass, dulichium ferns (bracken and royal), fireweed, haircap moss, loosestrife, mannagrass, marsh St. John's wort, mare's tail, mud rush, narrow-leaved goldenrod, needlegrass, nutgrass, pitchfork, plantain, ragweed, rattlesnakegrass, royal fern, rushes, smartweed, sorrel, summergrass, velvetgrass, wild strawberry, woolgrass, warty panicgrass		Casoron
Wild bean (if active, spot-treat with Stinger – see below for probable timing)		Casoron, then Devrinol
Clover		Stinger
Dodder		Casoron (< 50 lb/A)
Pitchforks, ragweed, summergrass		Princep or Caliber
Sphagnum moss		Iron sulfate
Barnyardgrass, broadleaf panicgrass, broomsedge, Canada rush, corn grass, cutgrass, dulichium, needlegrass, nut sedge, smokegrass, spike rush, summergrass, switchgrass, woolgrass		Evital
Dewberries (brambles)		Late water or fall flood
Barnyardgrass, corngrass, cutgrass, nutgrass, povertygrass, soft rush, summergrass, warty panicgrass		Devrinol
Bog renovation - all weeds		Vapam or Basamid
	JUNE THROUGH AUGUST	
Cinnamon fern, feather fern, sensitive fern		Iron sulfate and salt
Asters, cinquefoil, feather fern, marsh St. John's wort, sensitive fern		Iron sulfate
Actively growing, true grasses (annual and perennial) on producing bogs		Poast or Select Max
Ditch weeds (sprays)		Rodeo + nonionic surfactant; Roundup (Suppl. label needed w/ Roundup Ultra)
Water weeds (but not in ditches)		Diquat
Wild bean		Stinger or salt solution
Grasses on non-producing bogs Poast and Select may be used on producing bogs		Fusilade, Poast, or Select Max + surfactant
Tall weeds on bogs		Roundup, Weedar 64
	IN THE FALL AFTER HARVEST	
Aster, blue joint, cottongrass, cutgrass, fresh meadowgrass, loosestrife, marsh St. John's wort, mud rush, narrow-leaved goldenrod, needlegrass, nutgrass, ragweed, sphagnum moss, summergrass, woolgrass		Casoron Stinger vs. composite weeds
Barnyardgrass, Canada rush, corngrass, dulichium, needlegrass, nutgrass, summergrass, switchgrass	Evital	
Barnyardgrass, corn grass, cutgrass, nutgrass, povertygrass, spike rush		Devrinol
Bog renovation - all weeds		Vapam or Basamid
Pitchforks, ragweed, summergrass		Princep or Caliber
Spot-treatment with spray applications (if using RoundUp Ultra, need supplemental labels)		RoundUp WeatherMAX