



University of  
Massachusetts  
Amherst

## Antioxidant Response Mechanism in Apples during Post-Harvest Storage and Implications for Human Health Benefits

Item Type	Thesis (Open Access)
Authors	Adyanthaya, Ishan
DOI	<a href="https://doi.org/10.7275/312284">10.7275/312284</a>
Download date	2026-04-23 03:10:36
Link to Item	<a href="https://hdl.handle.net/20.500.14394/46637">https://hdl.handle.net/20.500.14394/46637</a>

ANTIOXIDANT RESPONSE MECHANISM IN APPLES DURING  
POST-HARVEST STORAGE AND IMPLICATIONS FOR HUMAN  
HEALTH BENEFITS

A Thesis Presented

by

ISHAN ADYANTHAYA

Submitted to the Graduate School of the  
University of Massachusetts Amherst in partial fulfillment  
of the requirements for the degree of

MASTER OF SCIENCE

September 2007

Molecular and Cellular Biology

ANTIOXIDANT RESPONSE MECHANISM IN APPLES DURING  
POST-HARVEST STORAGE AND IMPLICATIONS FOR HUMAN  
HEALTH BENEFITS

A Thesis Presented

by

ISHAN ADYANTHAYA

Approved as to style and content by:

---

Kalidas Shetty, Chair

---

Sallie Smith-Schneider, Member

---

Young-Cheul Kim, Member

---

Patricia Wadsworth, Director  
Molecular and Cellular Biology Program

## ACKNOWLEDGMENTS

I would like to thank Dr. Kalidas Shetty for giving me the opportunity to work in his lab. I am deeply indebted for his constant support and advice throughout my undergraduate and graduate career here at UMass. The opportunity to work in his lab has been an incredible learning experience of which I am sure will help me in the future. I would also like to thank both Young-In Kwon and Emmanouil Apostolidis not only for their never ending help and support in the lab but also their friendship. From my very first day in the lab Young has been an excellent guide with limitless patience. His constant insights and directions have made my every task, whether running and assay, writing a paper or making a presentation, much easier for which I will be eternally grateful. Both Manos and Young have made my life in lab not just easy but enjoyable. Finally I would like to thank my family for their constant support throughout my college career.

## TABLE OF CONTENTS

	Page
ACKNOWLEDGMENTS .....	iii
LIST OF TABLES .....	vi
LIST OF FIGURES .....	vii
CHAPTER	
1. INTRODUCTION .....	1
2. LITERATURE REVIEW .....	4
2.1. Phenolics: Synthesis, Functions and Health Benefits .....	4
2.2. Diabetes .....	9
2.3. Role of Free Radical Scavenging and Enzyme Antioxidant Activity in Post-Harvest Preservation of Fruits .....	11
2.4. Proline Linked Pentose Phosphate Pathway for Effective Antioxidant Activity and Phenolic Synthesis Enhancing Post-Harvest Preservation.....	15
3. OBJECTIVES.....	19
4. APPLE POST-HARVEST PRESERVATION LINKED TO PHENOLICS AND SOD ACTIVITY.....	20
4.1. Abstract.....	20
4.2. Introduction.....	21
4.3. Materials and Methods.....	25
4.3.1. Phenolic Extraction.....	25
4.3.2. Total Phenolics Assay.....	25
4.3.3. Antioxidant Activity by 1, 1-Diphenyl-2-Picrylhydrazyl Radical (DPPH) Inhibition Assay .....	26
4.3.4. Enzyme Extraction.....	26
4.3.5. Total Protein Assay.....	27
4.3.6. Superoxide Dismutase (SOD) Assay.....	27
4.3.7. HPLC Analysis of Proline .....	28
4.3.8. Proline Dehydrogenase (PDH) Assay.....	28
4.3.9. Succinate Dehydrogenase (SDH) Assay.....	29
4.3.10. Glucose-6-Phospahte Dehydrogenase (G-6-PDH) Assay .....	29
4.3.11. Guaiacol Peroxidase (GPX) Assay .....	30
4.3.12. Statistical Analysis.....	30
4.4. Results and Discussion .....	31

4.4.1. Phenolic-Linked Antioxidant Activity and Visual Observations .....	31
4.4.2. Superoxide Dismutase (SOD) Activity .....	35
4.4.3. Proline Linked Pentose Phosphate Pathway (PLPPP) for Energy and Phenolic Synthesis .....	37
<b>5. HEALTH BENEFITS OF APPLE PHENOLICS FROM POST-HARVEST STAGES FOR POTENTIAL DIABETES MANAGEMENT USING IN VITRO MODELS .....</b>	<b>45</b>
5.1. Abstract.....	45
5.2. Introduction.....	46
5.3. Materials and Methods.....	49
5.3.1. Phenolic Extraction.....	49
5.3.2. Total Phenolics Assay.....	49
5.3.3. Antioxidant Activity by 1, 1-Diphenyl-2-Picrylhydrazyl Radical (DPPH) Inhibition Assay .....	50
5.3.4. $\alpha$ -Glucosidase Inhibition Assay .....	50
5.3.5. $\alpha$ -Amylase Inhibition Assay.....	51
5.3.6. Statistical Analysis.....	52
5.4. Results and Discussion .....	52
5.4.1. Visual Observations.....	52
5.4.2. Phenolic Content of Apple.....	52
5.4.3. Antioxidant Activity .....	53
5.4.4. $\alpha$ -Glucosidase and $\alpha$ -Amylase Inhibition .....	54
<b>6. BIBLIOGRAPHY.....</b>	<b>67</b>

## LIST OF TABLES

Table	Page
1. Phenolic content and antioxidant activity of apples over 3 months .....	32

## LIST OF FIGURES

Figure	Page
1. Biosynthesis of phenolic phytochemicals.....	7
2. Common simple phenol, biphenyls, flavanoids and tannins in plants.....	8
3. Oxidative stress related hyperglycemia and complications.....	10
4. Reactive oxygen species and cellular homeostasis.....	13
5. Biological mechanisms of ROS formation and detoxification reactions.....	14
6. Proline synthesis coupled to the pentose phosphate pathway.....	16
7. Proline-linked pentose phosphate pathway for phenolic synthesis and efficient antioxidant response.....	18
8. Sample phenolic concentration ( $\mu\text{g/ml}$ ) and antioxidant activity.....	33
9. Visual observation of post-harvest changes in apples over 3 months.....	34
10. Superoxide dismutase specific activity (U/mg protein).....	36
11. Proline content of apples over 3 months ( $\mu\text{mol/mg protein}$ ).....	40
12. Proline dehydrogenase specific activity in apples over 3 months. (U/mg protein).....	41
13. Succinate dehydrogenase activity in apples over 3 months (nmol/mg protein).....	42
14. Glucose-6-phosphate dehydrogenase specific activity in apples over 3 months (nmol/mg protein).....	43
15. Guaiacol peroxidase specific activity in apples over 3 months (nmol/mg protein).....	44
16. Visual observation of post-harvest changes in apples over 3 months.....	57
17. Total phenolic content of apples water extraction ( $\mu\text{g/g FW}$ ).....	58
18. Total phenolic content of apples 12% ethanol extraction ( $\mu\text{g/g FW}$ ).....	59
19. DPPH inhibition. Apple water extracts.....	60
20. DPPH inhibition. Apple 12% ethanol extracts.....	61

21. $\alpha$ -Glucosidase inhibition. Apple water extracts .....	62
22. $\alpha$ -Glucosidase inhibition. Apple 12% ethanol extracts .....	63
23. Sample phenolic concentration ( $\mu\text{g/ml}$ ) and $\alpha$ -glucosidase inhibition. Apple peel water extracts .....	64
24. Sample phenolic concentration ( $\mu\text{g/ml}$ ) and $\alpha$ -glucosidase inhibition. Apple peel 12% ethanol extracts .....	65
25. Correlation between sample phenolic concentration ( $\mu\text{g/ml}$ ) and $\alpha$ -glucosidase inhibition .....	66

# CHAPTER 1

## INTRODUCTION

Phenolic phytochemicals are secondary metabolites synthesized by plants, which constitute an important part of the diet in both humans and animals (1-3). In plants these phenolics exhibit protective functions against environmental and biological stress such as high energy radiation exposure, bacterial infection or fungal attacks (4). In addition phenolics are also important for cell structure, signaling and pigmentation. Due to diversity of functions, phenolic phytochemicals are expressed in diverse cell and tissue types (4-6). Since phenolics have been linked to many protective functions in plants it is likely they play similar protective roles in fruits; making them a key factor in post-harvest preservation of fruits.

In addition growing evidence suggests that the intake of phenolics via fruits and vegetables is linked to reduced risk of chronic diseases like diabetes, cardiovascular disease and cancer (7-11). A study with many popular fruits found apples to contain the second highest total phenolics content after cranberries, but the highest soluble phenolics (12). Apples also had the second highest free radical scavenging-linked antioxidant activity among the fruits investigated (12). Apples are a popular fruit all over the world and have the potential to contribute as a significant source of phenolic antioxidants. In the United States, 22% of phenolic intake via fruit comes from apples (13). In Finland the main source of dietary phenolics are onions and apples and in the Netherlands tea, onions and apples are the biggest sources of phenolics (7, 9). Since the emerging oxidation-linked diseases can benefit from high intake of fruits and vegetables, high phenolic

antioxidant-containing apples have promise for enhancing human health through cellular protective functions.

There are more than 180 million people worldwide who have diabetes and 90% of these suffer from Non-insulin-dependant Type II diabetes (14). The World Health Organization (WHO) estimates that the number of people with diabetes will double by 2030 (14). The WHO states that Type II diabetes can be prevented by physical activity, healthy eating and prevention of obesity (14). In addition WHO figures show that about 80% of people with Type II diabetes are from low and middle income countries (14). Therefore there is a need for low cost and easily available methods for management of diabetes in order to improve the standard of living of most diabetes patients.

A number of previous studies have found phenolics from many common foods like capsicum, cinnamon and fenugreek to have the relevant phytochemical profile of,  $\alpha$ -glucosidase inhibition and low  $\alpha$ -amylase inhibition coupled with free radical scavenging-linked antioxidant activity, for potential diabetes management (15-17). This offers the potential for good postprandial blood glucose management via  $\alpha$ -glucosidase inhibition without the common side-effects associated with high  $\alpha$ -amylase inhibition (15-17). In addition these same foods have free radical scavenging-linked antioxidant activity which can help maintain the redox balance in susceptible cells (15-17). Since apple is a common fruit with no known side effects, any  $\alpha$ -glucosidase inhibiting effects, if found, are promising for Type II diabetes management.

Based on the above background this study explored the potential dual benefit of apple phenolics for better post-harvest preservation and health benefits. Specifically phenolic-linked changes were investigated during post-harvest storage of four varieties of

apple over three months. The objective was to determine the changes in phenolic content over the storage period and its relevance to post-harvest preservation and concurrently determine any anti-diabetes-linked health benefits that could be attributed to the phenolic content. The health relevant parameters investigated were *in vitro* antioxidant activity and inhibition of  $\alpha$ -glucosidase and  $\alpha$ -amylase relevant for glycemic index modulation. In addition understanding of how inducible phenolics and related antioxidant activity (both free radical and enzyme-linked) are coupled to the pentose phosphate pathway with positive consequences for post-harvest preservation of apples was investigated. This was done by evaluating antioxidant enzyme activity, proline content, phenolic content, free radical scavenging antioxidant activity and activity of key enzymes over a 3 month post-harvest storage period. The enzymes evaluated were glucose-6-phosphate dehydrogenase (G6PDH), succinate dehydrogenase (SDH), proline dehydrogenase (PDH), guaiacol peroxidase (GPX), superoxide dismutase (SOD) and catalase (CAT).

## CHAPTER 2

### LITERATURE REVIEW

#### **2.1. Phenolics: Synthesis, Functions and Health Benefits**

Phenolic phytochemicals are synthesized by a common biosynthetic pathway which incorporates precursors from both the shikimate and/or the acetate-malonate pathways (1, 18). The first step in the synthesis of phenolic phytochemicals is the commitment of glucose to the pentose phosphate pathway (PPP), converting glucose-6-phosphate irreversibly to ribulose-5-phosphate. This first committed step in the conversion to ribulose-5-phosphate is carried out by glucose-6-phosphate dehydrogenase (G6PDH). The conversion to ribulose-5-phosphate also produces reducing equivalents (NADPH) for cellular anabolic reactions (Fig. 1). PPP also generates erythrose-4-phosphate which along with phosphoenolpyruvate, from glycolysis, is channeled to the shikimate pathway to produce phenylalanine, which is directed through the phenylpropanoid pathway to produce phenolic phytochemicals (1, 6, 19-21); (Fig.2).

These phenolic phytochemicals have been shown to have a wide array of functions in plants. In plants biotic and abiotic stress has been shown to stimulate secondary metabolite synthesis which results in the production of phenolics (22, 23). Ozone exposure has been shown to increase transcript levels of enzymes involved in the phenolic synthesis and lignin pathways (24). Studies have linked increase in thermo-tolerance during hyperthermia to accumulation of heat shock proteins and phenolic metabolites (25). Many phenolics are induced in response to infection, wounding, nutritional stress, cold stress and UV irradiation (22, 23, 26-30). Phenolics can function as effective antioxidants by scavenging singlet oxygen and free radicals via their ability


to donate hydrogens from hydroxyl groups positioned around the aromatic ring (10, 31-34). A recent study of antioxidant activity in apples by Lee *et al.* 2003 (35) found that flavonoids like quercetin, epicatechin and procyanidin B<sub>2</sub> rather than vitamin C contributed significantly to total antioxidant activity (35).

In light of the vast array of cellular protective functions phenolics have in plants it is not surprising therefore that phenolics have diverse medicinal properties for human health applications. For example curcumin from *Curcuma longa* and *Curcuma mannga* and rosmarinic acid from *Rosmarinus officinalis* are used as antioxidants and anti-inflammatory compounds (36-41). Also lithospermic acid from *Lithospermum sp.* is used as antigonadotropic agent, proanthocyanidins from cranberry can be used to treat urinary tract infections and anethole from *Pimpinella anisum* is used as an antifungal agent (42-45).


Phenolic phytochemicals have also been found to have potential in the management of oxidative stress linked chronic diseases like diabetes, cancer and cardiovascular disease (5, 20, 46-48). In a study involving Empire apples found phenolics from the apples to have potential cancer chemopreventive activity due to their combined antioxidant and anti-tumor promoting activities (49). Similarly an epidemiological study by Knekt *et al.* 1997 (7) found that intake of dietary flavanoids showed an inverse association with lung cancer incidence (7). Another epidemiological study which examined the relationship between fruit and vegetable intake and the incidence of different cancers found that for most cancers the increase in risk for people with low fruit intake in the study was about twice as high as people with high intake (50). *In vitro* anti-proliferation studies with HepG2, human liver-cancer cells have shown phenolics from

many common fruits like cranberry, lemon, apples and strawberries to inhibit proliferation (12).

An epidemiological study by Knekt *et al* in 2001 (8) of more than 10,000 Finnish men and women showed intake of foods containing flavanoids was associated with lower risk of Type II diabetes. The study indicated the lower risk was associated with high intake of quercetin. The same study also states the strongest association between high flavonoid intake with lower risk for Type II diabetes was seen when the source of flavonoids was from apples and berries (8). Therefore a diet containing apples has the potential to reduce risk of Type II diabetes. Since apple is a common fruit with no known side effects, any  $\alpha$ -glucosidase inhibiting effects, if found, are promising for Type II diabetes management.



**Figure 1.** Biosynthesis of phenolic phytochemicals.



**Figure 2.** Common simple phenol, biphenyls, flavanoids and tannins in plants.


## 2.2. Diabetes

Hyperglycemia has been linked to the onset of insulin independent Type II diabetes (51). A good strategy for management of Type II diabetes is inhibition of enzymes that hydrolyze dietary polysaccharides in the gut which can significantly reduce the rise in blood sugar levels after a meal by reducing the absorption of monosaccharides by the enterocytes of the small intestine (52, 53). Enzymes that hydrolyze dietary polysaccharides and modulate gut absorption are pancreatic  $\alpha$ -amylase and  $\alpha$ -glucosidases (54, 55). However a common side effect of these enzyme inhibitory drugs like acarbose is the excessive inhibition of pancreatic  $\alpha$ -amylase which can result in abdominal distention, flatulence and diarrhea (56, 57). These side effects are caused by abnormal fermentation of unhydrolyzed polysaccharides by gut bacteria (56, 58). Therefore in order to reduce these side effects but still manage hyperglycemia a high  $\alpha$ -glucosidase inhibition and low  $\alpha$ -amylase inhibition is beneficial.

Acute complications in patients suffering from Type II diabetes are generally hyperglycemia-induced metabolic problems and infection. Long term effects of hyperglycemia are microvascular complications like nephropathy, diabetic neuropathy, sexual dysfunction and retinopathy and the macrovascular complication of hypertension (59); (Fig. 3). Recent studies have shown that hyperglycemia triggers generation of free radicals in mesangial cells in the renal glomerulus, neuron cells in peripheral nerves and capillary endothelial cells in the retina (60). The generation of free radicals in all these cells types causes oxidative stress which can be the cause of microvascular complications generally linked with hyperglycemia (60). Most cells are capable of reducing glucose transport inside the cell in hyperglycemic conditions so as to maintain a constant internal

glucose level, however the cells usually damaged by hyperglycemia were found to be inefficient in keeping their internal glucose levels constant (61, 62). Therefore it is not only important to control postprandial hyperglycemia but also keep in check any cellular redox imbalances to prevent diabetic complications.

In addition another study has shown quercetin, which is a phenolic abundant in apples, to be an aldose reductase (alditol: NADP+ oxidoreductase) inhibitor (63). Aldose reductase is the first enzyme of the polyol pathway. Glucose metabolism through the polyol pathway has been linked long-term diabetic complication like cataract, nephropathy, neuropathy and retinopathy (63).



**Figure 3.** Oxidative stress related hyperglycemia and complications.


### **2.3. Role of Free Radical Scavenging and Enzyme Antioxidant Activity in Post-Harvest Preservation of Fruits**

Reactive oxygen species (ROS) like superoxide radicals ( $O_2^{\cdot-}$ ), hydroxyl radicals ( $\cdot OH$ ) and hydrogen peroxide ( $H_2O_2$ ) are the products of oxidative dysfunctional biochemical reactions within cells. These ROS when left unchecked cause oxidative damage resulting in lipid peroxidation, protein denaturation and mutagenesis. An increase in ROS is linked to different types of stress such as drought, heat stress, metal toxicity, radiation exposure, pathogens and salinity (64-70); (Fig. 4). Further ROS is also involved in natural and induced senescence and cell death in plants (71-74). For example, studies have indicated increase in hydroperoxides during pepper, banana, pear and tomato ripening during which senescence is induced (75-77).


Plants counter harmful effects of ROS with antioxidants metabolites and enzymes. Antioxidants metabolites include; water soluble compounds like ascorbate, glutathione and flavonoids and lipid soluble compounds like carotenoids and tocopherols. Enzymes linked to antioxidant response include superoxide dismutase (SOD), catalase (CAT), monodehydroascorbate reductase (MDHAR), dehydroascorbate reductase (DHAR), ascorbate peroxidase (ASPX) and glutathione reductase (GR). SOD converts  $O_2^{\cdot-}$  to  $H_2O_2$  which is then reduced to  $H_2O$  by CAT or ASPX depending cellular localization (78-80); (Fig. 5). Both ascorbate and glutathione can also interact directly with and scavenge ROS (79). Oxidized ascorbate is reduced by DHAR, MDHAR and glutathione and oxidized glutathione is reduced by GR. Reduction of both oxidized glutathione and ascorbate by their respective enzymes is NADPH dependant (20). Studies have shown many phenolic compounds especially flavonoids e.g., quercetin, rutin and

catechin have free radical scavenging antioxidant activity (49, 81). In addition, studies in fava beans and peas have shown that even exogenous phenolics can stimulate antioxidant enzyme activity (6, 82).

Therefore, since ROS is involved in plant development, including fruit ripening and senescence, antioxidant response coupling phenolic synthesis and antioxidant enzyme response may be recruited to counter ROS and senescence (70, 80, 83-85). Previous studies with muskmelon fruits and sunflower seeds indicated that delayed senescence in specific tissue types correlated to high antioxidant enzyme response (86, 87). In order to couple cellular antioxidants like ascorbate, glutathione and phenolic phytochemicals with antioxidant enzymes for effective antioxidant response cellular reducing equivalents such as  $FADH_2$  and NADPH are required. Studies have found that reduced reductant levels increased the rate of senescence in a number of herbaceous species (73, 88).



**Figure 4.** Reactive oxygen species and cellular homeostasis.




**Figure 5.** Biological mechanisms of ROS formation and detoxification reactions (From Sigma-Aldrich website).

#### **2.4. Proline Linked Pentose Phosphate Pathway for Effective Antioxidant Activity and Phenolic Synthesis Enhancing Post-Harvest Preservation**

Another commonly seen stress response in plants is an increase in the synthesis of proline. Studies have shown proline biosynthesis to be stimulated in cases of water stress and salinity stress (89-93). Proline synthesis and accumulation is also seen during senescence, freezing tolerance, water stress, salt stress and dehydration in plant cells (94-99). It has been suggested that proline protects membranes in times of stress like those mentioned above (100, 101).

An alternative model for coupling proline synthesis with the PPP has been proposed where proline biosynthesis in response to stress can manage energy and reductant needs of anabolic pathways (20). This active metabolic role of proline could have implications for plant senescence where proline can act as an antioxidant or stimulate phenolic-linked antioxidant response (20, 97, 102). Proline is synthesized via the reduction of glutamate to  $\Delta^1$ -pyrroline-5-carboxylate (P5C) which is further reduced to proline, with both reactions using NADPH as a reductant. (20, 103, 104). Since the reduction of P5C in the cytosol requires NADPH an increase in the proline synthesis would result in a reduction in the NADPH/NADP<sup>+</sup> ratio which has been shown to activate G6PDH (105, 106). G6PDH catalyzes the first rate limiting step in the PPP, therefore it is possible that during the post-harvest storage period different stress factors induce proline synthesis which in turn stimulates the PPP (20, 107).



**Figure 6.** Proline synthesis coupled to the pentose phosphate pathway. G6PDH=Glucose-6-phosphate dehydrogenase, PDH=Proline dehydrogenase, P-5-C=  $\Delta^1$ -pyrroline-5-carboxylate,  $\alpha$ -KG =  $\alpha$ -ketoglutarate, TCA = tricarboxylic acid cycle

This stimulation of the PPP would result in more essential reducing equivalents in the form of NADPH for efficient antioxidant enzyme response. The PPP stimulation would also make more sugar phosphate precursors which along with the NADPH produced can support pathways for the synthesis of antioxidants, phenolic phytochemicals and other protective compounds (20, 107). The proline synthesized can function as an alternative reductant (in place of NADH) in mitochondrial oxidative phosphorylation to generate ATP (20, 104, 107) where proline dehydrogenase (PDH) catalyzes the first reaction of proline oxidation to P5C and linking the electron transport

chain (ETC) in the mitochondria with oxygen as the terminal electron acceptor. P5C is then hydrolyzed nonenzymatically and oxidized to glutamate by the NAD-dependent P5C dehydrogenase. Following oxidative deamination by glutamate dehydrogenase it flows into the TCA cycle through  $\alpha$ -ketoglutarate to generate NADH for oxidative phosphorylation or recycled back into the cytosol (20, 93, 108) (Fig. 6, 7).



## CHAPTER 3

### OBJECTIVES

- 1) To investigate the role of phenolic-linked antioxidant response in post-harvest preservation of apples.
- 2) To investigate the health benefits of phenolic-linked free radical scavenging activity of apple extracts for potential management of glycemic index related to Type II diabetes.

## CHAPTER 4

### APPLE POST-HARVEST PRESERVATION LINKED TO PHENOLICS AND SOD ACTIVITY

#### **4.1. Abstract**

The biochemical factors affecting post-harvest preservation in apples indicated that well-preserved varieties of apples had increased superoxide dismutase (SOD) activity initially and the activity declined during later storage as apples deteriorated. The SOD link to better preservation correlated with higher phenolic content and free-radical scavenging linked antioxidant activity. Well-preserved varieties were able to maintain a more stable pentose phosphate pathway (PPP) (measured by the activity of glucose-6-phosphate dehydrogenase, G6PDH) throughout the storage period. Proline content increased in all varieties with an increase in proline dehydrogenase (PDH) activity in the initial period indicating proline catabolism supporting potential ATP synthesis. During later storage succinate dehydrogenase (SDH) activity increased while PDH activity declined indicating a shift to tricarboxylic acid cycle and likely NADH generation for ATP synthesis. This shift coupled with the declining SOD activity coincides with rapid deterioration. The guaiacol peroxidase activity (GPX) activity generally declined in late stages indicating post-harvest deterioration.

Key words:

Apple post-harvest preservation, phenolic, antioxidant activity, superoxide dismutase, glucose-6-phosphate dehydrogenase, succinate dehydrogenase, proline dehydrogenase, guaiacol peroxidase.

## 4.2. Introduction

Phenolic enriched fruits and vegetables are important for both post-harvest preservation and human health benefits linked to the phenolic-associated antioxidant activity (109). A previous investigation with different varieties of apples indicated a positive link between phenolic content and better post-harvest preservation over 3 months. In the same study phenolic levels directly correlated with free radical scavenging antioxidant activity. Phenolic phytochemicals are secondary metabolites synthesized by plants, which constitute an important part of the diet in both humans and animals with potential health benefits (1-3). Phenolic phytochemicals are synthesized by a common biosynthetic pathway which incorporates precursors from both the shikimate and/or the acetate-malonate pathways (1, 18). The first step in the synthesis of phenolic phytochemicals is the commitment of glucose to the pentose phosphate pathway (PPP), converting glucose-6-phosphate irreversibly to ribulose-5-phosphate. This two step process also produces reducing equivalents (NADPH) for cellular anabolic reactions. PPP also generates erythrose-4-phosphate which is channeled to the shikimate pathways to produce phenylalanine, which is directed through the Phenylpropanoid pathway to produce phenolic phytochemicals (1, 6, 19-21).

Reactive oxygen species (ROS) like superoxide radicals ( $O_2^{\cdot-}$ ), hydroxyl radicals ( $\cdot OH$ ) and hydrogen peroxide ( $H_2O_2$ ) are the products of oxidative dysfunctional biochemical reactions within cells. These ROS when left unchecked cause oxidative damage resulting in lipid peroxidation, protein denaturation and mutagenesis. An increase in ROS is linked to different types of stress such as drought, heat stress, metal toxicity, radiation exposure, pathogens and salinity (64-70). Further ROS is also involved

in natural and induced senescence and cell death in plants (71-74). For example, studies have indicated increase in hydroperoxides during pepper, banana, pear and tomato ripening during which senescence is induced (75-77).

Plants counter harmful effects of ROS with antioxidant metabolites and enzymes. Studies have showed many phenolic compounds especially flavonoids e.g., quercetin, rutin and catechin have free radical scavenging antioxidant activity to counter ROS (49, 81). In addition, studies in fava beans and peas have shown that exogenous phenolics can stimulate antioxidant enzyme activity to potentially counter ROS (6, 82).

Therefore, since ROS is involved in plant development including fruit ripening and senescence antioxidant response coupling phenolics and antioxidant enzyme response may be recruited to counter ROS and senescence (70, 80, 83-85). Previous studies with muskmelon fruits and sunflower seeds indicated that delayed senescence in specific tissue types correlated to high antioxidant enzyme response (86, 87). In order to couple cellular antioxidants like ascorbate, glutathione and phenolic phytochemicals with antioxidant enzymes for effective antioxidant response cellular reducing equivalents such as FADH<sub>2</sub> and NADPH are required. Studies have found that reduced reductant levels increased the rate of senescence in a number of herbaceous species (73, 88). NADPH is regenerated by the PPP along with while synthesizing sugar phosphates, supplying precursors for phenolic biosynthesis. Therefore the enzymes of PPP play an important role in preservation of plant tissue.

An alternative model for coupling proline synthesis with the PPP has been proposed where proline biosynthesis in response to stress can manage energy and reductant needs of anabolic pathways (20). This active metabolic role of proline could

have implications for plant senescence where proline can act as an antioxidant or stimulate phenolic linked antioxidant response (20, 97, 102). Proline is synthesized via the reduction of glutamate to  $\Delta^1$ -pyrroline-5-carboxylate (P5C) which is further reduced to proline, with both reactions using NADPH as a reductant. (20, 103, 104). Since the reduction of P5C in the cytosol requires NADPH an increase in the proline synthesis would result in a reduction in the NADPH/NADP<sup>+</sup> ratio which has been shown to activate G6PDH (105, 106). G6PDH catalyzes the first rate limiting step in the PPP, therefore it is possible that during the post-harvest storage period different stress factors induce proline synthesis which in turn stimulates the PPP (20, 107). This stimulation of the PPP would result in more essential reductant equivalents in the form of NADPH for efficient antioxidant enzyme response. The PPP stimulation would also make more sugar phosphate precursors which along with the NADPH produced can support pathways for the synthesis of antioxidants, phenolic phytochemicals and other protective compounds (20, 107). The proline synthesized can function as an alternative reductant (in place of NADH) in mitochondrial oxidative phosphorylation to generate ATP (20, 104, 107) where proline dehydrogenase (PDH) catalyzes the first reaction of proline oxidation to P5C and linking the electron transport chain (ETC) in the mitochondria with oxygen as the terminal electron acceptor. P5C is then hydrolyzed nonenzymatically and oxidized to glutamate by the NAD-dependent P5C dehydrogenase. Following oxidative deamination by glutamate dehydrogenase it flows into the TCA cycle through  $\alpha$ -ketoglutarate to generate NADH for oxidative phosphorylation (20, 93, 108).

Based on the above rationale the objective of this investigation was to understand how inducible phenolics and related antioxidant activity (both free radical and enzyme-linked) are coupled to proline-linked PPP with positive consequences for post-harvest preservation of apples. This was done by evaluating antioxidant enzyme activity, proline content, phenolic content, free radical scavenging antioxidant activity and activity of key enzymes over a 3 month post-harvest storage period. The enzymes evaluated were glucose-6-phosphate dehydrogenase (G6PDH), succinate dehydrogenase (SDH), proline dehydrogenase (PDH), guaiacol peroxidase (GPX), superoxide dismutase (SOD) and catalase (CAT).

### **4.3. Materials and Methods**

The freshly picked apples were purchased from a local orchard (Clarkdale Orchards Deerfield, MA, US). Four of the popular varieties of apples were selected to be evaluated for this study; Cortland, Macintosh, Empire and Mutsu. These were stored in a cold room (5°C) for 3 months. Unless noted otherwise, all chemicals were purchased from Sigma Chemical Co. (St. Louis, MO).

#### **4.3.1. Phenolic Extraction**

Soluble phenolics were extracted from the peel and pulp of each variety separately. Phenolics were extracted using distilled water and 12% ethanol. The water extractions were done with 20 g of peel in 50 ml of water and 100 g of pulp in 50 ml of water. The ethanol extractions were done with 5 g of peel in 15 ml of 12% ethanol and 10 g of pulp in 20 ml of 12% ethanol. Apples were first peeled then weighed and mixed with either distilled water or 12% ethanol. This was then homogenized for 2 min using a blender. Resulting mix was collected and centrifuged for 5 min. Supernatant was collected and stored at -20°C. This was done every month until the 3<sup>rd</sup> month of storage.

#### **4.3.2 Total Phenolics Assay**

The total phenolic content was determined by an assay modified from Shetty *et al.* (110). Briefly, one milliliter of extract was transferred into a test tube and mixed with 1 ml of 95% ethanol and 5 ml of distilled water. To each sample 0.5 ml of 50% (v/v) Folin-Ciocalteu reagent was added and mixed. After 5 min, 1 ml of 5% Na<sub>2</sub>CO<sub>3</sub> was added to the reaction mixture and allowed to stand for 60 min. The absorbance was read at 725

nm. The absorbance values were converted to total phenolics and were expressed in micrograms equivalents of gallic acid per grams fresh weight (FW) of the sample. Standard curves were established using various concentrations of gallic acid in 95% ethanol.

#### **4.3.3. Antioxidant Activity by 1, 1-diphenyl-2-picrylhydrazyl radical (DPPH)**

##### **Inhibition Assay**

To 3 ml of 60  $\mu$ M DPPH in ethanol, 250  $\mu$ l of each extract was added, the decrease in absorbance was monitored at 517 nm until a constant reading was obtained. The readings were compared with the controls, which contained 250  $\mu$ l of 95% ethanol instead of the extract. The % inhibition was calculated by:

$$\% \text{ inhibition} = \left( \left[ \frac{A_{517}^{\text{Control}} - A_{517}^{\text{Extract}}}{A_{517}^{\text{Control}}} \right] \right) \times 100$$

#### **4.3.4. Enzyme Extraction**

For each variety of apple 2 g of pulp was ground in a mortar and pestle along with 2 ml enzyme extraction buffer. The mortar and pestle was kept cool in an ice bath during the extraction process. The sample was centrifuged at 12000xg for 15 min at 5 °C and stored on ice. The supernatant was used for further analysis.

#### **4.3.5. Total Protein Assay**

Protein content was measured by the method of Bradford assay (111). The dye reagent concentrate (Bio-Rad protein assay kit II, Bio-Rad Laboratory, Hercules, CA) was diluted 1:4 with distilled water. A volume of 5 mL of diluted dye reagent was added to 100  $\mu$ l cell extract. After vortexing and incubating for 5 min, the absorbance was measured at 595 nm against a 5 mL reagent blank and 100  $\mu$ l buffers using a UV-VIS Genesys spectrophotometer (Milton Roy, Inc., Rochester, NY).

#### **4.3.6. Superoxide Dismutase (SOD) Assay**

A competitive inhibition assay was performed that used xanthine-xanthine oxidase-generated superoxide to reduce nitroblue tetrazolium (NBT) to blue formazan. Spectrophotometric assay of SOD activity was carried out by monitoring the reduction of NBT at 560 nm (112). The reaction mixture contained 13.8 ml of 50 mM potassium phosphate buffer (pH 7.8) containing 1.33mM DETAPAC; 0.5 ml of 2.45 mM NBT; 1.7 ml of 1.8 mM xanthine and 40 IU/ml catalase. To 0.8 ml of reagent mixture 100  $\mu$ l of phosphate buffer and 100  $\mu$ l of xanthine oxidase was added. The change in absorbance at 560 nm was measured every 20 sec for 2 min and the concentration of xanthine oxidase was adjusted to obtain a linear curve with a slope of 0.025 absorbance per min. The phosphate buffer was then replaced by the enzyme extract and the change in absorbance was monitored every 20 sec for 2 min. One unit of SOD was defined as the amount of protein that inhibits NBT reduction to 50% of the maximum.

#### **4.3.7. HPLC Analysis of Proline**

High Performance Liquid Chromatography (HPLC) was performed using an Agilent 1100 liquid chromatograph equipped with a diode array detector (DAD 1100). The analytical column was a reverse phase Nucleosil C18, 250 mm x 4.6 mm with a packing material of 5  $\mu$ m particle size. The samples were eluted out in an isocratic manner with a mobile phase consisting of 20 mM potassium phosphate (pH 2.5 by phosphoric acid) at a flow rate of 1 ml/min and detected at 210 nm. L-Proline (Sigma chemicals, St. Louis, MO) dissolved in the 20 mM Potassium phosphate solution was used to calibrate the standard curve. The amount of proline in the sample was reported as  $\mu$ mol of proline per milligram of sample protein.

#### **4.3.8. Proline Dehydrogenase (PDH) Assay**

Modified method described by Costilow and Cooper (113) was used to assay the activity of proline dehydrogenase. Briefly, the enzyme reaction mixture contained 100 mM sodium carbonate buffer (pH 10.3), 20 mM L-proline solution and 10 mM NAD. To 1 ml of this reaction mixture, 200  $\mu$ l of enzyme extract was added. The increase in absorbance was followed at 340 nm for 3 min, at 32°C. The absorbance was noted at zero time and then after 3 min. In the spectrophotometric assay, one unit of enzyme activity is that amount causing an increase in absorbance of 0.01 per min at 340 nm (1.0 cm light path).

#### **4.3.9. Succinate Dehydrogenase (SDH) Assay**

Modified method described by Bregman (114) was used to assay the activity of succinate dehydrogenase. The cell suspension was diluted with 2.0 ml of cell extraction buffer. The sample was then assayed at room temperature for succinate dehydrogenase activity. The assay mixture consists of the following: 1.01 ml of 0.4 M potassium phosphate buffer (pH 7.2); 40  $\mu$ l of 0.15 M sodium succinate (pH 7.0); 40  $\mu$ l of 0.2 M sodium azide; and 10  $\mu$ l of 6.0 mg/ml DCPIP. This mixture was used to obtain baseline (zero) of the spectrophotometer reading at 600 nm wavelength. To 1.1 ml of this mixture, 200  $\mu$ l of the sample was added. The rate of change in absorbance per minute was used to quantify the enzyme in the mixture using the extinction co-efficient of DCPIP ( $19.1 \text{ mM}^{-1}\text{cm}^{-1}$ ).

#### **4.3.10. Glucose-6-Phosphate Dehydrogenase (G6PDH) Assay**

A modified version of the assay described by Deutsch (115) was followed. The enzyme reaction mixture containing 5.88  $\mu$ mol  $\beta$ -NADP, 88.5  $\mu$ mol  $\text{MgCl}_2$ , 53.7  $\mu$ mol glucose-6-phosphate, and 0.77 mmol maleimide was prepared. This mixture was used to obtain baseline (zero) of the spectrophotometer reading at 340 nm wavelength. To 1 ml of this mixture, 50  $\mu$ l of the sample was added. The rate of change in absorbance per minute was used to quantify the enzyme in the mixture using the extinction co-efficient of NADPH ( $6.22 \text{ mM}^{-1}\text{cm}^{-1}$ ).

#### **4.3.11. Guaiacol Peroxidase (GPX) Assay**

Modified version of assay developed by Laloue *et al.* (116) was used. Briefly, the enzyme reaction mixture contained 0.1 M potassium phosphate buffer (pH 6.8), 50 mM guaiacol solution and 0.2 mM hydrogen peroxide. To 1 mL of this reaction mixture, 50  $\mu$ L of enzyme extract was added. The absorbance was noted at zero time and then after 5 min. The rate of change in absorbance per minute was used to quantify the enzyme in the mixture using the extinction co-efficient of the oxidized product tetraguaiacol ( $26.6 \text{ mM}^{-1} \text{ cm}^{-1}$ ).

#### **4.3.12. Statistical Analysis**

Analysis at every time point was carried out in triplicates. Means, standard errors and standard deviations were calculated using Microsoft Excel 2003.

## **4.4. Results and Discussion**

### **4.4.1. Phenolic-Linked Antioxidant Activity and Visual Observations**


In all varieties, the phenolic content of peel extracts was much higher than pulp extracts in both water and 12% ethanol extraction (109). Similarly in most cases we see much higher free radical scavenging activity in peel extracts than pulp extracts was observed (Tab. 1). In general we see that Empire variety had consistently low phenolic levels during the storage period. Comparatively Macintosh and Cortland had higher phenolic content and Mutsu maintained consistent intermediate phenolic levels throughout the storage period. A clear link between phenolic content and antioxidant activity was observed. When phenolic content increased in antioxidant activity was also higher. For example Macintosh peel water extract at 2 months indicated a strong positive correlation between phenolic content and antioxidant activity (Fig. 8). This link is not surprising as phenolics are known to be free radical scavengers (49, 81). In addition the visual observations indicated that the varieties like Macintosh and Cortland which had high phenolic content also preserved better during the storage period (Fig. 9).

We also observed that Empire variety which had lower phenolic content deteriorated much faster than the other varieties. Empire variety began to deteriorate after 1 month of storage whereas other varieties remain firm until 2 months of storage after which they too began to deteriorate. After 3 months of storage, all apples had completely deteriorated. In many fruits, increase in hydroperoxides due to increasing ROS levels has been observed during senescence (75-77). Since the antioxidant activity was directly linked to the phenolic content it is logical to assume varieties with higher phenolic


content preserved better due to their improved free radical scavenging-linked antioxidant response systems.

**Table 1.** Phenolic content and antioxidant activity of apples over 3 months.

Total Phenolics Water Extraction	0 Month	1 Month	2 Month	3 Month
Cortland peel	689 µg/g	518 µg/g	677 µg/g	689 µg/g
Macintosh peel	305 µg/g	360 µg/g	1248 µg/g	1074 µg/g
Mutsu peel	299 µg/g	303 µg/g	477 µg/g	569 µg/g
Empire peel	285 µg/g	214 µg/g	447 µg/g	668 µg/g
Cortland pulp	49 µg/g	27 µg/g	44 µg/g	61 µg/g
Macintosh pulp	30 µg/g	37 µg/g	53 µg/g	39 µg/g
Mutsu pulp	49 µg/g	40 µg/g	12 µg/g	58 µg/g
Empire pulp	33 µg/g	22 µg/g	31 µg/g	38 µg/g
DPPH Inhibition% Water Extraction	0 Month	1 Month	2 Month	3 Month
Cortland peel	26%	6%	6%	12%
Macintosh peel	10%	10%	30%	3%
Mutsu peel	2%	1%	6%	6%
Empire peel	9%	5%	11%	17%
Cortland pulp	7%	5%	5%	6%
Macintosh pulp	7%	3%	5%	6%
Mutsu pulp	6%	6%	6%	5%
Empire pulp	7%	5%	6%	8%
Total Phenolics 12% Ethanol Extraction	0 Month	1 Month	2 Month	3 Month
Cortland peel	554 µg/g	235 µg/g	783 µg/g	739 µg/g
Macintosh peel	692 µg/g	301 µg/g	1039 µg/g	817 µg/g
Mutsu peel	588 µg/g	211 µg/g	622 µg/g	574 µg/g
Empire peel	466 µg/g	242 µg/g	525 µg/g	657 µg/g
Cortland pulp	123 µg/g	133 µg/g	98 µg/g	146 µg/g
Macintosh pulp	60 µg/g	103 µg/g	175 µg/g	134 µg/g
Mutsu pulp	55 µg/g	103 µg/g	131 µg/g	108 µg/g
Empire pulp	42 µg/g	66 µg/g	77 µg/g	74 µg/g
DPPH Inhibition% 12% Ethanol Extraction	0 Month	1 Month	2 Month	3 Month
Cortland peel	32%	23%	23%	3%
Macintosh peel	44%	45%	38%	4%
Mutsu peel	24%	25%	17%	20%
Empire peel	27%	10%	23%	19%
Cortland pulp	15%	12%	10%	10%
Macintosh pulp	11%	10%	10%	12%
Mutsu pulp	11%	12%	11%	11%
Empire pulp	8%	10%	9%	8%



**Figure 8.** Sample phenolic concentration ( $\mu\text{g/ml}$ ) and antioxidant activity. A) Peel water extract. B) Peel 12% ethanol extract.



**Figure 9.** Visual Observation of post-harvest changes in apples over 3 months. A) Macintosh B) Empire: 1) 0 Month 2-4) 3 Month.

#### **4.4.2. Superoxide Dismutase (SOD) Activity**

Previous studies in muskmelon and sunflower seeds indicated antioxidant enzyme response delayed senescence (86, 87). In Empire variety which had the worst post-harvest preservation of all varieties in this study deterioration began in the first month of storage itself where as all other varieties were well preserved until the second month of storage. Therefore it not surprising to observe that Empire variety had poor SOD activity compared to other varieties (Fig. 10). Over the storage period all varieties showed an increase in SOD activity after one month of storage followed by reduced activity. Empire variety, however, had a steady almost linear drop in SOD activity. As Phenolic levels have already been shown to boost antioxidant enzyme function in numerous plants (6, 82) the lower SOD activity in Empire variety could be attributed to its lower phenolic content. Conversely we also observe that apples with higher phenolic content like Macintosh had high SOD response. In general there is a positive correlation between phenolic content and SOD activity. However in Macintosh, a well preserved variety in later stages of storage (2-3 months) SOD activity was reduced, while phenolic content is still high. This indicates that combination of both phenolic metabolite and antioxidant enzyme response are essential for better post-harvest preservation.




Figure 10. Superoxide dismutase specific activity (U/mg protein).

#### **4.4.3. Proline Linked Pentose Phosphate Pathway (PLPPP) for Energy and Phenolic Synthesis.**

During the initial month of storage different stress factors seem to induce a rise in proline biosynthesis (Fig. 11). This stimulation likely maintains lower steady state G6PDH activity in Cortland and Macintosh (Fig. 14). This indicates no new glucose is utilized for anabolic pathways. The increase in proline synthesis also induces PDH activity (Fig. 12). The increase in PDH activity would catalyze more proline catabolism allowing proline to act as a substitute reductant for NADH for mitochondrial oxidative phosphorylation for energy needs. This would allow for synthesis of ATP with less oxidative stress in tissues. However after 1 month of storage proline synthesis and PDH activity start to decline (Fig. 11 and 12) and SDH activity rises in almost all varieties during this period (Fig. 13) indicating a switch to the TCA cycle for ATP synthesis. This switch to the TCA cycle would likely generate more ROS therefore more oxidative stress. This higher ROS potential combined with falling SOD activity in the later stages (Fig. 10) likely leads to the rapid deterioration of apples seen towards the end of our 3 month storage period. During the 3<sup>rd</sup> month of storage, very low or no enzyme activity in all varieties for all enzymes was observed (Fig. 11-15).


In Empire and Mutsu varieties however a dramatic drop in G6PDH activity in the first month of storage was observed indicating that the PPP is less active and glucose mobilization is drastically reduced for anabolic pathways (Fig. 14). A less active PPP would result in less phenolic synthesis and this can be confirmed by the low phenolic content in both Mutsu and Empire varieties when compared to Macintosh and Cortland varieties where more stable low steady state G6PDH activity was exhibited for a longer

part of the storage period. Therefore, even though Empire and Mutsu have successfully switched to proline catabolism for energy needs they do not maintain an active PPP for anabolic support. This inactive PPP results in lower levels of reductant and phenolics synthesis or remobilization in the system causing poor antioxidant response (free radical and enzyme-linked) leaving these varieties open to oxidative damage via ROS. In addition in Empire variety deterioration begins earlier possibly because of the poor SOD activity which showed a steady fall in activity unlike all other varieties (Fig. 10).


The GPX activity over the storage period showed a general downward trend in almost all cases except for Mutsu variety which showed an initial increase in the first month followed by a decline like all other varieties (Fig. 15). GPX polymerizes synthesizes phenolic phytochemicals to form lignins (116). Lignins function as structural support and are usually synthesized during growth periods. It is possible that here lignin synthesis via GPX is reduced so as to have more free phenolics for antioxidant functions in the tissue to respond to oxidative stress linked post-harvest deterioration.

From the above results phenolic metabolite and enzyme factors affect the post-harvest preservation of apples. However further study is needed to discern the exact interconnection between each of the factors. Insights gleaned can be used to create various treatments that would boost innate systems (proline-linked PPP driven phenolics synthesis) responsible for post-harvest preservation. This would also help elucidate markers such high phenolic content that would allow for selection and breeding of varieties that would preserve better. The indication that phenolic help prolong post-harvest storage is very promising, as many studies have already shown phenolics to have a vast array of health benefits. Phenolics have been shown to reduce the risk of many


chronic oxidative stress linked diseases like Type II diabetes, cardiovascular disease and cancer (7-11). This dual benefit of phenolics would allow not only for selection of better preserving varieties of apples but also varieties with health benefits as well.




**Figure 11.** Proline content of apples over 3 months ( $\mu\text{mol/mg protein}$ ).




**Figure 12.** Proline dehydrogenase specific activity in apples over 3 months. (U/mg protein).



**Figure 13.** Succinate dehydrogenase activity in apples over 3 months (nmol/mg protein).



**Figure 14.** Glucose-6-phosphate dehydrogenase specific activity in apples over 3 months (nmol/mg protein).



**Figure 15.** Guaiacol peroxidase specific activity in apples over 3 months (nmol/mg protein).

## CHAPTER 5

### HEALTH BENEFITS OF APPLE PHENOLICS FROM POST-HARVEST STAGES FOR POTENTIAL DIABETES MANAGEMENT USING *IN VITRO* MODELS

#### 5.1. Abstract

Increasing number of studies have shown that regular intake of fruits and vegetables have clear links to reduced risk of chronic diseases like diabetes and cardiovascular disease. The beneficial effects in many cases have been attributed to the phenolic and antioxidant content of the fruits and vegetables. Apples are a major source of fiber and contain good dietary phenolics with antioxidant function. Previous epidemiological studies have indicated that intake of apples reduces the risk of developing Type II diabetes. Our studies indicate that this reduced risk is potentially due to modulation of postprandial glucose increase by phenolics present in apples via inhibition of  $\alpha$ -glucosidase. Phenolic content was evaluated during 3 months of post-harvest storage of four varieties of apples and results indicated positive linkage to enhanced post-harvest preservation and  $\alpha$ -glucosidase inhibition. These *in vitro* results along with existing epidemiological studies provide strong biochemical rationale for further animal or human clinical studies.

Key words:

Apple post-harvest preservation, phenolic, antioxidant activity, Type II diabetes, anti-diabetic,  $\alpha$ -glucosidase.

## 5.2. Introduction

Growing evidence has suggested that the intake of fruits and vegetables is linked to reduced risk of chronic diseases like diabetes, cardiovascular disease and cancer (7-11). A recent study found apples to contain the second highest total phenolics content after cranberries, but the highest soluble phenolics (12). Apples also had the second highest free radical scavenging-linked antioxidant activity among the fruits investigated (12). Since the emerging oxidation-linked diseases can benefit from high intake of fruits and vegetables, high phenolic antioxidant-containing apples have the promise for enhancing human health through cellular protective functions. Apples are a popular fruit all over the world and have the potential to contribute as a significant source of phenolic antioxidants. In the United States, 22% of phenolic intake via fruit comes from apples (13). In Finland the main source of dietary phenolics are onions and apples and in the Netherlands tea, onions and apples are the biggest sources of phenolics (7, 9).

Abnormally high postprandial glucose levels or hyperglycemia has been linked to the onset of insulin independent diabetes mellitus (51). A good strategy for management of Type II diabetes is inhibition of enzymes that hydrolyze dietary polysaccharides in the gut which can significantly reduce the rise in blood sugar levels after a meal by reducing the absorption of monosaccharides by the enterocytes of the small intestine (52, 53). Enzymes that hydrolyze dietary polysaccharides and modulate gut absorption are pancreatic  $\alpha$ -amylase and  $\alpha$ -glucosidases (54, 55). However a common side effect of these enzyme inhibitory drugs like acarbose is the excessive inhibition of pancreatic  $\alpha$ -amylase which can result in abdominal distention, flatulence and diarrhea (56, 57). These side effects are caused by abnormal fermentation of unhydrolyzed polysaccharides by gut

bacteria (56, 58). Therefore in order to reduce these side effects but still manage hyperglycemia a high  $\alpha$ -glucosidase inhibition and low  $\alpha$ -amylase inhibition is beneficial.

Acute complications in patients suffering from Type II diabetes are generally hyperglycemia-induced metabolic problems and infection. Long term effects of hyperglycemia are microvascular complications like nephropathy, diabetic neuropathy, sexual dysfunction and retinopathy and the macrovascular complication of hypertension (59). Recent studies have shown that hyperglycemia triggers generation of free radicals in mesangial cells in the renal glomerulus, neuron cells in peripheral nerves and capillary endothelial cells in the retina (60). The generation of free radicals in all these cells types causes oxidative stress which can be the cause of microvascular complications generally linked with hyperglycemia (60). Most cells are capable of reducing glucose transport inside the cell in hyperglycemic conditions so as to maintain a constant internal glucose level, however the cells usually damaged by hyperglycemia were found to be inefficient in keeping their internal glucose levels constant (61, 62). Therefore it is not only important to control postprandial hyperglycemia but also keep in check any cellular redox imbalances to prevent diabetic complications. Phenolic phytochemicals have been found to have potential in the management of oxidative stress linked chronic diseases and therefore have potential for managing complications of diabetes (5, 20, 46-48, 118).

A growing number of studies from our laboratory have found many common foods to have the correct profile of  $\alpha$ -glucosidase inhibition and low  $\alpha$ -amylase inhibition and free radical scavenging-linked antioxidant activity (15-17). This offers the potential for good postprandial blood glucose management without the common side-effects

associated with high  $\alpha$ -amylase inhibition. In addition these same foods have free radical scavenging-linked antioxidant activity which can help maintain the redox balance in susceptible cells.

An epidemiological study by Knekt *et al* in 2001 (8) of more than 10,000 Finnish men and women showed intake of foods containing flavanoids was associated with lower risk of Type II diabetes. The study indicated the lower risk was associated with high intake of quercetin. The same study also states the strongest association between high flavonoid intake with lower risk for Type II diabetes was seen when the source of flavonoids was from apples and berries (8). Therefore a diet containing apples has the potential to reduce risk of Type II diabetes. Since apple is a common fruit with no known side effects, any  $\alpha$ -glucosidase inhibiting effects, if found, are promising for Type II diabetes management.

Based on the above background this study explores the potential phenolic-linked anti-diabetic benefits of apples. Specifically phenolic-linked changes were investigated during post-harvest storage of four varieties of apple over three months (119). The objective of this study was to determine how the changes in phenolic content over the storage period which has relevance to post-harvest preservation (119) also has relevance for enhanced anti-diabetes-linked health benefits that could be attributed to the phenolic content. The health relevant parameters investigated were free radical-linked antioxidant activity and *in vitro* inhibition of enzymes  $\alpha$ -glucosidase and  $\alpha$ -amylase that are relevant to glycemic index control associated with Type II diabetes management.

### **5.3. Materials and Methods**

The freshly picked apples were purchased from a local orchard (Clarkdale Orchards Deerfield, MA, US). Four of the popular varieties of apples were selected to be tested; Cortland, Macintosh, Empire and Mutsu. These were stored in a cold room (5°C) for 3 months.

#### **5.3.1. Phenolic Extraction**

Soluble phenolics were extracted from the peel and pulp of each variety separately. Phenolics were extracted using distilled water and 12% ethanol. The water extractions were done with 20 g of peel in 50 ml of water and 100 g of pulp in 50 ml of water. The ethanol extractions were done with 5 g of peel in 15 ml of 12% ethanol and 10 g of pulp in 20ml of 12% ethanol. Apples were first peeled then cut and weighed. Peel and pulp was then mixed with either distilled water or 12% ethanol. This was then homogenized for 2 min using a blender. Resulting mix was collected and centrifuged for 5 min. Supernatant was collected and stored at -20°C. This was done every month until the 3<sup>rd</sup> month of storage.

#### **5.3.2. Total Phenolics Assay**

The total phenolic content was determined by an assay modified from Shetty *et al.* (109). Briefly, one milliliter of extract was transferred into a test tube and mixed with 1 ml of 95% ethanol and 5 ml of distilled water. To each sample 0.5 ml of 50% (v/v) Folin-Ciocalteu reagent was added and mixed. After 5 min, 1 ml of 5% Na<sub>2</sub>CO<sub>3</sub> was

added to the reaction mixture and allowed to stand for 60 min. The absorbance was read at 725 nm. The absorbance values were converted to total phenolics and were expressed in micrograms equivalents of gallic acid per grams fresh weight (FW) of the sample. Standard curves were established using various concentrations of gallic acid in 95% ethanol.

### **5.3.3. Antioxidant Activity by 1, 1-Diphenyl-2-Picrylhydrazyl Radical (DPPH) Inhibition Assay**

To 3 ml of 60  $\mu$ M DPPH in ethanol, 250  $\mu$ l of each extract was added, the decrease in absorbance was monitored at 517 nm until a constant reading was obtained. The readings were compared with the controls, which contained 250  $\mu$ l of 95% ethanol instead of the extract. The % inhibition was calculated by:

$$\% \text{ inhibition} = \left( \left[ \frac{A_{517}^{Control} - A_{517}^{Extract}}{A_{517}^{Control}} \right] \right) \times 100$$

### **5.3.4. $\alpha$ -Glucosidase Inhibition Assay**

$\alpha$ -Glucosidase (EC 3.2.1.20) was purchased from Sigma Chemical Co. A volume of 50  $\mu$ l of sample solution and 100  $\mu$ l of 0.1 M phosphate buffer (pH 6.9) containing  $\alpha$ -glucosidase solution (1.0 U/ml) were incubated in 96 well plates at 25 $^{\circ}$ C for 10 min. After pre-incubation, 50  $\mu$ l of 5 mM p-nitrophenyl- $\alpha$ -D-glucopyranoside solution in 0.1 M phosphate buffer (pH 6.9) was added to each well at timed intervals. The reaction mixtures were incubated at 25 $^{\circ}$ C for 5 min. Before and after incubation, absorbance

readings were recorded at 405 nm by micro-array reader (Thermomax, Molecular device Co., Sunnyvale, CA) and compared to a control which had 50  $\mu$ l of buffer solution in place of the extract. The  $\alpha$ -glucosidase inhibitory activity was expressed as inhibition % and was calculated as follows:

$$\% \text{ inhibition} = \left( \left[ \frac{\Delta A_{405}^{Control} - \Delta A_{405}^{Extract}}{[\Delta A_{405}^{Control}]} \right] \right) \times 100$$

### 5.3.5. $\alpha$ -Amylase inhibition Assay

Porcine pancreatic  $\alpha$ -amylase (EC 3.2.1.1) was purchased from Sigma Chemical Co. A total of 500  $\mu$ l of extract and 500  $\mu$ l of 0.02 M sodium phosphate buffer (pH 6.9 with 0.006 M sodium chloride) containing  $\alpha$ -amylase solution (0.5 mg/ml) were incubated at 25 $^{\circ}$ C for 10 min. After pre-incubation, 500  $\mu$ l of a 1% starch solution in 0.02 M sodium phosphate buffer (pH 6.9 with 0.006 M sodium chloride) was added to each tube at timed intervals. The reaction mixtures were then incubated at 25 $^{\circ}$ C for 10 min. The reaction was stopped with 1.0 ml of dinitrosalicylic acid color reagent. The test tubes were then incubated in a boiling water bath for 5 min, cooled to room temperature. The reaction mixture was then diluted after adding 10 ml distilled water and absorbance was measured at 540 nm.

$$\% \text{ inhibition} = \left( \left[ \frac{A_{540}^{Control} - A_{540}^{Extract}}{[A_{540}^{Control}]} \right] \right) \times 100$$

### **5.3.6. Statistical Analysis**

Analysis at every time point was carried out in triplicates. Means, standard errors and standard deviations were calculated using Microsoft Excel 2003.

## **5.4. Results and Discussion**

### **5.4.1 Visual Observations**

On visual examination of the post-harvest stored apples were observed for visual changes over a 3 month period (Fig. 16). Empire variety began to spoil much earlier than the other varieties. After the first month itself all the Empire apples had started to become soft and completely deteriorate. All other apple varieties investigated remained firm until two months, after which by the third month they began to deteriorate as well.

### **5.4.2. Phenolic Content of Apples**

In all cases the phenolics content of peel extracts was much higher than that of pulp extracts (Fig. 17, 18) (119). Further 12% ethanol extracts had higher phenolic content than water extracts across all varieties and tissue types except in Macintosh in later stages of storage.

In peel extracts Macintosh had the highest water extractable phenolics ( $>1000 \mu\text{g/g}$  Fresh weight, FW) in the later part of the storage period (2<sup>nd</sup> and 3<sup>rd</sup> month). Cortland had intermediate content over the entire storage period ( $>600 \mu\text{g/g}$  FW). Empire and Mutsu in general had low phenolic content ( $400\text{-}600 \mu\text{g/g}$  FW) in the early stages (0-2<sup>nd</sup> month) with an increase towards intermediate content ( $>600 \mu\text{g/g}$  FW) in the last month of storage. In the 12% ethanol extracts (peel), Macintosh in general had higher ( $>800 \mu\text{g/g}$

FW) of phenolic content. Cortland was again intermediate ( $>600 \mu\text{g/g FW}$ ) and Empire and Mutsu on the lower side ( $<600 \mu\text{g/g FW}$ ).

In the pulp extracts the water extractable phenolics were generally less than  $50\mu\text{g/g FW}$  across all varieties, except for Macintosh Cortland and Mutsu in the later stages of storage. In the 12% extracts (pulp) Macintosh, Mutsu and Cortland in general had over  $100 \mu\text{g/g FW}$  in later stages, where as Empire was consistently lower at less than  $100 \mu\text{g/g FW}$  through out the storage period.

### **5.4.3. Antioxidant Activity**

The peel samples generally had higher free radical scavenging-linked antioxidant activity, except for Mutsu in the water extract (Fig. 19, 20). The peel in 12% ethanol extracts of Macintosh was superior with initial high inhibition of 45% that gradually reduced after the first month with a low final inhibition ( $<5\%$ ). Other varieties had an intermediate inhibition (25% to 30%) that was maintained through most of the storage period with the exception of empire which had a drop to 10% inhibition in the first month but regained close to original inhibition after and Cortland which had a drop in inhibition% on the third month to less than 5%.

In water extracts of peel, Macintosh had an initial low inhibition (10%) that increased to 30% by the second month of storage before dropping down to less than 5%. Cortland had a high initial inhibition ( $\sim 25\%$ ) which dropped substantially in the first month of storage itself (5% to 10%), however inhibition increased over the rest of the storage period but never to initial levels. Empire and Mutsu both had low initial inhibition levels ( $<10\%$ ) which dropped slightly in the first month but then began to rise

for the rest of the storage period, Empire had a intermediate final inhibition (15% to 20%) but Mutsu still had a low inhibition level of 5%.

The pulp extracts, both water and 12% ethanol, generally had low DPPH inhibition that we was maintained steadily over the storage period.

#### **5.4.4. $\alpha$ -Glucosidase and $\alpha$ -Amylase Inhibition**

Peel extracts in general had higher  $\alpha$ -glucosidase inhibitory activity than pulp (21, 22). In the 12% ethanol extracts of peel, Macintosh had the highest  $\alpha$ -glucosidase inhibition (75%-90%) over the three month storage period. Cortland was the next highest (60%-80%) over storage time. Empire and Mutsu had an intermediate initial inhibition of around 60% followed by lower inhibition for rest of the storage period (<50%). The water extracts of peel generally had higher inhibition in the later stages of storage with Macintosh having the highest inhibition (>90%) at two months storage.

The pulp extracts in 12% ethanol had consistent low levels of inhibition (20% - 40%) over the storage period. The pulp water extracts had intermediate levels of inhibition (40% - 60%) but this was maintained over the storage period.

In all samples no detectable  $\alpha$ -amylase inhibition was observed.

The phenolic content especially in late storage was highest in Macintosh and this was reflected in better post-harvest storage of this variety based on visual observations (119). Empire which had the lowest phenolic content deteriorated most rapidly with Cortland and Mutsu being intermediate.


It is significant that sample phenolic concentration ( $\mu\text{g/ml}$ ) was proportional to the  $\alpha$ -glucosidase inhibition trend this indicates a clear link between the two (Pearson Correlation Coefficient: 0.875, Fig. 23-25). In case where a dramatic rise or fall in phenolic concentration was observed a similar rise or fall the  $\alpha$ -glucosidase inhibition followed. This trend is most evident in Macintosh where the phenolic content and  $\alpha$ -glucosidase inhibition of the peel extracts show the most promising health relevant potential. Further evidence of this trend is the fact that, like phenolic content, inhibitory activity in the peel samples is much higher than that of pulp samples across all varieties. A similar relationship is seen between the antioxidant activity and phenolic content.

In this study Macintosh and Cortland appear to be superior varieties as they can preserve better which is linked to their phenolic content which in turn is closely linked to  $\alpha$ -glucosidase inhibition. This means well stored apples with high phenolic content have potential for some glycemic index modulation while contributing antioxidants that can positively influence tissues susceptible to glucose-linked oxidative stress. The lack of  $\alpha$ -amylase inhibition is also promising because this prevents common side effects that are associated with carbohydrates hydrolyzing enzyme inhibitors but still allows for glycemic modulation by allowing control over polysaccharide breakdown to monosaccharides.



Further advantages of this study is the indication towards mechanism of action where dietary phenolics with  $\alpha$ -glucosidase inhibitory activity can counter negative effects of glucose-induced oxidative dysfunction a microvascular complication of Type II diabetes. This occurs via direct glycemic modulation by phenolics via inhibition of starch hydrolyzing enzymes and antioxidant activity which can help prevent oxidative damage in susceptible tissues. In this study these activities,  $\alpha$ -glucosidase inhibition and

antioxidant activity; have shown strong ties to the phenolic content in the apples. Though these *in vitro* studies may not indicate clinical benefits but they provide strong biochemical rationale for animal and human clinical studies. These studies along with already existing epidemiological studies provide the biochemical basis for further clinical investigation.



The other interesting observation emerging out of this study is the correlation between high phenolic content apple varieties to extended post-harvest preservation. This also translates into health beneficial consequences in the context of higher antioxidant activity and Type II diabetes linked  $\alpha$ -glucosidase inhibition. Therefore strategies to understand phenolic-linked post-harvest preservation and natural treatments to extend this preservation in selected varieties, such as Macintosh and Cortland in this study, can be basis for food ingredient design for health benefits. These strategies can then be extended to prolong post-harvest preservation and enhance phenolic linked health benefits of a wide variety of fruits and vegetables.



**Figure 16.** Visual observation of post-harvest changes in apples over 3 months. A) Macintosh B) Empire: 1) 0 Month 2-4) 3 Month.




**Figure 17.** Total phenolic content of apples water extraction ( $\mu\text{g/g}$  FW).  
 A) Peel samples. B) Pulp samples.




**Figure 18.** Total phenolic content of apples 12% ethanol extraction ( $\mu\text{g/g}$  FW).  
 A) Peel samples. B) Pulp samples.

A)




B)




**Figure 19.** DPPH inhibition. Apple water extracts. A) Peel samples. B) Pulp samples.

A)




B)




**Figure 20.** DPPH inhibition. Apple 12% ethanol extracts.  
A) Peel samples. B) Pulp samples.

A)




B)




**Figure 21.**  $\alpha$ -glucosidase inhibition. Apple water extracts.  
A) Peel samples. B) Pulp samples.

A)




B)




**Figure 22.**  $\alpha$ -glucosidase inhibition. Apple 12% ethanol extracts.


A) Peel samples. B) Pulp samples.



**Figure 23.** Sample phenolic concentration ( $\mu\text{g/ml}$ ) and  $\alpha$ -glucosidase inhibition. Apple peel water extracts.



**Figure 24.** Sample phenolic concentration ( $\mu\text{g/ml}$ ) and  $\alpha$ -glucosidase inhibition. Apple peel 12% ethanol extracts.



**Figure 25.** Correlation between sample phenolic concentration (µg/ml) and α-glucosidase inhibition.

## BIBLIOGRAPHY

1. Mann J. *Secondary Metabolism*. Oxford Chemistry Series. Oxford: Clarendon Press; 1978.
2. Bravo L. Phenolic phytochemicals: chemistry, dietary sources, metabolism, and nutritional significance. *Nutr. Rev.* 1998; 56: 317–333.
3. Crozier A, Burns J, Aziz AA, Stewart AJ, Rabiasz HS, Jenkins GI, Edwards CA, Lean M. Antioxidant flavanols from fruits, vegetables and beverages: measurements and bioavailability. *Biol. Res.* 2000; 33: 79–88.
4. Briskin DP. Medicinal plants and phytomedicines. Linking plant biochemistry and physiology to human health. *Plant Physiol.* 2000; 124: 507-514
5. Shetty K. Biotechnology to harness the benefits of dietary phenolics; Focus on *Lamiaceae*. *Asia Pacif. J. Clini. Nutri.* 1997; 6: 162-171.
6. Vattem DA, Randhir R, Shetty K. Cranberry phenolics-mediated elicitation of antioxidant enzyme response in fava bean (*vicia faba*) sprouts. *J. of Food. Biochem.* 2005; 29: 41-70.
7. Knekt P, Jarvinen R, Seppanen R, et al. Dietary flavonoids and the risk of lung cancer and other malignant neoplasms. *Am. J. Epidemiol.* 1997; 146: 223-230.
8. Knekt P, Kumpulainen J, Jarvinen R, et al. Flavonoid intake and risk of chronic diseases. *Am. J. Clin. Nutr.* 2002; 76: 560-568.
9. Hertog M, Feskens E. Dietary antioxidant flavonoids and risk of coronary heart disease. *Lancet.* 1993; 342: 1007-1011.
10. Hertog M, Kromhout D, Aravanis C, et al. Flavonoid intake and long-term risk of coronary heart disease and cancer in the seven counties study. *Arch. Intern. Med.* 1995; 155: 381-386.
11. Boyer J, Liu RH. Apple phytochemicals and their health benefits. *Nutrition Journal.* 2004; 3: 5-9.
12. Sun J, Chu Y, Wu X, Liu RH. Antioxidant and antiproliferative Activities of Common Fruits. *J. Agric Food Chem.* 2002; 50: 7449-7454.
13. Vinson JA, Su X, Zubik L, Bose P. Phenol antioxidant quantity and quality in foods: fruits. *J. Agric Food Chem.* 2001; 49: 5315-5321.

14. World Health Organization Website. Diabetes Factsheet. Factsheet No312. 2006. Available at: <http://www.who.int/mediacentre/factsheets/fs312/en/>. Accessed May 5 2007.
15. Kwon Y-I, Vatter DA, Shetty K. Evaluation of clonal herbs of *Lamiaceae* species for management of diabetes and hypertension. *Asia Pac. J. Clin. Nutr.* 2006; 15: 107-118.
16. Kwon Y-I, Apostolidis E, Shetty K. Evaluation of pepper (*Capsicum annuum*) for management of diabetes and hypertension. *J. Food Biochemistry*. 2006; In press.
17. McCue P, Kwon Y-I, Shetty K.. Anti-amylase, anti-glucosidase and anti-angiotension I converting enzyme potential of select foods. *J. Food Biochem.* 2005; 29: 278-294.
18. Strack D. Phenolic metabolism. In: Deyand PM, Harborne JB, eds. *Plant Biochemistry*. San Diego, CA: Academic Press; 1997: 387–416.
19. Chugh LK, Sawhney SK. Effects of cadmium on activities of some enzymes of glycolysis and pentose phosphate pathway in pea. *Biol.Plantarum*. 1999; 42: 401-407.
20. Shetty K, Wahlqvist ML. A Model for the role of the proline-linked pentose-phosphate pathway in phenolic phytochemical biosynthesis and mechanism of action for human health and environmental applications. *Asia Pacific J. Clin. Nutr.* 2004; 13: 1-24.
21. Shetty P, Aatallah MT, Shetty K. Stimulation of total phenolics, L-DOPA and antioxidant activity through proline-linked pentose phosphate pathway in response to proline and its analogue in germinating fava beans (*Vicia faba*). *Process Biochem.* 2003; 38: 1707-17.
22. Dixon R, Harrison M, Lamb C. Early events in the activation of plant defence responses. *Annl. Rev. of Phytopath.* 1994; 32: 479-501
23. Dixon R, Paiva N. Stress-induced phenylpropanoid metabolism. *Plant Cell.* 1995; 7: 1085-1097.
24. Brooker F, Miller J. Phenylpropanoid metabolism and phenolic composition of soybean [*Glycine max* (L) Merr.] leaves following exposure to ozone. *J. Exp. Bot.* 1998; 49: 1191-1202.
25. Zimmerman YZ, Cahill PR. Heat shock and thermotolerance in plant and animal embryogenesis. *New Biology* 1991; 3: 641-650.

26. Rhodes JM, Wooldorton LSC. In: Kahl G, ed. The biosynthesis of phenolic compounds in wounded plant storage tissues. Biochemistry of wounded plant tissues. Berlin: W de Gruyter; 1978: 286.
27. Hahlbrock, Scheel D. Physiology and molecular biology of phenylpropanoid metabolism. *Plant Mol. Biol.* 1989; 40: 347-369.
28. Christie PJ, Alefenito MR, Walbot V. Impact of low temperature stress on general phenylpropanoid and anthocyanins pathways: Enhancement of transcript abundance and anthocyanin pigmentation in maize seedlings. *Planta.* 1994; 194: 541-549.
29. Beggs CJ, Kuhn K, Bocker R, Wellmann E. Phytochrome induced flavanoid biosynthesis in mustard (*Sinapsis alba* L) cotyledons: Enzymatic control and differential regulation of anthocyanin and quercetin formation. *Planta.* 1987; 172: 121-126.
30. Lois R, Buchanan BB. Severe sensitivity to ultraviolet light radiation in *Arabidopsis* mutant deficient in flavanoid accumulation. Mechanisms of UV-resistance in *Arabidopsis*. *Planta.* 1994; 194: 504-509.
31. Rice-Evans CA, Miller NJ, Bolwell PG, Bramley PM, Pridham JB. The relative antioxidant activities of plant-derived polyphenolic flavonoids. *Free Rad Res.* 1995; 22: 375-383.
32. Hertog MGL, Hollman PCH, Katan MB. Content of potentially anticarcinogenic flavonoids of 28 vegetables and 9 fruits commonly consumed in the Netherlands. *J Agric Food Chem.* 1992; 40: 2379-2383.
33. Jorgensen LV, Madsen HL, Thomsen MK, Dragsted LO, Skibsted LH. Regulation of phenolic antioxidants from phenoxyl radicals: An ESR and electrochemical study of antioxidant hierarchy. *Free Radical Res.* 1999; 30: 207-220.
34. Foti M, Piattelli M, Amico V, Ruberto G. Antioxidant activity of phenolic meroditerpenoids from marine algae. *J Photochem Photobiol.* 1994; 26: 159-164.
35. Lee K, Kim Y, Kim D, Lee H, Lee C. Major phenolics in apple and their contribution to total antioxidant capacity. *J. Agric. Food Chem.* 2003; 51: 6516-6520.
36. Huang MT, Lysz T, Ferraro T, Conney AH. Inhibitory effects of curcumin on tumor promotion and arachidonic acid metabolism in mouse epidermis. In: Wattenberg L, Lipkin M, Boone CW, Kelloff GJ, eds. *Cancer chemoprevention.* Boca Raton, FL: CRC Press; 1992: 375-391.

37. Masuda T, Jitoe A. Antioxidative and anti-inflammatory compounds from tropical gingers: Isolation, structure determination, and activities of cassumunins A, B and C, new complex curcuminoids from *Zingiber cassumunar*. *J Agric Food Chem* 1994; 42: 1850-1856.
38. Osawa T, Sugiyama Y, Inayoshi M, Kawakishi S. Antioxidant activity of tetrahydrocurcuminoids. *Biosci Biotechnol Biochem* 1995; 59:1609-1612.
39. Lim GP, Chu T, Yang F, Beech W, Frauschy SA, Cole GM. The curry spice curcumin reduces oxidative damage and amyloid pathology in an Alzheimer transgenic mouse. *J Neuroscience* 2001; 21: 8370-8377.
40. Jitoe A, Masuda T, Tengah GP, Suprpta DN, Gara IW, Nakatani N. Antioxidant activity of tropical ginger extracts and analysis of the contained curcuminoids. *J Agric Food Chem* 1992; 40: 1337-1340.
41. Peake PW, Pussel BA, Martyn P, Timmermans V, Charlesworth JA. The inhibitory effect of rosmarinic acid on complement involves the C5 convertase. *Int J Immunopharmac*. 1991; 13: 853-857.
42. Winterhoff H, Gumbinger HG, Sourgens, H. On the antiogonadotropic activity of *Lithospermum* and *Lycopus* species and some of their phenolic constituents. *Planta Medica* 1988; 54: 101-106.
43. Howell AB, Vorsa N, Der Marderosian A, Foo LY. Inhibition of adherence of P-fimbriated *Escherichia coli* to uroepithelial-cell surfaces by proantho-cyanidin extracts from cranberries. *N Engl J Med*. 1998; 339: 1085-1086.
44. Howell AB, Foxman B. Cranberry juice and adhesion of antibiotic-resistant uropathogens. *JAMA* 2002; 287:3082- 3083.
45. Himejima M, Kubo I. Fungicidal activity of polygodial in combination with anethol and indole against *Candida albicans*. *J Agric Food Chem*. 1993; 41: 1776-1779.
46. Shetty K and Labbe RL. Food-borne pathogens, health and role dietary phytochemicals. *Asia Pac. J. Clin. Nutr.* 1998; 7: 270-276.
47. Shetty K. Phytochemicals: Biotechnology of phenolic phytochemicals for food preservatives and functional food applications. In: Francis FJ, eds. *Wiley Encyclopedia of Food Science and Technology*, 2<sup>nd</sup> Edition. New York: Wiley Publishers; 1999: 1901-1909.
48. Shetty K. Biosynthesis of rosmarinic acid and applications in medicine. *J Herbs Species and medicinal Plants*. 2001; 8: 161-181.

49. Kang NM, Lee KW, Lee SJ, Lee CY, Lee HJ. Effects of phenolics in Empire apples on hydrogen peroxide-induced inhibition of gap junctional intercellular communication. *BioFactors*. 2004; 21: 361-365.
50. Block G, Patterson B. and Subar A. Fruit, vegetables, and cancer prevention: a review of the epidemiological evidence. *Nutr. Cancer*. 1992; 18: 1-29.
51. Fonseca V. Clinical significance of targeting postprandial and fasting hyperglycemia in managing type 2 diabetes mellitus. *Curr. Med. Res. Opin*. 2003; 19: 635-641.
52. Puls W, Keup U, Krause H, Thomas PG, Hoffmeister F. Glucosidase inhibition: a new approach to the treatment of diabetes, obesity and hyperlipoproteinemia. *Naturwissen Schafte*. 1977; 64: 36- 537.
53. Ratner RE. Controlling postprandial hyperglycemia. *Am. J. Cardiol*. 2001; 88: 26-31.
54. Bischoff H. Pharmacology of glucosidase inhibitor. *Eur. J. Clin. Invest*. 1994; 24(3): 3-10.
55. Harris M.I. and Zimmer P. Classification of diabetes mellitus and other categories of glucose intolerance. In: Alberti et al eds. *International textbook of diabetes mellitus*. London: John Wiley; 1992: 3-18.
56. Puls W, Keup U. Metabolic studies with an amylase. In: Howard A, ed. *Recent advances in obesity research*. Vol 1. London: Newman Publisher; 1975: 391
57. Bischoff H, Puls W, Krause HP, Schutt H, Thomas G. Pharmacological properties of the novel glucosidase inhibitors BAY m 1099 (miglitol) and BAY o 1248. *Diabetes Res. Clin. Pract*. 1985; 1: 53-62.
58. Horii S, Fukasse K., Matsuo T, Kameda K., Asano N, Masui Y. Synthesis and  $\alpha$ -D-glucosidase inhibitory activity of N-substituted valiolamine derivatives as potent oral antidiabetic agents. *J. Med. Chem*. 1987; 29: 1038-1046.
59. Nishikawa T, Edelstein D, Du XL, Yamagishi S, et al. Normalizing mitochondrial superoxide production blocks three pathways of hyperglycemic damage. *Nature*. 2000; 404: 787–790.
60. Brownlee M. The pathobiology of diabetic complications. *Diabetes*. 2005; 54: 1615-1625.
61. Helig CW, Concepcion LA, Riser BL, Freytag SO, Zhu M, Cortes P. Overexpression of glucose transporters in rat mesangial cells cultured in a normal glucose milieu mimics the diabetic phenotype. *J Clin Invest*. 1995; 96:1802–1814.

62. Kaiser N, Sasson S, Feener, EP, et al. Differential regulation of glucose transport and transporters by glucose in vascular endothelial and smooth muscle cells. *Diabetes*. 1993; 42: 80–89.
63. Costantino L, Rastelli G, Gamberini MC, et al. 1-Benzopyran-4-one antioxidants as aldose reductase inhibitors. *J Med Chem*. 1999; 42: 1881-1893.
64. Bolwell GP, Wojtaszek P. Mechanisms for the generation of reactive oxygen species in plant defense: a broad perspective. *Physiol. Mol. Plant Pathol*. 1997; 51: 347-366.
65. Karpinski S, Reynolds H, Karpinska B, Wingsle G, Creissen G, Mullineaux P. Systemic signaling and acclimation in response to excess excitation energy in *Arabidopsis*. *Nature*. 1999; 284: 654-657.
66. Quartacci MF, Cosi E, Navari-Izzo F. Lipids and NADH dependent superoxide production in plasma membrane vesicles from roots of wheat germ under copper deficiency or excess. *J Exp Bot*. 2001; 52: 77-84.
67. Del Rio LA, Corpas FJ, Sandalio LM, Palma JM, Gomez M, Barroso JB. Reactive oxygen species, antioxidant systems and nitric oxide in peroxisomes. *J. Exp. Bot*. 2002; 53: 1255-1272.
68. Dat J, Vandenabeele S, Vranova E, Van Montagu M, Inze, D, Van Breusegem F. Dual action of the active oxygen species during plant stress responses. *Cell. Mol. Life Sci*. 2000; 57: 779- 795.
69. Hernandez JA, Ferrer MA, Jimenez A, Ros-barcelo A, Sevilla F. Antioxidant systems and O<sub>2</sub>-/H<sub>2</sub>O<sub>2</sub> production in the apoplast of *Pisum sativum* L. leaves: its relation with NaCl induced necrotic lesions in minor veins. *Plant Physiol*. 2001; 127: 817-831.
70. Jimenez A, Hernandez JA, Pastori G, Del Rio LA, Sevilla F. Role for the ascorbate-glutathione cycle of mitochondria and peroxisomes in the senescence of pea leaves. *Plant Physiol*. 1998; 118: 1327-1335.
71. Doillard MJ, Paulin A, Massot JC. Free radical production, catalase, and superoxide dismutase activities and membrane integrity during senescence of petal of cut carnations (*Dianthus caryophyllus*). *Physiologia Plantarum*. 1987; 71: 197-202.
72. Bartoli CG, Simontachhi M, Montaldi E, Puntarulo S. Oxidative stress, antioxidant capacity and ethylene production during ageing of cut carnation (*Dianthus caryophyllus*) petals. *J.Exp. Bot*. 1996; 297: 595-601.

73. Philosoph-Hadas S, Meir S, Akiri B, Kanner K. Oxidative defense systems in leaves of three edible herb species in relation to their senescence rate. *J of Agric and Food Chem.* 1994; 42: 2376-81.
74. Thompson JE, Brown JH, Paliyath G, Todd JF, Yao K. Membrane phospholipid catabolism primes the production of activated oxygen in senescing tissue. In: Pell E, Steffen K, eds. *Active oxygen/oxidative stress and plant metabolism.* Rockville, MD: ASPP; 1991: 57-66.
75. Frenkel C. Role of hydroperoxides in the onset of senescence processes in plant tissues. In: Hultin HO, Milner M, Eds; *Postharvest Biology and Biotechnology.* Westport, Spychalla : Food and Nutrition Press; 1978: 433-448.
76. Thompson JE, Ledge RL, Barber RF. The role of free radicals in senescence and wounding. *New Phytol.* 1987; 105: 317-344.
77. Rogiers SY, Kumar MGN, Knowles NR. Maturation and ripening of fruit of *Amelanchier alnifolia* Nutt. Are accompanied by increasing oxidative stress. *Ann. Bot.* 1998; 81: 203-211.
78. Hodges DM, Andrews CJ, Johnson DA, Hamilton RI. Antioxidant compound responses to chilling stress in differentially sensitive inbred maize lines. *Phytophologia plantarum.* 1996; 98: 685-92.
79. Foyer CH, Descourvieres P, Kunert KJ. Protection against oxygen radicals: an important defence mechanism studied in transgenic plants. *Plant Cell Environ* 1994; 17: 507-523.
80. Hodges DM, Forney CF. The effects of ethylene, depressed oxygen and elevated carbon dioxide on antioxidant profiles of senescing spinach leaves. *J Exp Bot.* 2000 ; 51 : 645-655.
81. Zhang J, Stanley RA, Adaim A, Melton LD, Skinner MA. 2006. Free radical scavenging and cytoprotective activities of phenolic antioxidants. *Mol. Nutr. Food Res.* 50, 996-1005.
82. Duval B, Shetty K. The stimulation of phenolics and antioxidant activity in pea (*pisum sativum*) elicited by genetically transformed anise root extract. *J. Food Biochem.* 2000; 25: 361-77.
83. Jimenez A, Creissen G, Kular B, et al. Changes in oxidative processes and components of the antioxidant system tomato fruit ripening. *Planta.* 2002; 214: 751-758.

84. Jimenez A, Romojaro F, Gomez JM, Llanos MR, Sevilla F. Antioxidant Systems and Their Relationship with the Response of Pepper Fruits to Storage at 20 °C. *J. Agric. Food Chem.* 2003; 51: 6293-6299.
85. Pastori GM, Del-Rio LA. Natural senescence of pea leaves: and activated oxygen-mediated function for peroxisomes. *Plant Physiol.* 1997; 113: 411-418.
86. Lacan D, Baccou J-C. High levels of antioxidant enzymes correlate with delayed senescence in non-netted muskmelon fruits. *Planta.* 1998; 204: 377-82.
87. Bailly C, Benamer A, Corbineau F, Come D. Changes in the malondialdehyde content and in superoxide dismutase, catalase and glutathione reductase activities in sunflower seeds as related to deterioration during accelerated aging. *Physiologia Plantarum.* 1996; 97: 104- 10.
88. Meir S, Kanner K, Akiri B, Philosoph-Hadas S. Determination and involvement of aqueous reducing compounds in oxidative defense systems of various senescing leaves. *J. of Agric. and Food Chem.* 1995; 43: 1813-19.
89. Stewart CR, Larher F. Accumulation of amino acids and related compounds in relation to environmental stress. In: Mifflin BJ, ed. *The Biochemistry of Plants.* Vol 5. New York: Academic Press; 1980: 609-635.
90. Thompson JF. Arginine synthesis, proline synthesis, and related process. In: Mifflin BJ, ed. *The Biochemistry of Plants.* Vol 5. New York: Academic Press; 1980: 375-403.
91. Rhodes D. Metabolic Responses to stress. In: Davies DD, ed. *Biochemistry of Plants.* Vol 12. New York: Academic Press; 1987: 201-241.
92. Taylor CB. Proline and water deficit: ups and downs. *The Plant Cell* 1996; 8:1221-1224.
93. Hare PD, Cress WA. Metabolic implications of stress induced proline accumulations in plants. *Plant Growth Regulation* 1997; 21: 79-102.
94. Rhodes D, Handa S, Bressan RA. Metabolic changes associated with adaptation of plant cells to water stress. *Plant Physiol* 1986; 82: 890-902.
95. Sudhakar C, Reddy PS, Veeranjanyulu K. Effects of salt stress on enzymes of proline synthesis and oxidation in green gram (*Phaseolus aureus* Roxb.) seedlings. *J Plant Physiol* 1993; 141: 621-623.
96. LaRosa PC, Rhodes D, Rhodes JC, Bressan RA, Csonka LN. Elevated accumulation of proline in NaCl-adapted tobacco cells is not due to altered  $\alpha$ -pyrroline-5-carboxylate reductase. *Plant Physiol* 1991; 96: 245-250.

97. Reddy PS, Veeranjanyulu K. Proline metabolism in senescing leaves of horsegram (*Macrotyloma uniflorum* Lam.). *J Plant Physiol.* 1991; 137: 381-383.
98. Kiyosue T, Yoshiba Y, Yamaguchi-Shinozaki K, Shinozaki K. A nuclear gene, encoding mitochondrial proline dehydrogenase, an enzyme involved in proline metabolism but downregulated by dehydration in Arabidopsis. *The Plant Cell* 1996; 8: 1323-1335.
99. Xin Z, Browse J. Eskimo 1 mutants of Arabidopsis are constitutively freezing tolerant. *Proc Natl Acad Sci USA* 1998; 95: 7799-7804.
100. Paleg LG, Douglas TJ, van Daal A, Keech DB. Proline, betaine and other osmotic solutes protect enzymes against heat inactivation. *Aust J Plant Physiol* 1981; 8: 107-114.
101. Santoro MM, Liu Y, Khan SMA, Hou LX, Bolen DW. Increased thermal stability of Proteins in the presence of naturally occurring osmolytes. *Biochemistry* 1992; 31:5278-5283.
102. Smirnoff N, Cumbes QJ. Hydroxyl radical scavenging activity of compatible solutes. *Phytochemistry.* 1989; 28: 1057-1060.
103. Hagedorn CH, Phang JM. Transfer of reducing equivalents into mitochondria by the interconversions of proline and  $\alpha$ - pyrroline-5-carboxylate. *Arch Biochem Biophysics.* 1983; 225: 95-101.
104. Phang JM. The regulatory functions of proline and pyrroline-5-carboxylic acid. *Curr. Top Cell Regul.* 1985; 25:92-132.
105. Copeland L, Turner JF. The regulation of glycolysis and the pentose-phosphate pathway. In: Stumpf P and Conn EE, eds. *The Biochemistry of Plants*, Vol. 11. New York: Academic Press; 1987: 107-125.
106. Lenzian KJ. Modulation of glucose-6-phosphate dehydrogenase by NADPH, NADP<sup>+</sup> and dithiothreitol at variable NADPH/NADP<sup>+</sup> ratios in an illuminated reconstituted spinach (*Spinacia oleracea* L.) chloroplast system. *Planta.* 1980; 148, 1-6.
107. Shetty K. Role of proline-linked pentose phosphate pathway in biosynthesis of plant phenolics for functional food and environmental applications; A review. *Process Biochem.* 2004; 39: 789-803.
108. Krishnan N, Becker DF. Oxygen reactivity of PutA from *Helicobacter* species and proline-linked oxidative stress. *J. Bacteriol.* 2006; 88: 1227-1235.

109. Adyanthaya I, Kwon Y-I, Apostolidis E, Shetty K. Health benefits of apple phenolics from post-harvest stages for diabetes management. *J. Food Biochemistry*. 2007; Submitted for Publication.
110. Shetty K, Curtis OF, Levin RE, Witkowsky R, Ang W. Prevention of vitrification associated with in vitro shoot culture of oregano (*Origanum vulgare*) by *Pseudomonas* spp. *Journal of Plant Physiology*. 1995; 147: 447-451.
111. Bradford M. A rapid and sensitive method for the quantification of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem*. 1976; 72: 248-254.
112. Oberley LW, Spitz DR. Assay of SOD activity in tumor tissue. In: *Methods in Enzymology*. Academic Press; 1984: 457-461.
113. Costilow RN, Cooper D. Identify of proline dehydrogenase and Pyrroline-5-carboxylic acid reductase in *Clostridium sporogenes*. *J. Bacteriol*. 1978; 134: 139-145.
114. Bregman AA. *Laboratory investigations in cell biology*. NewYork: John Wiley and Sons Inc; 1987.
115. Deutsch J. Glucose-6-phosphate dehydrogenase. In: *Methods of enzymatic analysis*. 3<sup>rd</sup> ed. Verlag chemie; 1983: 191-197.
116. Laloue H, Weber-Lofti F, Lucau-Danila A, Gullema TP. Identification of ascorbate and guaiacol peroxidase in needle chloroplasts of spruce tree. *Plant Physiol. Biochem*. 1997; 35: 341-346.
117. Morales M, Barcelo, A. R. A basic peroxidase isoenzyme from vacuoles and cell walls of *Vitis vinifera*. *Phytochemistry*. 1997; 45: 229-232.
118. Serdula MK, Byres MAH, Simoes E, Mendlein JM, Coates RJ. The association between fruit and vegetable intake and chronic disease risk factors. *Epidemiol*. 1996; 7: 161-165.
119. Adyanthaya I, Kwon Y-I, Apostolidis E, Shetty K. Apple post-harvest preservation linked to phenolics and SOD activity. *J. Food Biochemistry*. 2007; Submitted for Publication.