



University of  
Massachusetts  
Amherst

## **NON-LINEAR EFFECTS IN THE FORMATION OF DNA DAMAGE IN MEDAKA FISH FIBROBLAST CELLS CAUSED BY COMBINED ACTION OF CADMIUM AND IONIZING RADIATION**

Item Type	article;article
Authors	Grygoryev, Dmytro;Moskalenko, Oleksandr;Zimbrick, John D
Download date	2025-01-21 16:35:14
Link to Item	<a href="https://hdl.handle.net/20.500.14394/20546">https://hdl.handle.net/20.500.14394/20546</a>

## NON-LINEAR EFFECTS IN THE FORMATION OF DNA DAMAGE IN MEDAKA FISH FIBROBLAST CELLS CAUSED BY COMBINED ACTION OF CADMIUM AND IONIZING RADIATION

**Dmytro Grygoryev, Oleksandr Moskalenko, John D. Zimbrick** □ Department of Environmental and Radiological Health Sciences, Colorado State University, Fort Collins, CO

□ Ionizing radiation-induced formation of genomic DNA damage can be modulated by nearby chemical species such as heavy metal ions, which can lead to non-linear dose response. To investigate this phenomenon, we studied cell survival and formation of 8-hydroxyguanine (8-OHG) base modifications and double strand breaks (DSB) caused by combined action of cadmium (Cd) and gamma radiation in cultured medaka fish (*Oryzias latipes*) fibroblast cells. Our data show that the introduction of Cd leads to a significant decrease in the fraction of surviving cells and to increased sensitivity of cells to ionizing radiation (IR). Cd also appears to cause non-linear increases in radiation-induced yields of 8-OHG and DSB as dose-yield plots of these lesions exhibit non-linear S-shaped curves with a sharp increase in the yields of lesions in the 10-20  $\mu\text{M}$  range of Cd concentrations. The combined action of ionizing radiation and Cd leads to increased DNA damage formation compared to the effects of the individual stressors. These results are consistent with a hypothesis that the presence of Cd modulates the efficiency of DNA repair systems thus causing increases in radiation-induced DNA damage formation and decreases in cell survival.

*Keywords:* Cadmium, ionizing radiation, DNA, double-strand breaks, 8-hydroxyguanine, Medaka fish cells.

### INTRODUCTION

Relatively few mechanistic studies have been carried out on the interactions of radiation and chemicals in tissues or whole organisms. In part, this is due to the complexity of the changes in cellular processes perturbed by combinations of ionizing radiation and chemicals. How can we approach the design of mechanistic studies on combined exposures in model systems to help elucidate molecular mechanisms underlying a possible variety of additive, synergistic or antagonistic responses? One approach is to select a chemical agent that acts in cells similarly to ionizing radiation. In the present work we selected combinations of cadmium, a soluble metal ion found in numerous waste sites around the United States, and gamma radiation from Cesium-137, a radioactive isotope found in a number of the same waste sites.

Address correspondence to John D. Zimbrick, Department of Environmental and Radiological Health Sciences, Colorado State University, Fort Collins, CO. Email: Zimbrick@colostate.edu

Cd acts via a number of mechanisms, which have also been associated with the actions of ionizing radiation. For example, Cd can activate oncogenes (Fang *et al.* 2002), induce apoptotic cell death (Szuster-Ciesielska *et al.* 2000; Watjen *et al.* 2002; Pulido and Parrish 2003; Jimi *et al.* 2004), induce reactive oxygen radical species (Szuster-Ciesielska *et al.* 2000; McMurray and Tainer 2003; Waisberg *et al.* 2003; Bertin and Averbeck 2006), inhibit DNA repair (Yang *et al.* 1996; Hartwig *et al.* 2002a, b; Hartwig and Schwerdtle 2002; Waalkes 2003; Waisberg *et al.* 2003; Durham and Snow 2006; Giaginis *et al.* 2006), and induce metallothionein (MT) formation in tissues (Chan and Cherian 1992). The diverse mechanisms whereby Cd can induce damage, as well as alter biological responses caused by other contaminants, suggest that Cd can potentially affect all stages of the carcinogenic process, including initiation, promotion and progression (Waalkes 2003) just as has been shown for ionizing radiation.

#### **Genotoxic and genomic studies on combined exposure to ionizing radiation and Cd**

Very little is known about the combined effects of ionizing radiation and metals, such as Cd, on DNA damage and gene expression in mammalian cells or tissues. A recent study (Hornhardt *et al.* 2006) on human lymphoblastoid cells included exposures to combined levels of Cd and Cs gamma-rays. Their data on the formation of micronuclei as a marker suggest that a combination of Cd and radiation had an additive effect on the frequency of micronuclei formation. It was also shown that exposure to 10  $\mu\text{M}$  Cd led to a significant increase in  $\gamma$ -radiation induced 8-OHG formation in human fibroblast GM00637 and HeLa S3 cells by inhibiting the expression of an 8-oxoguanine DNA glycosylase (Youn *et al.* 2005). A synergistic increase in enzyme activities of lactate dehydrogenase, alkaline phosphatase, and acid phosphatase in Wistar rats under combined action of Cd and  $\gamma$ -radiation was also reported (Salovsky *et al.* 1993). It has been suggested that Cd inhibits the repair of potentially lethal damage caused by UV-irradiation in cultured human and simian cells (Nocentini 1987). In combination with gamma irradiation, Cd has been shown to increase the level of metallothioneins in the bone marrow and liver of mice (Koterov *et al.* 1997). Increases in the levels of lipid peroxidation in tissues, accumulation of its end products in biosubstrates, and activation of catalase and peroxidase oxidoreductases in blood of test animals have also been shown (Honcharuk *et al.* 1995).

Cd fed to rats as a dietary supplement slightly enhanced the cytogenetic damage caused by chronic and acute  $\gamma$ -radiation and decreased the cytogenetic adaptive response induced by chronic  $\gamma$ -radiation. (Zaichkina *et al.* 2001). An antagonistic effect of the combined action of Cd and  $\gamma$ -radiation also was reported as mice treated with Cd 48 h prior to irradiation exhibited decreased levels of DNA damage in peripheral blood lym-

phocytes and splenocytes in comparison with untreated mice. (Privezentsev *et al.* 1996). In addition, the pre-irradiation administration of Cd to mice is reported to have accelerated repair processes in the bone marrow and spleen as well as increased the number of leukocytes and thrombocytes in the peripheral blood (Mackova *et al.* 1996).

It is possible that such broad and high Cd activity, associated with the ability of this metal to bind to proteins and DNA and to change the conformation of these macromolecules (Predki and Sarkar 1994; Asmuss *et al.* 2000a; Hartwig 2001; Kopera *et al.* 2004; Witkiewicz-Kucharczyk and Bal 2006), can lead to nonlinear or even hormetic dose-response relationships in combination with ionizing radiation.

In this paper we present the results of some of our initial studies on the combined effects of Cd and ionizing radiation exposures on DNA damage in cultured medaka fish fibroblast cells.

## MATERIALS AND METHODS

### Cell culture treatments

CAB fibroblast medaka fish (*Oryzias latipes*) cell line was provided by Dr. Wendy Kuhne (Colorado State University) with kind permission of Dr. Hiroshi Mitani (The University of Tokyo). Cells were incubated in T75 (Corning, NY) flasks in cell culture medium (L15 Leibovitz Medium (HyClone, Logan, UT), 20% Fetal Bovine Serum (HyClone, Logan, UT), 0.5 U/ml Penicillin – Streptomycin (HyClone, Logan, UT)) with 0 - 50  $\mu\text{M}$   $\text{CdCl}_2$  at 30° C for 24 hrs. After a Cd treatment, they were  $\gamma$ -irradiated at the Colorado State University  $^{137}\text{Cs}$  gamma radiation facility at a dose-rate of 2.5 Gy/min with doses from 0 to 5 Gy for each Cd concentration. After irradiation, cells were immediately placed on ice. For Cd-only treatments cells were incubated with  $\text{CdCl}_2$  for 48 and 72 hours and used immediately. For the colony formation assay approximately 300 cells were plated in each 16mm well in 24-well tissue culture plates (Falcon; Becton Dickinson & Co, Oxnard, CA) and incubated with 0 - 50  $\mu\text{M}$   $\text{CdCl}_2$  at 30° C for 24 hrs. Immediately following this treatment cells were  $\gamma$ -irradiated with doses from 0 to 15 Gy at a dose-rate of 2.5 Gy/min.

### DNA isolation

DNA from the cells was extracted by the chaotropic NaI method (Helbock *et al.* 1998). Approximately  $7.5 \times 10^5$  cell fractions were used for preparation of agarose plugs (Wang *et al.* 1995).

### Colony formation assay

After treatment, cells were incubated for 14 days to allow for the growth of clonogenic colonies. Colonies were fixed with methanol and

treated with Crystal Violet stain. Colonies with 50 or more cells were counted (Herskind *et al.* 2000; Potts *et al.* 2003; Akudugu *et al.* 2004; Nair *et al.* 2004; Youn *et al.* 2005; Franken *et al.* 2006).

#### **Determination of 8-hydroxy-2'-deoxyguanosine (8-OHdG) levels in isolated DNA**


The quantitation of 8-OHdG levels in the isolated DNA samples was carried out via high-performance liquid chromatography with electrochemical detection (HPLC-ECD) (Nicotera and Bardin 1998; Pouget *et al.* 2000). The assay performance was tested by quantitation of 8-OHdG formation in  $\gamma$ -irradiated aqueous solutions of salmon sperm DNA (Sigma, St. Louis). Aqueous solutions of 1 $\mu$ g/ $\mu$ l DNA in 10mM NaClO<sub>4</sub> were  $\gamma$ -irradiated with 5 doses from one to 100 Gy at a dose rate of 30 Gy/hour. After enzymatic digestion of DNA to nucleosides (Helbock *et al.* 1998) HPLC analysis was done with a Waters 2690 Alliance System equipped with an Atlantis<sup>TM</sup> dC<sub>18</sub> reversed-phase column and a Waters 2487 dual  $\lambda$  UV absorbance detector. An ESA Model 5600A CoulArray electrochemical detector with 8 electrochemical cells received the eluent from the UV detector and provided the ECD spectra used in the analysis. For each sample, the amount of DNA injected onto the column was estimated using the UV signal of unmodified nucleosides after appropriate calibration. The amount of 8-OHdG was determined using the ECD signal after appropriate calibration with a commercially available standard (Berry & Associates Inc., Dexter, MI ). The radiation yield of 8-OHdG obtained from the yield – dose dependence was  $1.1 \pm 0.05$  8-OHdG/ $10^5$  dG Gy<sup>-1</sup> and is in good agreement with values found in the literature (Dizdaroglu 1985; Fuciarelli *et al.* 1990; Douki *et al.* 1997; Svoboda and Harms-Ringdahl 1999).

#### **Quantitation of double-strand breaks in isolated DNA**

A CHEF Mapper XA Pulsed Field Electrophoresis System (Bio-Rad, Hercules, CA) was used for measurements of double strand break (DSB) formation. DNA fragments in the size region of 200 kb – 6 Mb were separated according to a previously published protocol (Elia and Nichols 1993). DNA fragments in the 200 bp – 200 kbp region were resolved by a modified method (Carle *et al.* 1986). Quantation of double-strand breaks was done by number average length analysis (Sutherland *et al.* 2003a, b; Gradzka and Iwanenko 2005).

#### **Detection of apoptosis**

Cells were incubated with several Cd concentrations in the 10 – 50  $\mu$ M range at 30°C for 24 hrs. After incubation, cells were harvested and DNA was extracted with an Apoptotic DNA Ladder Extraction kit (BioVision,



**FIGURE 1.** The effect of Cd on the survival of medaka cells: (1) The dependence of cell survival fraction (SF) on Cd concentration. The cells were incubated for 24 hours with various Cd concentrations; The data are fit very well by a quadratic model (Chadwick and Leenhouts 1973; Hethcote *et al.* 1976), reduced chi-square < 0.005; (2) Apoptotic medaka cell DNA fragmentation patterns caused by Cd exposures: Lane (1) - 1 kb DNA Ladder; lane (2) - 100 bp DNA Ladder; lane (3) - Control DNA from the medaka cells without any treatment ; lane (4) - DNA from the medaka cells incubated with CAM; lanes (5) – (9) - DNA from the medaka cells incubated with 10, 20, 30, 40 and 50  $\mu\text{M}$  of Cd correspondingly.

Mountain View, CA) and analyzed by electrophoresis on a 1.2 % agarose gel (Kang *et al.* 2006). DNA extracted from medaka cells treated with Camptothecin (CAM) was used as positive control for apoptotic DNA fragmentation (Tonini *et al.* 2004).


*Statistical analysis.* All data displayed graphically are expressed as mean  $\pm$  SD values obtained from 3-5 repeats. Linear regression analysis and non linear curve fitting were done with Microcal Origin software (OriginLab, Northampton, MS).

## RESULTS AND DISCUSSION

### Effect of Cadmium on Medaka Cell Survival and DNA Damage Formation

We have performed studies of Cd and combined Cd and  $\gamma$ -irradiation effects on medaka CAB (*Oryzias latipes*) fibroblasts. Endpoints measured consisted of cell survival, induction of apoptosis and quantitation of DNA damage markers (8OHdG and double strand break [DSB] formation).

Cell survival data resulting from Cd exposures of medaka fibroblast cells are presented in Figure 1. These data indicate that acute (24 hour) exposures of the cells to Cd concentrations up to 10  $\mu\text{M}$  did not produce statistically significant changes in the survival fraction (SF). However, Cd concentrations in the range of 20 to 50  $\mu\text{M}$  resulted in decreases in SF up to a factor of 30. (Fig. 1 [1]). The data were analyzed by use of a non-linear curve fitting analysis based on a quadratic model (Chadwick and



**FIGURE 2.** The effect of Cd on the yield of DNA damage markers in treated medaka cells: (1) The dependence of the yield of 8OHdG on time of exposure to Cd (lesions per  $\mu\text{M}$ ); (2) The dependence of the yield of DSB on time of exposure to Cd (lesions per  $\mu\text{M}$ ).

Leenhouts 1973; Hethcote *et al.* 1976). The quadratic model provided an accurate (reduced chi-square  $< 0.005$ ) fit to the experimental data.

Gel electrophoresis results (Fig. 1 [2]) show apoptotic DNA fragmentation caused by incubation of the cells with Cd. The fragmentation takes place only if Cd concentrations are higher than  $10 \mu\text{M}$ . DNA extracted from medaka cells treated with Camptothecin (CAM) was used as a positive control for apoptotic DNA fragmentation (Tonini *et al.* 2004).

Incubation of cells with Cd also leads to increases in levels of DNA damage markers (8-hydroxyguanine and DSB). This effect depended not only on concentration but also on exposure time. (Fig. 2).

Quantitation of DNA damage markers - 8-OHG (Fig. 2 [1]) and DSB (Fig. 2 [2]) - in medaka cells indicated that the yields of DNA lesions per Cd concentration unit ( $\mu\text{M}$ ) strongly depended on the exposure time.

Similar cytotoxic and genotoxic effects of Cd have been widely described in the literature (Latinwo *et al.* 1997; Mikhailova *et al.* 1997; McMurray and Tainer 2003; Filipic and Hei 2004; Bertin and Averbek 2006; Martelli *et al.* 2006; Badisa *et al.* 2007). However, the underlying mechanisms that could be responsible for such effects are still not clearly understood. The toxicity produced by Cd exposures could be due to several simultaneous processes such as induction of apoptosis (Szuster-Ciesielska *et al.* 2000; Watjen *et al.* 2002; Pulido and Parrish 2003; Jimi *et al.* 2004), indirect generation of reactive oxygen species (Szuster-Ciesielska *et al.* 2000; Galan *et al.* 2001; McMurray and Tainer 2003; Waisberg *et al.* 2003; Watanabe *et al.* 2003), and interference with DNA repair (Yang *et al.* 1996; Asmuss *et al.* 2000b; Hartwig *et al.* 2002b; Hartwig and Schwerdtle 2002; Jin *et al.* 2003; Potts and Pasqualotto 2003; Potts *et*

*al.* 2003; Waalkes 2003; Waisberg *et al.* 2003; Youn *et al.* 2005; Durham and Snow 2006; Giaginis *et al.* 2006). Moreover, Cd exposure can affect cell cycle progression, proliferation, differentiation, and DNA replication (Dong *et al.* 2001; Fang *et al.* 2002; Yang *et al.* 2004; Oh and Lim 2006).

The data presented in Fig. 1 indicate that Cd exposures induced apoptosis and reduction in SF in medaka fibroblast cells in a concentration-dependent manner. The induction of apoptotic cell death by Cd has also been demonstrated in some other mammalian and fish cell lines. (Achanzar *et al.* 2000; Szuster-Ciesielska *et al.* 2000; Lag *et al.* 2002; Watjen *et al.* 2002; Pulido and Parrish 2003; Jimi *et al.* 2004; Risso-de Faverney *et al.* 2004; Gonzalez *et al.* 2006). Such induction can occur through two main pathways: extrinsic and intrinsic. The extrinsic pathway is triggered by ligand-induced activation of the death receptors and the intrinsic pathway is induced by cellular stress signals (Watjen *et al.* 2002).


The presence of a shoulder on the survival-dose curve (Fig.1 [1]) and the absence of detectable apoptosis (Fig. 1 [2]) at Cd concentrations up to 10  $\mu\text{M}$  may be consistent with the involvement of metallothioneins. Metallothioneins – cysteine-rich heavy metal-binding proteins – can play a protective role against Cd toxicity (Andrews 2000). Cd-induced overexpression of metallothioneins has been detected in mammalian cells (Li *et al.* 2005) as well as in fish tissues (Woo *et al.* 2006). Overexpression of metallothioneins in the cells was shown to reduce the effects of Cd at lower (up to 10  $\mu\text{M}$ ) but not at higher (20–50  $\mu\text{M}$ ) concentrations (Li *et al.* 2005). Further, low concentrations of Cd can produce stimulation and hormesis effects under certain conditions (Weis and Weis 1986; von Zglinicki *et al.* 1992; Damelin *et al.* 2000; Gaddipati *et al.* 2003; Fulladosa *et al.* 2005).

The dose-dependent accumulation of unrepaired DNA damage and the concomitant increases in the damage yields per  $\mu\text{M}$  of Cd in the Cd-exposed cells with time (Fig. 2) appear to be consistent with a mechanism involving increases in reactive oxygen species (ROS) production and/or suppression of DNA repair by Cd similarly to previously reported results (Dally and Hartwig 1997; Mikhailova *et al.* 1997; Filipic and Hei 2004; Pan and Zhang 2006; Emmanouil *et al.* 2007).

Although Cd does not generate free radicals directly (Lloyd *et al.* 1998; Valko *et al.* 2005), it can do so indirectly by facilitating increases in lipid peroxidation (Filipic and Hei 2004), decreases in intracellular glutathione content, decreases in activities of cellular antioxidant enzymes including superoxide dismutase, peroxidase and catalase, (Hussain *et al.* 1987; Waisberg *et al.* 2003), and increases in iron and copper concentrations (Koizumi *et al.* 1992; Casalino *et al.* 1997), all of which can lead to increasing levels of ROS.

On the other hand, Cd has been shown to inhibit DNA repair processes such as base-excision repair (BER) (Dally and Hartwig 1997; Potts *et al.*





**FIGURE 3.** The effect of combined action of Cd and  $\gamma$ -radiation on the survival of medaka cells: (1) The dependence of the cell survival fraction (SF) on radiation dose after 24 hours incubation with various Cd concentrations. The data are fit very well by a quadratic model (Chadwick and Leenhouts 1973; Hethcote *et al.* 1976), reduced chi-square  $< 0.07$ ; (2) The effect of Cd concentration on radiation Lethal Dose (LD<sub>50</sub>).


2001; Potts *et al.* 2003; Youn *et al.* 2005), nucleotide-excision repair (NER), and mismatch repair (MMR) (Jin *et al.* 2003). Additionally, Cd inhibits repair of oxidative DNA damage by inhibiting the expression of 8-oxoguanine DNA glycosylase (Potts and Pasqualotto 2003; Potts *et al.* 2003; Youn *et al.* 2005). The data presented in the next section illustrate the interference of Cd with DNA repair systems.

### Combined Effects of Cadmium and Ionizing Radiation on Cell Survival and Formation of DNA Damage


Cell survival data obtained from combined Cd and  $\gamma$ -irradiation exposures of medaka fibroblast cells are presented in Figure 3. It can be seen that a prior 24 hour exposure to Cd significantly increases the sensitivity of cells to  $\gamma$ -radiation (Fig. 3 [1]). The data were plotted by use of a non-linear curve fitting analysis based on a quadratic model (Chadwick and Leenhouts 1973; Hethcote *et al.* 1976). The quadratic model provided an accurate (reduced chi-square  $< 0.07$ ) fit to the experimental data. As Cd exposure concentrations are increased up to 30  $\mu\text{M}$ , the median lethal radiation dose (LD<sub>50</sub>) decreases by a factor of 3 (Fig. 3 [2]). Further increases in the concentration of Cd from 30 to 50  $\mu\text{M}$  no longer affect the LD<sub>50</sub>.

Medaka cells treated with combined Cd and  $\gamma$ -irradiation exhibit increases in levels of DNA damage markers compared with the levels observed after exposures to single contaminants. These increases can be seen in the bar graphs presented in Figures 4(1) and 4(2) as well as in the three-dimensional graphs shown in Figures 5(1) and 5(2).

Combined Cd/IR effects in Medaka cell DNA damage




**FIGURE 4.** The effect of combined Cd and  $\gamma$ -radiation exposures on the levels of DNA damage markers in medaka cells: (1) Quantitation of 8OHdG lesions in medaka cells produced in control cells incubated for 24 hours; cells incubated for 24 hours with Cd (20  $\mu$ M); cells irradiated with  $\gamma$ -rays (1 Gy); and cells treated with a combination of Cd and  $\gamma$ -rays; (2) Quantitation of DSB lesions produced in cells treated as shown on graph. The control is the background level of DSB frequency in the medaka cell DNA.



**FIGURE 5.** The effect of Cd exposures on the levels of DNA damage markers caused by ionizing radiation in treated medaka cells. The cells were incubated for 24 hours with various Cd concentrations prior to irradiation. (1) The dependence of the 8OHdG/ $10^5$ dG level on radiation dose and Cd concentration: X-axis - Cd concentration in  $\mu$ M; Y-axis - radiation dose in Gy; Z-axis - 8OHdG/ $10^5$ dG level. (2) The dependence of the DSB/Mbp on radiation dose and Cd concentration: X-axis - Cd concentration in  $\mu$ M; Y-axis - radiation dose in Gy; Z-axis - DSB/Mbp.

In the Graphs shown in Figure 5, it can be seen that the slopes of the dose-response curves for the formation of both 8OHdG and DSB increase with increasing Cd concentrations.

The radiation-induced yields of 8OHdG and DSB in medaka cell DNA were determined by linear regression analyses from the slopes of the lin-



**FIGURE 6.** The effect of Cd exposure on the yield of DNA damage markers in treated medaka cells. The cells were incubated for 24 hours with various Cd concentrations prior to irradiation (1) The dependence of the radiation yield of 8OHdG (lesions per Gy) on Cd concentration.; (2) The dependence of the DSB radiation yield (lesions per Gy) on Cd concentration.

ear portions of the dose–response graphs for various Cd concentrations (Fig. 6). These dose-response graphs show that the Cd concentration dependence has a nonlinear character with curves for both 8OHdG and DSB yields being similar in shape. Increases in Cd concentration up to 10 μM do not produce changes in the yields of DNA damage markers. In the concentration range from 10 to 20 μM, the yields of markers increase sharply by a maximum factor of approximately two. No further significant changes are found for Cd concentrations above 20 μM.

As mentioned above, Cd can impair the efficiency of DNA repair systems, which will lead to increases in the sensitivity of cells to ionizing radiation and in the yields of radiation-induced DNA damage as the concentration of Cd is increased. (Fig. 3 [2]; Fig. 5). Higher yields of DNA damage observed after a combined exposure of cells to Cd and ionizing radiation compared with the yields observed after exposures to the single agents may be consistent with the action of two parallel processes - the indirect formation of additional ROS by Cd along with the direct radiation-induced formation of ROS and a greater reduction in efficiency of repair of DNA damage caused by both radiation and Cd-generated ROS.

It has been shown that Cd can reduce or even inhibit DNA binding of a number of proteins involved in DNA repair processes (Bertin and Averbeck 2006) such as xeroderma pigmentosum A (XPA) protein required for recognition of DNA lesions in NER system (Hartmann and Hartwig 1998; Hartwig 1998; Hartwig *et al.* 2002a; Hartwig and Schwerdtle 2002); formamidopyrimidine glycosylase (Fpg) that recognizes and repairs 8-oxo-7,8-dihydroguanine base modifications in the Base Excision

Repair (BER) system in *Escherichia coli* (O'Connor *et al.* 1993), a mammalian homologue of Fpg (Potts *et al.* 2003) 8-oxoguanine-DNA glycosylase-1 (OGG1), and specificity protein-1 (Sp1) transcription factor (Watkin *et al.* 2003). The mechanism underlying the ability of Cd to inhibit binding of DNA repair proteins is consistent with displacement of zinc (Zn) by Cd in the Zn finger structure of these proteins (Predki and Sarkar 1994; Asmuss *et al.* 2000a; Hartwig 2001; Kopera *et al.* 2004; Witkiewicz-Kucharczyk and Bal 2006). Also, Bertin and Averbeck (Bertin and Averbeck 2006) have noted that proteins thought to be involved in homologous recombination (HR) and non-homologous end joining (NHEJ) such as BRCA1 (Miki *et al.* 1994), ATR (Villard *et al.* 1997), and DNA ligase III alpha and beta (Martin and MacNeill 2002) can be inhibited by Cd. It has not yet been demonstrated (Bertin and Averbeck 2006) that Cd can inhibit the repair of DNA DSBs in mammalian cells or tissues, although such inhibition has been observed in the repair of single-strand breaks (SSB) induced by external agents (Littlefield *et al.* 1994; Lynn *et al.* 1997; Pruski and Dixon 2002; Fatur *et al.* 2003; Hengstler *et al.* 2003). In contrast, inhibition of DNA SSB repair in HeLa cells by Cd was not detected (Dally and Hartwig 1997).

We have demonstrated in this study that Cd not only causes the formation of DNA DSB in medaka cells (Fig. 4 [2]) but also affects DSB repair (Fig. 6 [2]). In the latter figures, the dose-response curves for radiation-induced yields versus Cd concentration are non-linear for both 8OHdG and DSB yields up to 10  $\mu\text{M}$  of Cd followed by sharp increases in yields in the narrow concentration range from 10  $\mu\text{M}$  to 20  $\mu\text{M}$  and then followed by saturation of the yields in the 20-50  $\mu\text{M}$  interval. The finding that both dose-response relationships have similar character suggests that the underlying mechanisms can be similar or the same. For example, one such mechanism could involve the competitive displacement (Scatchard 1949) of Zn by Cd in the Zn finger structures. As stated previously, over-expression of metallothioneins can reduce the effects of Cd at relatively low concentrations up to 10  $\mu\text{M}$  Cd (Li *et al.* 2005) and can be responsible for the 0 – 10  $\mu\text{M}$  Cd threshold observed in Fig. 6.

## CONCLUSIONS

Summarizing the data presented above we conclude that Cd exposures significantly reduce the survival of medaka fish fibroblast cells and cause concentration-dependent increases in DNA damage markers such as 8OHG and DSB. Further, Cd decreases the observed  $\text{LD}_{50}$  of  $\gamma$ -irradiated cells and increases the radiation yields of the DNA damage markers. The dose-response curves for these effects are nonlinear. The mechanisms underlying these dose-responses likely involve several processes including the interaction of Cd with metallothioneins and competitive binding of Cd to proteins involved in DNA repair systems.

## ACKNOWLEDGMENTS

This work was supported by NIH/NCI T32 Training Grant CA – 09236-29

## REFERENCES

- Achanzar WE, Achanzar KB, Lewis JG, Webber MM and Waalkes MP. 2000. Cadmium induces c-myc, p53, and c-jun expression in normal human prostate epithelial cells as a prelude to apoptosis. *Toxicol Appl Pharmacol* 164:291-300
- Akudugu JM, Bell RS, Catton C, Davis AM, O'Sullivan B, Waldron J, Wunder JS and Hill RP. 2004. Clonogenic survival and cytokinesis-blocked binucleation of skin fibroblasts and normal tissue complications in soft tissue sarcoma patients treated with preoperative radiotherapy. *Radiother Oncol* 72:103-112
- Andrews GK. 2000. Regulation of metallothionein gene expression by oxidative stress and metal ions. *Biochem Pharmacol* 59:95-104
- Asmuss M, Mullenders LH, Eker A and Hartwig A. 2000a. Differential effects of toxic metal compounds on the activities of Fpg and XPA, two zinc finger proteins involved in DNA repair. *Carcinogenesis* 21:2097-2104
- Asmuss M, Mullenders LH and Hartwig A. 2000b. Interference by toxic metal compounds with isolated zinc finger DNA repair proteins. *Toxicol Lett* 112-113:227-231
- Badisa VL, Latinwo LM, Odewumi CO, Ikediobi CO, Badisa RB, Ayuk-Takem LT, Nwoga J and West J. 2007. Mechanism of DNA damage by cadmium and interplay of antioxidant enzymes and agents. *Environ Toxicol* 22:144-151
- Bertin G and Averbeck D. 2006. Cadmium: cellular effects, modifications of biomolecules, modulation of DNA repair and genotoxic consequences (a review). *Biochimie* 88: 1549-1559
- Carle GF, Frank M and Olson MV. 1986. Electrophoretic separations of large DNA molecules by periodic inversion of the electric field. *Science* 232:65-68
- Casalino E, Sblano C and Landriscina C. 1997. Enzyme activity alteration by cadmium administration to rats: the possibility of iron involvement in lipid peroxidation. *Arch Biochem Biophys* 346:171-179
- Chadwick KH and Leenhouts HP. 1973. A molecular theory of cell survival. *Phys Med Biol* 18:78-87
- Chan HM and Cheria MG. 1992. Protective roles of metallothionein and glutathione in hepatotoxicity of cadmium. *Toxicology* 72:281-290
- Dally H and Hartwig A. 1997. Induction and repair inhibition of oxidative DNA damage by nickel(II) and cadmium(II) in mammalian cells. *Carcinogenesis* 18:1021-1026
- Damelin LH, Vokes S, Whitcutt JM, Damelin SB and Alexander JJ. 2000. Hormesis: a stress response in cells exposed to low levels of heavy metals. *Hum Exp Toxicol* 19:420-430
- Dizdaroglu M. 1985. Formation of an 8-hydroxyguanine moiety in deoxyribonucleic acid on gamma-irradiation in aqueous solution. *Biochemistry* 24:4476-4481
- Dong S, Shen HM and Ong CN. 2001. Cadmium-induced apoptosis and phenotypic changes in mouse thymocytes. *Mol Cell Biochem* 222:11-20
- Douki T, Martini R, Ravanat JL, Turesky RJ and Cadet J. 1997. Measurement of 2,6-diamino-4-hydroxy-5-formamidopyrimidine and 8-oxo-7,8-dihydroguanine in isolated DNA exposed to gamma radiation in aqueous solution. *Carcinogenesis* 18:2385-2391
- Durham TR and Snow ET. 2006. Metal ions and carcinogenesis. *Exs*:97-130
- Elia MC and Nichols WW. 1993. Application of programmable, autonomously controlled electrode (PACE) technology to the development of an improved pulsed field gel electrophoresis assay for DNA double-strand breaks in mammalian cells. *Int J Radiat Biol* 63:7-11
- Emmanouil C, Sheehan TM and Chipman JK. 2007. Macromolecule oxidation and DNA repair in mussel (*Mytilus edulis* L.) gill following exposure to Cd and Cr(VI). *Aquat Toxicol* 82:27-35
- Fang MZ, Mar W and Cho MH. 2002. Cadmium affects genes involved in growth regulation during two-stage transformation of Balb/3T3 cells. *Toxicology* 177:253-265
- Fatur T, Lah TT and Filipic M. 2003. Cadmium inhibits repair of UV-, methyl methanesulfonate- and N-methyl-N-nitrosourea-induced DNA damage in Chinese hamster ovary cells. *Mutat Res* 529:109-116

- Filipic M and Hei TK. 2004. Mutagenicity of cadmium in mammalian cells: implication of oxidative DNA damage. *Mutat Res* 546:81-91
- Franken NA, Rodermond HM, Stap J, Haveman J and van Bree C. 2006. Clonogenic assay of cells in vitro. *Nat Protoc* 1:2315-2319
- Fuciarelli AF, Wegher BJ, Blakely WF and Dizdaroglu M. 1990. Yields of radiation-induced base products in DNA: effects of DNA conformation and gassing conditions. *Int J Radiat Biol* 58:397-415
- Fulladosa E, Murat JC and Villaescusa I. 2005. Effect of cadmium(II), chromium(VI), and arsenic(V) on long-term viability- and growth-inhibition assays using *Vibrio fischeri* marine bacteria. *Arch Environ Contam Toxicol* 49:299-306
- Gaddipati J, Rajeshkumar N, Grove J, Maharaj S, Centeno J, Maheshwari R and BJ. W. 2003. Low-Dose Cadmium Exposure Reduces Human Prostate Cell Transformation in Culture and Up-Regulates Metallothionein and MT-1G mRNA. *Nonlinearity in Biology, Toxicology, and Medicine* 1:199 - 212
- Galan A, Garcia-Bermejo L, Troyano A, Vilaboa NE, Fernandez C, de Blas E and Aller P. 2001. The role of intracellular oxidation in death induction (apoptosis and necrosis) in human promonocytic cells treated with stress inducers (cadmium, heat, X-rays). *Eur J Cell Biol* 80:312-320
- Giaginis C, Gatzidou E and Theocharis S. 2006. DNA repair systems as targets of cadmium toxicity. *Toxicol Appl Pharmacol* 213:282-290
- Gonzalez P, Baudrimont M, Boudou A and Bourdineaud JP. 2006. Comparative effects of direct cadmium contamination on gene expression in gills, liver, skeletal muscles and brain of the zebrafish (*Danio rerio*). *Biometals* 19:225-235
- Gradzka I and Iwanenko T. 2005. A non-radioactive, PFGE-based assay for low levels of DNA double-strand breaks in mammalian cells. *DNA Repair (Amst)* 4:1129-1139
- Hartmann M and Hartwig A. 1998. Disturbance of DNA damage recognition after UV-irradiation by nickel(II) and cadmium(II) in mammalian cells. *Carcinogenesis* 19:617-621
- Hartwig A. 1998. Carcinogenicity of metal compounds: possible role of DNA repair inhibition. *Toxicol Lett* 102-103:235-239
- Hartwig A. 2001. Zinc finger proteins as potential targets for toxic metal ions: differential effects on structure and function. *Antioxid Redox Signal* 3:625-634
- Hartwig A, Asmuss M, Blessing H, Hoffmann S, Jahnke G, Khandelwal S, Pelzer A and Burkle A. 2002a. Interference by toxic metal ions with zinc-dependent proteins involved in maintaining genomic stability. *Food Chem Toxicol* 40:1179-1184
- Hartwig A, Asmuss M, Ehleben I, Herzer U, Kostelac D, Pelzer A, Schwerdtle T and Burkle A. 2002b. Interference by toxic metal ions with DNA repair processes and cell cycle control: molecular mechanisms. *Environ Health Perspect* 110 Suppl 5:797-799
- Hartwig A and Schwerdtle T. 2002. Interactions by carcinogenic metal compounds with DNA repair processes: toxicological implications. *Toxicol Lett* 127:47-54
- Helbock HJ, Beckman KB, Shigenaga MK, Walter PB, Woodall AA, Yeo HC and Ames BN. 1998. DNA oxidation matters: the HPLC-electrochemical detection assay of 8-oxo-deoxyguanosine and 8-oxo-guanine. *Proc Natl Acad Sci U S A* 95:288-293
- Hengstler JG, Bolm-Audorff U, Faldum A, Janssen K, Reifnath M, Gotte W, Jung D, Mayer-Popken O, Fuchs J, Gebhard S, Bienfait HG, Schlink K, Dietrich C, Faust D, Epe B and Oesch F. 2003. Occupational exposure to heavy metals: DNA damage induction and DNA repair inhibition prove co-exposures to cadmium, cobalt and lead as more dangerous than hitherto expected. *Carcinogenesis* 24:63-73
- Herskind C, Johansen J, Bentzen SM, Overgaard M, Overgaard J, Bamberg M and Rodemann HP. 2000. Fibroblast differentiation in subcutaneous fibrosis after postmastectomy radiotherapy. *Acta Oncol* 39:383-388
- Hethcote HW, McLarty JW and Thames HD, Jr. 1976. Comparison of mathematical models for radiation fractionation. *Radiat Res* 67:387-407
- Honcharuk Ie H, Bardov VH, Kartysh AP, Iavorov's'yi, Korshun MM and Verpiv's'kyi OM. 1995. An experimental study of the mechanism of the combined action on the body of ionizing radiation, pesticides, nitrates and lead and cadmium salts. *Lik Sprava* 5-6:7-12
- Hornhardt S, Gomolka M, Walsh L and Jung T. 2006. Comparative investigations of sodium arsenite, arsenic trioxide and cadmium sulphate in combination with gamma-radiation on apoptosis, micronuclei induction and DNA damage in a human lymphoblastoid cell line. *Mutat Res* 600:165-176

- Hussain T, Shukla GS and Chandra SV. 1987. Effects of cadmium on superoxide dismutase and lipid peroxidation in liver and kidney of growing rats: in vivo and in vitro studies. *Pharmacol Toxicol* 60:355-358
- Jimi S, Uchiyama M, Takaki A, Suzumiya J and Hara S. 2004. Mechanisms of cell death induced by cadmium and arsenic. *Ann NY Acad Sci* 1011:325-331
- Jin YH, Clark AB, Slebos RJ, Al-Refai H, Taylor JA, Kunkel TA, Resnick MA and Gordenin DA. 2003. Cadmium is a mutagen that acts by inhibiting mismatch repair. *Nat Genet* 34:326-329
- Kang SG, Brown AL and Chung JH. 2006. Oxygen tension regulates the stability of insulin receptor substrate (IRS)-1 through caspase-mediated cleavage. *J Biol Chem* 282:6090-6097
- Koizumi T, Li ZG and Tatsumoto H. 1992. DNA damaging activity of cadmium in Leydig cells, a target cell population for cadmium carcinogenesis in the rat testis. *Toxicol Lett* 63:211-220
- Kopera E, Schwerdtle T, Hartwig A and Bal W. 2004. Co(II) and Cd(II) substitute for Zn(II) in the zinc finger derived from the DNA repair protein XPA, demonstrating a variety of potential mechanisms of toxicity. *Chem Res Toxicol* 17:1452-1458
- Koterov AN, Trebenok ZA, Pushkareva NB and Nikol'skii AV. 1997. The diverse effects of the combined action of cadmium chloride and ionizing radiation on the content of metallothioneins in mouse bone marrow and liver. *Radiats Biol Radioecol* 37:196-201
- Lag M, Westly S, Lerstad T, Bjornsrud C, Refsnes M and Schwarze PE. 2002. Cadmium-induced apoptosis of primary epithelial lung cells: involvement of Bax and p53, but not of oxidative stress. *Cell Biol Toxicol* 18:29-42
- Latinwo LM, Ikediobi CO, Singh NP, Sponholtz G, Fasanya C and Riley L. 1997. Comparative studies of in vivo genotoxic effects of cadmium chloride in rat brain, kidney and liver cells. *Cell Mol Biol (Noisy-le-grand)* 43:203-210
- Li J, Liu Y and Ru B. 2005. Effect of metallothionein on cell viability and its interactions with cadmium and zinc in HEK293 cells. *Cell Biol Int* 29:843-848
- Littlefield NA, Hass BS, James SJ and Poirier LA. 1994. Protective effect of magnesium on DNA strand breaks induced by nickel or cadmium. *Cell Biol Toxicol* 10:127-135
- Lloyd DR, Carmichael PL and Phillips DH. 1998. Comparison of the formation of 8-hydroxy-2'-deoxyguanosine and single- and double-strand breaks in DNA mediated by fenton reactions. *Chem Res Toxicol* 11:420-427
- Lynn S, Lai HT, Kao SM, Lai J and Jan KY. 1997. Cadmium inhibits DNA strand break rejoining in methyl methanesulfonate-treated CHO-K1 cells. *Toxicol Appl Pharmacol* 144:171-176
- Mackova NO, Lenikova S, Fedorocko P, Brezani P and Fedorockova A. 1996. Effects of cadmium on haemopoiesis in irradiated and non-irradiated mice: 2. Relationship to the number of circulating blood cells and haemopoiesis. *Physiol Res* 45:101-106
- Martelli A, Rousselet E, Dycke C, Bouron A and Moulis JM. 2006. Cadmium toxicity in animal cells by interference with essential metals. *Biochimie* 88:1807-1814
- Martin IV and MacNeill SA. 2002. ATP-dependent DNA ligases. *Genome Biol* 3: reviews3005.1-3005.7
- McMurray CT and Tainer JA. 2003. Cancer, cadmium and genome integrity. *Nat Genet* 34:239-241
- Mikhailova MV, Littlefield NA, Hass BS, Poirier LA and Chou MW. 1997. Cadmium-induced 8-hydroxydeoxyguanosine formation, DNA strand breaks and antioxidant enzyme activities in lymphoblastoid cells. *Cancer Lett* 115:141-148
- Miki Y, Swensen J, Shattuck-Eidens D, Futreal PA, Harshman K, Tavtigian S, Liu Q, Cochran C, Bennett LM, Ding W, Bell R, Rosenthal J, Hussey C, Tran T, McClure M, Frye C, Hattier T, Phelps R, Haugen-Strano A, Katcher H, Yakumo K, Gholami Z, Shaffer D, Stone S, Bayer S, Wray C, Bogden R, Dayananth P, Ward J, Tonin P, Narod S, Bristow PK, Norris FH, Helvering L, Morrison P, Rostock P, Lai M, Barrett JC, Lewis C, Neuhausen S, Cannon-Albright L, Goldgar D, Wiseman R, Kamb A and Skolnick MH. 1994. A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1. *Science* 266:66-71
- Nair HK, Rao KV, Aalinkel R, Mahajan S, Chawda R and Schwartz SA. 2004. Inhibition of prostate cancer cell colony formation by the flavonoid quercetin correlates with modulation of specific regulatory genes. *Clin Diagn Lab Immunol* 11:63-69
- Nicotera TM and Bardin S. 1998. Electrochemical detection of 8-hydroxy-2-deoxyguanosine levels in cellular DNA. *Methods Mol Biol* 108:181-190
- Nocentini S. 1987. Inhibition of DNA replication and repair by cadmium in mammalian cells. Protective interaction of zinc. *Nucleic Acids Res* 15:4211-4225

- O'Connor TR, Graves RJ, de Murcia G, Castaing B and Laval J. 1993. Fpg protein of *Escherichia coli* is a zinc finger protein whose cysteine residues have a structural and/or functional role. *J Biol Chem* 268:9063-9070
- Oh SH and Lim SC. 2006. A rapid and transient ROS generation by cadmium triggers apoptosis via caspase-dependent pathway in HepG2 cells and this is inhibited through N-acetylcysteine-mediated catalase upregulation. *Toxicol Appl Pharmacol* 212:212-223
- Pan L and Zhang H. 2006. Metallothionein, antioxidant enzymes and DNA strand breaks as biomarkers of Cd exposure in a marine crab, *Charybdis japonica*. *Comp Biochem Physiol C Toxicol Pharmacol* 144:67-75
- Potts JM and Pasqualotto FF. 2003. Seminal oxidative stress in patients with chronic prostatitis. *Andrologia* 35:304-308
- Potts RJ, Bespalov IA, Wallace SS, Melamede RJ and Hart BA. 2001. Inhibition of oxidative DNA repair in cadmium-adapted alveolar epithelial cells and the potential involvement of metallothionein. *Toxicology* 161:25-38
- Potts RJ, Watkin RD and Hart BA. 2003. Cadmium exposure down-regulates 8-oxoguanine DNA glycosylase expression in rat lung and alveolar epithelial cells. *Toxicology* 184:189-202
- Pouget JP, Douki T, Richard MJ and Cadet J. 2000. DNA damage induced in cells by gamma and UVA radiation as measured by HPLC/GC-MS and HPLC-EC and Comet assay. *Chem Res Toxicol* 13:541-549
- Predki PF and Sarkar B. 1994. Metal replacement in "zinc finger" and its effect on DNA binding. *Environ Health Perspect* 102 Suppl 3:195-198
- Privezentsev KV, Sirota NP and Gaznev AI. 1996. Effects of combined action of Cd and gamma radiation on DNA damage and repair in lymphoid tissues of mice. *Radiats Biol Radioecol* 36:234-240
- Pruski AM and Dixon DR. 2002. Effects of cadmium on nuclear integrity and DNA repair efficiency in the gill cells of *Mytilus edulis* L. *Aquat Toxicol* 57:127-137
- Pulido MD and Parrish AR. 2003. Metal-induced apoptosis: mechanisms. *Mutat Res* 533:227-241
- Risso-de Faverney C, Orsini N, de Sousa G and Rahmani R. 2004. Cadmium-induced apoptosis through the mitochondrial pathway in rainbow trout hepatocytes: involvement of oxidative stress. *Aquat Toxicol* 69:247-258
- Salovsky P, Shopova V, Dancheva V, Marev R and Pandurska A. 1993. Enhancement of the pneumotoxic effect of cadmium acetate by ionizing radiation in the rat. *Environ Health Perspect* 101 Suppl 2:269-274
- Scatchard G. 1949. The attraction of proteins for small molecules and ions. *Annals of the New York Academy of Sciences* 51:660-672
- Sutherland BM, Bennett PV, Georgakilas AG and Sutherland JC. 2003a. Evaluation of number average length analysis in quantifying double strand breaks in genomic DNAs. *Biochemistry* 42:3375-3384
- Sutherland BM, Georgakilas AG, Bennett PV, Laval J and Sutherland JC. 2003b. Quantifying clustered DNA damage induction and repair by gel electrophoresis, electronic imaging and number average length analysis. *Mutat Res* 531:93-107
- Svoboda P and Harms-Ringdahl M. 1999. Protection or sensitization by thiols or ascorbate in irradiated solutions of DNA or deoxyguanosine. *Radiat Res* 151:605-616
- Szuster-Ciesielska A, Stachura A, Slotwinska M, Kaminska T, Sniezko R, Paduch R, Abramczyk D, Filar J and Kandefer-Szerszen M. 2000. The inhibitory effect of zinc on cadmium-induced cell apoptosis and reactive oxygen species (ROS) production in cell cultures. *Toxicology* 145:159-171
- Tonini T, Gabellini C, Bagella L, D'Andrilli G, Masciullo V, Romano G, Scambia G, Zupi G and Giordano A. 2004. pRb2/p130 decreases sensitivity to apoptosis induced by camptothecin and doxorubicin but not by taxol. *Clin Cancer Res* 10:8085-8093
- Valko M, Morris H and Cronin MT. 2005. Metals, toxicity and oxidative stress. *Curr Med Chem* 12:1161-1208
- Villard L, Lossi AM, Cardoso C, Proud V, Chiaroni P, Colleaux L, Schwartz C and Fontes M. 1997. Determination of the genomic structure of the XNP/ATRAX gene encoding a potential zinc finger helicase. *Genomics* 43:149-155
- von Zglinicki T, Edwall C, Ostlund E, Lind B, Nordberg M, Ringertz NR and Wroblewski J. 1992. Very low cadmium concentrations stimulate DNA synthesis and cell growth. *J Cell Sci* 103 (Pt 4):1073-1081
- Waalkes MP. 2003. Cadmium carcinogenesis. *Mutat Res* 533:107-120



- Waisberg M, Joseph P, Hale B and Beyersmann D. 2003. Molecular and cellular mechanisms of cadmium carcinogenesis. *Toxicology* 192:95-117
- Wang J, Delgado DA and Deen DF. 1995. Use of pulsed-field gel electrophoresis to investigate factors influencing the measurement of DNA double-strand breaks in human brain tumour specimens. *Int J Radiat Biol* 67:153-160
- Watanabe M, Henmi K, Ogawa K and Suzuki T. 2003. Cadmium-dependent generation of reactive oxygen species and mitochondrial DNA breaks in photosynthetic and non-photosynthetic strains of *Euglena gracilis*. *Comp Biochem Physiol C Toxicol Pharmacol* 134:227-234
- Watjen W, Cox M, Biagioli M and Beyersmann D. 2002. Cadmium-induced apoptosis in C6 glioma cells: mediation by caspase 9-activation. *Biometals* 15:15-25
- Watkin RD, Nawrot T, Potts RJ and Hart BA. 2003. Mechanisms regulating the cadmium-mediated suppression of Sp1 transcription factor activity in alveolar epithelial cells. *Toxicology* 184:157-178
- Weis P and Weis JS. 1986. Cadmium acclimation and hormesis in *Fundulus heteroclitus* during fin regeneration. *Environ Res* 39:356-363
- Witkiewicz-Kucharczyk A and Bal W. 2006. Damage of zinc fingers in DNA repair proteins, a novel molecular mechanism in carcinogenesis. *Toxicol Lett* 162:29-42
- Woo S, Yum S, Jung JH, Shim WJ, Lee CH and Lee TK. 2006. Heavy Metal-Induced Differential Gene Expression of Metallothionein in Javanese Medaka, *Oryzias javanicus*. *Mar Biotechnol (NY)* 8:654-662
- Yang PM, Chiu SJ, Lin KA and Lin LY. 2004. Effect of cadmium on cell cycle progression in Chinese hamster ovary cells. *Chem Biol Interact* 149:125-136
- Yang SW, Becker FF and Chan JY. 1996. Inhibition of human DNA ligase I activity by zinc and cadmium and the fidelity of ligation. *Environ Mol Mutagen* 28:19-25
- Youn CK, Kim SH, Lee DY, Song SH, Chang IY, Hyun JW, Chung MH and You HJ. 2005. Cadmium down-regulates human OGG1 through suppression of Sp1 activity. *J Biol Chem* 280:25185-25195
- Zaichkina SI, Rozanova OM, Aptikaeva GF, Akhmadieva A, Klovov D and Smirnova EN. 2001. Induction of cytogenetic damages by combined action of heavy metal salts, chronic and acute gamma irradiation in bone marrow cells of mice and rats. *Radiat Biol Radioecol* 41:514-518