



University of
Massachusetts
Amherst

Energy criterion for the spectral stability of discrete breathers

Item Type	article;article
Authors	Kevrekidis, Panos;Cuevas-Maraver, Jesus;Pelinovsky, Dmitry
Download date	2024-11-03 08:42:26
Link to Item	https://hdl.handle.net/20.500.14394/34341

University of Massachusetts Amherst

From the Selected Works of Panos Kevrekidis

2016

Energy criterion for the spectral stability of discrete breathers

Panos Kevrekidis

Jesus Cuevas-Maraver

Dmitry Pelinovsky



Available at: https://works.bepress.com/panos_kevrekidis/401/

An energy criterion for the spectral stability of discrete breathers

Panayotis G. Kevrekidis

Department of Mathematics and Statistics, University of Massachusetts, Amherst, MA 01003-9305, USA

Jesús Cuevas–Maraver

Grupo de Física No Lineal, Departamento de Física Aplicada I,

Universidad de Sevilla. Escuela Politécnica Superior, C/ Virgen de África, 7, 41011-Sevilla, Spain

Instituto de Matemáticas de la Universidad de Sevilla (IMUS). Edificio Celestino Mutis. Avda. Reina Mercedes s/n, 41012-Sevilla, Spain

Dmitry E. Pelinovsky

Department of Mathematics, McMaster University, Hamilton, Ontario, Canada, L8S 4K1

Department of Applied Mathematics, Nizhny Novgorod State Technical University, 24 Minin Street, Nizhny Novgorod, Russia

Discrete breathers are ubiquitous structures in nonlinear anharmonic models ranging from the prototypical example of the Fermi-Pasta-Ulam model to Klein-Gordon nonlinear lattices, among many others. We propose a *general* criterion for the emergence of instabilities of discrete breathers analogous to the well-established Vakhitov-Kolokolov criterion for solitary waves. The criterion involves the change of monotonicity of the discrete breather's energy as a function of the breather frequency. Our analysis suggests and numerical results corroborate that breathers with increasing (decreasing) energy-frequency dependence are generically unstable in soft (hard) nonlinear potentials.

Introduction. Discrete breathers, also referred to as intrinsic localized modes, are time-periodic and exponentially localized in space coherent structures that have been extensively studied over the last three decades; see, e.g., [1, 2]. Their relevance has been recognized not only theoretically but, importantly, via physical experiments in areas as diverse as Josephson junction arrays [3, 4], micro-mechanical cantilever arrays [5, 6], coupled antiferromagnetic layers [7], electrical transmission lines [8], halide-bridged transition metal complexes [9], and torsionally-coupled pendula [10] among numerous others. Remarkably, their areas of purview continue to grow with a recent example being, e.g., granular crystals in material science [11, 12]. Essentially, it is recognized that broad classes of nonlinear dynamical lattices, including the paradigmatic (for nonlinear science) case of the Fermi-Pasta-Ulam (FPU) problem [13, 14], as well as that of Klein-Gordon (KG) chains support a plethora of such states.

Since the energy function is typically the only conserved quantity for the FPU and KG chains, stability criteria that are well-established for solitary waves, such as the famous Vakhitov–Kolokolov (VK) slope condition [15], *do not apply* to classify their stability. As a result, most studies of stability of discrete breathers chiefly rely on numerical experiments and a qualitative analysis of eigenvalues in the Floquet-Bloch spectra of the time-periodic linearization operators [16–19]. Some analytical results on the stability of discrete breathers for KG lattices were obtained by using the limit of small coupling between nearest lattice sites, typically referred to as the anti-continuum (AC) limit [20]. In this limit, asymptotic stability of the fundamental (single-site) breathers was established in [21]. Spectral stability of excited (multi-site) breathers was classified near the AC limit in the work of [22–24], depending on the phase difference in the nonlinear oscillations between different sites of the lattice. More recently, nonlinear instability of spectrally stable two-site breathers was shown in [25]. Nevertheless, an overarching criterion of breather stability tantamount to the VK criterion remains un-

known up to now.

In this work, we fill in this important void by deriving a universal energy criterion *both* for the KG and FPU lattices. In particular, we show that a transition from stability to instability of a discrete breather will occur at frequency ω , where the *energy-frequency dependence features an extremum*, i.e., at $H'(\omega) = 0$, where H is the breather's energy. The previously known lattices that exhibit energy thresholds for discrete breathers like in [26, 27] represent case examples of such an instability transition. Yet, here we illustrate the generality of such a conclusion both through an analytical theory and through a number of prototypical numerical examples (KG, monoatomic FPU, and diatomic FPU). In the vicinity of the bifurcation point, where $H'(\omega) = 0$, our asymptotic analysis and numerical computations suggest the following general conclusion: Breathers with increasing (decreasing) energy-frequency dependence are generically unstable in soft (hard) nonlinear potentials. On the other hand, breathers with decreasing (increasing) energy-frequency dependence in soft (hard) potentials are generally free of the instability associated with this criterion, yet they may experience other instability forms (including e.g. period doublings, oscillatory instabilities, etc. [1, 2]). Let us mention that here, the potential is referred to as hard (soft) when the energy-frequency dependence of individual oscillators is monotonically increasing (decreasing) [28].

Mathematical Setup. We consider a one-dimensional (1D) chain of nonlinear oscillators under Newtonian dynamics:

$$\ddot{u}_n + V'(u_n) = W'(u_{n+1} - u_n) - W'(u_n - u_{n-1}), \quad (1)$$

where n is defined on a 1D lattice, V is an on-site (substrate) potential and W is the inter-site potential for nearest-neighbor interaction. Both V and W are assumed smooth. The associ-




FIG. 1: Breathers in a 2D KG lattice with a hard quartic potential in the case of $C = 0.5$ (left panels) and a Morse potential with $C = 0.2$ (right panels). The top panels show the profile of two unstable breathers with a portion of the unit circle shown in the inset, corresponding to $C = 0.5$, $\omega = 2.3$ (left) and $C = 0.2$, $\omega = 0.992$ (right). Central panels shows the energy-frequency dependence, whereas the bottom panels display the Floquet multipliers with $|\mu| > 1$ (i.e., associated with instability) versus ω .

ated energy function for the lattice (1) is given by

$$H = \sum_{n \in \mathbb{Z}} \frac{1}{2} \dot{u}_n^2 + V(u_n) + W(u_{n+1} - u_n). \quad (2)$$

If $W'(u) = Cu$ with coupling constant C while V satisfies $V'(0) = 0$ and $V''(0) > 0$, the chain is referred to as the Klein–Gordon (KG) lattice. If $V'(u) = 0$ while W satisfies $W'(0) = 0$ and $W''(0) > 0$, the chain is referred to as the Fermi–Pasta–Ulam (FPU) lattice. For clarity, we describe our results for the KG lattice and draw parallels to the FPU case.

Discrete breathers of the KG lattice are T -periodic solutions with $u_n(t+T) = u_n(t)$ for every n . Setting the breather frequency to $\omega = 2\pi/T$, we can normalize the period of the breather to 2π using $u_n(t) = U_n(\tau)$, where $\tau = \omega t$ and $U_n(\tau + 2\pi) = U_n(\tau)$. The profile U_n also depends on frequency ω . We then have

$$\omega^2 U_n''(\tau) + V'(U_n(\tau)) = C(\Delta U)_n(\tau), \quad (3)$$

where $(\Delta U)_n$ denotes the discrete Laplacian. The spectral stability of discrete breathers is determined by the linearized equations of motion

$$\ddot{w}_n + V''(u_n)w_n = C(\Delta w)_n, \quad (4)$$

where w_n is a perturbation to u_n . According to the Floquet theory, we are looking for solutions of the linearized equation (4) in the form $w_n(t) = e^{\lambda t} W_n(\tau)$, where $\lambda \in \mathbb{C}$ is a spectral parameter and $W_n(\tau + 2\pi) = W_n(\tau)$. The spectral stability problem is then

$$\omega^2 W_n''(\tau) + 2\lambda\omega W_n'(\tau) + \lambda^2 W_n(\tau) + V''(U_n(\tau))W_n(\tau) = C(\Delta W)_n(\tau). \quad (5)$$

The (continuous) spectral bands can be identified on the unit circle in terms of the Floquet multipliers $\mu = e^{\lambda T}$. To be precise, the two bands are located at $\mu_{\pm}(\theta) = e^{\pm i\omega(\theta)T}$, where $\omega(\theta) = \sqrt{1 + 4C \sin^2(\frac{\theta}{2})}$, $\theta \in [-\pi, \pi]$. We assume that the two bands are bounded away from the unit multiplier $\mu_0 = 1$, which corresponds to the isolated eigenvalue $\lambda_0 = 0$ in the spectral problem (5). Because of the translational invariance symmetry (in time), we note that the isolated eigenvalue $\lambda_0 = 0$ is at least double. Indeed, the eigenvector $W_n(\tau) = U_n'(\tau)$ satisfies (5) for $\lambda = 0$. Furthermore, the generalized eigenvector $\tilde{W}_n(\tau) = \partial_{\omega} U_n(\tau)$ satisfies the derivative of (5) in λ for $\lambda = 0$ given by

$$(L\partial_{\omega} U)_n(\tau) = 2\omega U_n''(\tau), \quad (6)$$

where

$$(LW)_n(\tau) = C(\Delta W)_n(\tau) - V''(U_n(\tau))W_n(\tau) - \omega^2 W_n''(\tau)$$

is the linearized operator for the spectral problem (5).

Let us assume that the kernel of L is exactly one-dimensional with the eigenvector $W_n(\tau) = U_n'(\tau)$. This assumption is generally satisfied because no other symmetry exists in the lattice (1) besides the translational symmetry in time. The most typical scenario of a discrete breather becoming unstable occurs when a pair of Floquet multipliers μ on the unit circle coalesces at $\mu_0 = 1$ and splits along the real axis. At the critical point, the eigenvalue $\lambda_0 = 0$ of the spectral problem (5) is assumed to have a higher-than-two-algebraic multiplicity. It is exactly that condition which will provide us with the energy criterion for spectral stability of discrete breathers, as follows.

The condition that $\lambda_0 = 0$ is at least quadruple (by Hamiltonian symmetry, it has an even multiplicity) is equivalent to the Fredholm condition of existence of a solution to the second derivative of (5) in λ for $\lambda = 0$. Using the projection technique [28] yields the solvability condition in the form

$$0 = \int_0^{2\pi} \sum_{n \in \mathbb{Z}} U_n'(\tau) [2\omega \partial_{\omega} U_n'(\tau) + U_n'(\tau)] d\tau = TH'(\omega),$$

where $H(\omega)$ is the time-independent breather energy that follows from (2). The higher multiplicity condition (signaling the potential transition between stability and instability) is thus satisfied if ω is a critical point of the breather energy $H(\omega)$.

The solvability condition $H'(\omega) = 0$ cannot be satisfied in the AC limit, where the individual oscillator is always stable with $H'(\omega) > 0$ for hard potentials and $H'(\omega) < 0$ for soft




FIG. 2: Breathers in a monoatomic FPU chain with $\alpha = -1$, $\beta = 1$. Left (right) panels corresponds to the Sievers–Takeno (Page) mode. The top panels show the breather profiles, in the strain variable, for $\omega = 2.1$. The middle panel shows the energy-frequency dependence, whereas the bottom panel displays modulus of the Floquet multipliers with $|\mu| > 1$ versus ω .

potentials [28]. However, far from the AC limit such a bifurcation may (and often does) occur. If at the critical point, $\lambda_0 = 0$ is exactly quadruple, i.e., if a pair of simple Floquet multipliers coalesces with the double unit multiplier $\mu_0 = 1$ at $H'(\omega) = 0$, then an expansion of the eigenvalue problem (5) near the bifurcation point yields:

$$\lambda^2 T H'(\omega) + \lambda^4 M + \mathcal{O}(\lambda^6) = 0, \quad (7)$$

where $M \neq 0$. Then, if $M > 0$, the breathers are stable if $H'(\omega) > 0$ and unstable if $H'(\omega) < 0$, whereas if $M < 0$, then the breathers are stable if $H'(\omega) < 0$ and unstable if $H'(\omega) > 0$. Detailed asymptotic analysis [28] suggests that the former case is intrinsic for hard potentials and the latter case is typical for soft potentials, at least in the small-amplitude limit of KG breathers.

The same conclusion is also drawn in the FPU case when reformulated in terms of the strain variable $r_n = u_{n+1} - u_n$, because it is the strain variable that decays to zero at infinity for FPU breathers [28].

Numerical illustrations: 2D KG breathers. We consider a two-dimensional (2D) version of the KG lattice with the hard ϕ^4 potential $V(u) = u^2/2 + u^4/4$ [6] and the soft Morse potential $V(u) = (\exp(-u) - 1)^2/2$. The latter has been ubiquitously utilized for the study of breathers in DNA denaturation settings where it is used to model the hydrogen bond




FIG. 3: Gap breathers in a diatomic FPU chain for a *hard* potential with $\alpha = -1$, $\beta = 1$, $\epsilon = 0.8$ (left) and a *soft* potential with $\alpha = 0$, $\beta = -1$ and $\epsilon = 0.7$ (right). The top panels show the breather profiles, in the strain variable, for $\omega = 1.7$ (left) and $\omega = 1.4$ (right). Blue (red) dots correspond to the more (less) massive particles. The middle panel shows the energy-frequency dependence, whereas the bottom panel displays the modulus of the Floquet multipliers with $|\mu| > 1$ versus ω .

connecting the two bases in a pair [29].

Fig. 1 shows the energy-frequency dependence for a fixed coupling constant C , as well as the most unstable real Floquet multiplier (recall that instability is tantamount to $|\mu| > 1$) for both hard and soft potentials. We observe a perfect correlation, as prescribed by the theory, between the stability changes and energy extrema. Indeed, the breather is stable (unstable) at the regions of increasing (decreasing) energy $H(\omega)$ for hard potentials, and this trend is reversed for soft potentials.

Notice that in the case of the hard potential, the breather is still stable for every ω past the upper limit shown in Fig. 1. However, in the case of the Morse potential, an instability emerges for ω below the lower limit of the figure. This instability not predicted by our energy criterion pertains to the exchange of instability (precursor of breather mobility) that typically occurs within the Morse potential [30].

Numerical Illustrations: 1D FPU breathers. We consider both monoatomic and diatomic FPU chains [14]. In general, these chains are modeled by the FPU equation

$$M_n \ddot{u}_n = W'(u_{n+1} - u_n) - W'(u_n - u_{n-1}), \quad (8)$$

with M_n being the particle masses. We choose $V(u) = u^2/2 + \alpha u^3/3 + \beta u^4/4$. In the monoatomic case, $M_n = 1$

for all sites, whereas, in the diatomic case, $M_n = 1$ for n even and $M_n = 1/\epsilon^2$ for n odd, where ϵ^2 is the parameter for mass ratio of the diatomic FPU chain [18, 19].

It was demonstrated in [27], for the *monoatomic chain*, that the large-amplitude breathers possess a minimum of $H(\omega)$ since their amplitude does not tend to zero at the band edge $\omega \rightarrow 2$. The energy threshold exists when α is taken below a critical value of $\alpha_c = -\sqrt{3}/2 \approx -0.86$ (for $\beta = 1$). However, in [27], the instability past the energy minimum was not considered. Here we show that the energy threshold results in the change of stability of discrete breathers.

As is typically the case in both FPU and KG chains, there are two principal breathers, the so called Sievers–Takeno (bond-centered) and Page (site-centered) modes. The former is, in general, exponentially unstable. Fig. 2 shows, as dictated by our stability criterion for hard potentials, that an exponential instability arises at the energy minimum for *both* modes when $\omega \rightarrow 2$. In the Page mode, this transition manifests itself as the appearance of an exponential instability of the previously stable structure. In the already unstable Sievers–Takeno mode, a second unstable Floquet multiplier appears as $\omega \rightarrow 2$ (for a secondary instability which rapidly overtakes the previous one as the instability with the largest growth rates).

In the *diatomic case*, there is an opening of a frequency gap within the phonon spectrum,

$$2\epsilon^2 W''(0) < \omega^2 < 2W''(0).$$

This allows the existence of breathers with frequency ω in the gap of the phonon spectrum (so-called *gap breathers*). Such structures can exist even in the case of soft potentials [31], bifurcating from the bottom of the optical phonon band; see also [11] for a relevant experimental manifestation of such modes. For the soft potential, see the right panels on Fig. 3, no global energy minimum exists but extrema in the energy-frequency curve may occur even if $\alpha = 0$. In a full agreement

with the energy criterion for soft potentials, the instability of such gap breathers is perfectly correlated with the increasing energy-frequency dependence,

Finally, gap breathers also exist for hard potentials, bifurcating from the top of the acoustic band, see the left panels on Fig. 3. Their stability and energetic properties are similar to the breathers in the monoatomic FPU lattice, also necessitating a non-zero α for the existence of energy minima.

Conclusions. In this work we have presented a systematic and general energy criterion for spectral stability of breathers in nonlinear dynamical lattices. The energy stability criterion for discrete breathers is strongly reminiscent of the VK criterion for solitary waves; in fact, as illustrated in [28], it *reduces* to the VK criterion in the small amplitude limit where the breathers can be approximated as solitary waves. In view of that, the proposed criterion can be considered as the definitive analogue of the VK criterion for breathers.

We have then corroborated the validity of the energy criterion for stability of discrete breathers via a wide range of models, both KG and FPU, both 1D and 2D, both homogeneous and heterogeneous, showcasing that its generality transcends the specific such properties of the model. It follows from our numerical results that the breathers are unstable in hard (soft) potentials if the energy-frequency dependence is decreasing (increasing) and stable otherwise.

Admittedly, a general classification of instabilities of breathers (more generally of periodic orbits, including non-localized ones, such as plane waves in Hamiltonian systems) in the same spirit as the well developed theory of solitary waves of the nonlinear Schrödinger equation is still incomplete. Nevertheless, the present criterion we believe, constitutes an important step towards future work in this direction, and on understanding nonlinear stability of breathers in lattices.

-
- [1] S. Aubry, *Physica D* **103**, 201 (1997).
 [2] S. Flach and A.V. Gorbach, *Phys. Rep.* **467**, 1 (2008).
 [3] E. Trías, J.J. Mazo and T.P. Orlando, *Phys. Rev. Lett.* **84**, 741 (2000).
 [4] P. Binder, D. Abramov, A.V. Ustinov, S. Flach and Y. Zolotaryuk, *Phys. Rev. Lett.* **84**, 745 (2000).
 [5] M. Sato, B.E. Hubbard, A.J. Sievers, B. Ilic, D.A. Czaplewski and H.G. Craighead, *Phys. Rev. Lett.* **90**, 044102 (2003).
 [6] M. Sato, B.E. Hubbard, L.Q. English, A.J. Sievers, B. Ilic, D.A. Czaplewski and H.G. Craighead, *Chaos* **13**, 702 (2003).
 [7] U.T. Schwarz, L.Q. English and A.J. Sievers, *Phys. Rev. Lett.* **83**, 223 (1999).
 [8] L.Q. English, R. Basu Thakur and R. Stearrett, *Phys. Rev. E* **77**, 066601 (2008).
 [9] B. I. Swanson, J. A. Brozik, S. P. Love, G. F. Strouse, A. P. Shreve, A. R. Bishop, W.-Z. Wang, and M. I. Salkola, *Phys. Rev. Lett.* **82**, 3288 (1999).
 [10] J. Cuevas, L. Q. English, P. G. Kevrekidis, and M. Anderson *Phys. Rev. Lett.* **102**, 224101 (2009).
 [11] N. Boechler, G. Theocharis, S. Job, P.G. Kevrekidis, M.A. Porter and C. Daraio, *Phys. Rev. Lett.* **104**, 244302 (2010).
 [12] C. Chong, F. Li, J. Yang, M. O. Williams, I. G. Kevrekidis, P. G. Kevrekidis, and C. Daraio *Phys. Rev. E* **89**, 032924 (2014).
 [13] E. Fermi, J. Pasta, and S. Ulam, *Tech. Rep. Los Alamos Nat. Lab. LA1940* (1955).
 [14] D. K. Campbell, P. Rosenau, and G. M. Zaslavsky, *Chaos* **15**, 015101 (2005).
 [15] N.G. Vakhitov and A.A. Kolokolov, *Radiophys. Quantum Electron.* **16** 783 (1973).
 [16] R.S. MacKay and J.-A. Sepulchre, *Physica D* **119**, 148 (1998).
 [17] J.L. Marín, S. Aubry, and L.M. Floría, *Physica D* **113**, 283 (1998).
 [18] T. Cretegny, R. Livi and M. Spicci, *Physica D* **119**, 88 (1998).
 [19] G. James and P. Noble, *Physica D* **196**, 124 (2004).
 [20] R.S. MacKay and S. Aubry, *Nonlinearity* **7**, 1623 (1994).
 [21] D. Bambusi, *Comm. Math. Phys.* **324**, 515 (2013).
 [22] J.F.R. Archilla, J. Cuevas, B. Sánchez-Rey, and A. Álvarez,

- Physica D **180**, 235 (2003).
- [23] V. Koukoulouyannis and P.G. Kevrekidis, *Nonlinearity* **22**, 2269 (2009).
- [24] D.E. Pelinovsky and A. Sakovich, *Nonlinearity* **25**, 3423 (2012).
- [25] J. Cuevas–Maraver, P.G. Kevrekidis, and D.E. Pelinovsky, DOI: 10.1111/sapm.12107. *Stud. Appl. Math.* (2015).
- [26] S. Flach, K. Kladko and R.S. MacKay, *Phys. Rev. Lett.* **78**, 1207 (1997).
- [27] B. Sánchez–Rey, G. James, J. Cuevas and J.F.R. Archilla, *Phys. Rev. B* **70**, 014301 (2004).
- [28] See the Supplementary Material at [url], which includes Refs. [32], [33], [34], [35], [36], [37] and [38], for (i) the derivation of expansion (7), (ii) an analysis of small-amplitude Klein-Gordon breathers, (iii) an extension of the result to FPU lattices, (iv) an energy criterion in the anti-continuous limit, and (v) a brief description of the numerical methods used for calculating discrete breathers.
- [29] M. Peyrard and A.R. Bishop, *Phys. Rev. Lett.* **62**, 2755 (1989).
- [30] S. Aubry and T. Cretegny, *Physica D* **119**, 34 (1998).
- [31] G. James and M. Kastner, *Nonlinearity* **20**, 631 (2007).
- [32] D. Pelinovsky, T. Penati, and S. Paleari, arXiv:1509.06389.
- [33] G. James, *J. Nonlin. Sci.* **13** (2003), 27–63.
- [34] G. James, B. Sánchez-Rey, and J. Cuevas, *Rev. Math. Phys.* **21** (2009), 1–59.
- [35] W.-X. Qin and X. Xiao, *Nonlinearity* **20** (2007), 2305–2317.
- [36] D.E. Pelinovsky and G. Schneider, arXiv:1603.05463.
- [37] J. Cuevas, J.F.R. Archilla, and F.R. Romero, *J. Phys. A: Math. Theor.* **44**, 035102 (2011).
- [38] J.F.R. Archilla, R.S. MacKay, and J.L. Marín, *Physica D* **134**, 406 (1999).